

DETERMINATION OF PHYSICAL AND BIOMETRIC PROPERTIES OF ONION BULBS IN RELATION TO DESIGN OF DIGGER CUM WINDROWER

SHIDDANAGOUDA YADACHI* AND KIRAN NAGAJJANAVAR

Department of Agril. Engineering, CHEFT Devihosur, Haveri - 581 110, Karnataka

e-mail: shiddu86@gmail.com

KEYWORDS

Shape index
AMD
GMD
Bulk density

Received on :
26.06.2020

Accepted on :
13.08.2020

*Corresponding
author

ABSTRACT

The physical and biometric properties play vital role in design of any soil operating machine. In view of designing the various components of digger-cum-windrower, physical and biometric properties were determined for three varieties of onion bulbs. The critical dimensions, viz. equatorial diameter, polar diameter, thickness, geometric mean diameter, arithmetic mean diameter, cross sectional area, shape index, bulk density, true density, weight of bulb, angle of repose were determined. Geometric and arithmetic mean diameters were identical for all three varieties, which could help designing soil separator effectively. The Arka Bhim was observed to be denser than that of two other varieties. The maximum cross sectional area was recorded for both Arka Kalyan and N 53 ($1.74 \pm 0.05 \text{ cm}^2$ and $1.75 \pm 0.07 \text{ cm}^2$) against Arka Bhim ($1.50 \pm 0.04 \text{ cm}^2$). Highest values of true density and bulk density were recorded for Arka Bhim. Shape index was under 1.5 for all the three varieties and bulb was considered to be nearly spherical. The Arka Bhim with its maximum frictional coefficient experienced less flow behavior. The measured properties as a whole helped optimizing design values of different components and effective fabrication of prototype.

INTRODUCTION

Onion (*Allium Cepa* L.) is an important vegetable crop among the root vegetable crops in all countries. India is the second largest producer of onion after China with the production of 24.4 million tonnes in an area of 1.3 mha during 2018-19. Onion crop has attained major significance of cash crop rather than a vegetable crop as it has high export potential. This is the most unique vegetable consumed by almost sections of society throughout the year. Onion is extensively cultivated in the North-Karnataka districts, viz. Gadag, Dharwad, Belgaum, Bagalkot and Vijayapur. The onion root system is fibrous, spreading just beneath the soil surface to a distance of 45 cm (Al-Jamal *et al.*, 2001). Onions play a significant role in preventing heart diseases and other ailments (Augusti, 1990).

Of late, farmers cultivate underground vegetable crops over the raised beds in order to save the irrigation water, energy and harvesting of roots with minimum damage. The underground crops, viz. potato, onion, turmeric and ginger are the crops which are planted on the raised beds to facilitate healthier root growth, development and better productivity as well. Manual harvesting of root, tuberous and rhizomatous crops is uneconomical, laborious and time consuming one. Harvesting is a labour oriented operation, which accounts to 21.4 per cent of total production cost (Jadhav *et al.*, 1995). Timely harvesting is very essential in order to get higher market price and harvesting bulbs with less physical damage. The delay in harvesting may affect the keeping quality of onion bulbs. After maturity, onion bulbs are harvested with tractor operated diggers-cum-harvesters or manually after loosening

the soil. Onion diggers are available, however, the dug onions are supposed to be picked by manual labours as the existing diggers leave the onions submerged inside the soil. To avoid onion submerging in soil during digging, there is a need to develop windrower mechanism for digger. Design of any machine or equipment which interacts with soil and crops simultaneously requires the knowledge of physical and biometric dimensions (Somashekar and Nagesha 2010). The physical, frictional, biometric and engineering properties of bulb decides the design values of various functional parts of onion digger-cum-windrower. Availability of these properties is limited for onion bulb and also physical properties of different varieties of onion crop very essential. Some researchers have studied physical, engineering and mechanical properties of onion bulbs from the view point of design of onion diggers. The physical properties of any crop play a key role in designing agricultural processing machinery with higher standards and properly (Tabatabaefar and Rajabipour, 2005; Karababa, E. 2006). Bahnasawy *et al.* (2004) and Rani *et al.* (2006) studied biometric properties of different cultivars of onion. These properties included shape index, geometric mean diameter, arithmetic mean diameter, frontal surface area, cross sectional area, volume, unit mass, density, angle of repose, rolling angle, crushing load and puncture resistance. Bahnasawy (2007) and Arunkumar *et al.* (2014) also studied some physical and mechanical properties of garlic, viz. linear dimensions, shape index and surface area, friction angle, coefficient of static friction etc. for garlic. Abdel-Ghaffar and Hindey (1984) tested Egyptian onion of four different sizes (small, medium, large, and extra-large). They found that the

mean polar diameters were 40.45, 47, 47.94, and 52.40 mm and the mean equatorial diameters were 39.07, 50.03, 56 and 60.40 mm for the same previous order. The mean mass, bulb density and bulk density were 177 g, 0.976 g/cm³, and 0.586 g/cm³, respectively. There was ample demand for growing of cultivars like Arka Kalyan, Arka Bhim and N-53 in North Karnataka and borders districts of Maharashtra. Literature related to the properties, viz. Arka Kalyan, Arka Bhim and N-53 is limited. Hence, the study was planned to determine physical, frictional and biometric properties of onion bulbs of Arka Kalyan, Arka Bhim and N-53, in order to attain the design values of different components of digger-cum-windrower.

MATERIALS AND METHODS

The study was conducted in the food engineering laboratory of College of Horticultural Engineering and Food Technology, UHS Bagalkote, Karnataka during 2019-20. Three varieties of onion extensively grown in different parts of Karnataka state were selected for the study (Fig.1). Ten onion bulbs were randomly chosen from the lot of fresh bulbs for the measuring properties. The physical properties, viz. equatorial diameter (De), polar diameter (Dp), thickness(t), geometric mean diameter (Dgm), arithmetic mean diameter (Dam), cross sectional area (Acs), shape index, bulk density, true density, weight of bulb, angle of repose were determined (Fig 2). In order to provide aeration, onion bulbs were stored in jute bag.

Moisture content

The moisture content of freshly harvested onion bulbs were determined by oven method. Bulbs were dried in an air ventilated oven at 105°C for 48h and the observations related to weights were taken on an electronic balance of precision of 0.01 g. Moisture content was calculated in wet basis. (AOAC, 1990).

Physical, frictional and biometric properties

Twenty onion bulbs were randomly selected from the lot of each variety and their linear dimensions such as polar diameter, equatorial diameter and transverse diameter were measured. The vernier caliper (Mitutoyo 530-312) of ±0.01

mm accuracy was used for measurement of linear dimensions. Polar diameter (Dp) is the distance between the onion crown and the point of root attachment to the onion bulb (Fig.2). Equatorial diameter (De) is the largest diameter perpendicular to the polar diameter and transverse diameter (Dt) is the smallest diameter perpendicular to polar diameter.

Mean diameters and cross sectional area

The geometric mean diameter (Dgm), arithmetic mean diameter (Dam) and cross sectional area (Acs) of onion bulb were calculated using the following equations;

$$D_{gm} = (De \cdot Dp \cdot Dt)^{1/3} \dots\dots\dots(1)$$

$$D_{am} = \left(\frac{De + Dp + Dt}{3} \right) \dots\dots\dots(2)$$

$$A_{cs} = \frac{\pi}{4} \left[\left(\frac{De + Dp + Dt}{3} \right)^2 \right] \dots\dots\dots(3)$$

Where, De–Equatorial diameter, mm; Dp– Polar diameter, mm; Dt –Transverse diameter, mm

Shape index

Shape index is a measure of the shape of onion bulbs and it is computed according to the following equation:

$$\text{Shape index} = \frac{De}{\sqrt{Dp \cdot Dt}} \dots\dots\dots(4)$$

RESULTS AND DISCUSSION

The properties were determined at moisture content of 80.57 ± 0.95 per cent (wb) for the freshly harvested onion bulbs. The descriptive statistics such as mean, standard deviation and CV were computed for all the properties determined in this study. There were no previous works accomplished related to physical properties of these varieties Arka Kalyan, Arka Bhim and N53.

Shape and size

The linear measurements are very crucial in finalizing the

Table 1: Details of physical and engineering properties measured

Variety	Physical and Engineering properties
Arka kalyana	Equatorial diameter, Polar diameter, Thickness, GMD, AMD, Cross sectional area, Shape index, Bulk density, True density, Weight of bulb, Angle of repose
Arka Bhim	
N-53	

Table 2: Properties of onion bulbs to determine shape index

Variety	Descriptive Statistics	Equatorial diameter (De), cm	Polar diameter (Dp), cm	Transverse diameter, cm	Shape index
Arka Kalyan	Mean	7.27	6.42	6.21	1.15
	SD	0.33	0.37	0.3	0.07
	CV, %	4.54	5.81	4.77	6.24
Arka Bhim	Mean	6.47	5.49	5.21	1.21
	SD	0.3	0.31	0.29	0.07
	CV, %	4.6	5.61	5.63	5.93
N 53	Mean	7.27	6.47	6.37	1.13
	SD	0.43	0.35	0.22	0.06
	CV, %	5.9	5.43	3.39	5.57

Table 3 : Physical and engineering properties of onion bulbs

Variety	Descriptive Statistics	GMD (Dgm), cm	AMD (Dam), cm	CS area (Acs), cm ²	Mass of bulb (g)
Arka Kalyan	Mean	6.61	6.63	1.74	157.9
	SD	0.18	0.18	0.05	6.01
	CV, %	2.68	2.67	2.67	3.81
Arka Bhim	Mean	5.68	5.72	1.5	131.6
	SD	0.16	0.16	0.04	6.42
	CV, %	2.82	2.85	2.85	4.87
N 53	Mean	6.67	6.7	1.75	168
	SD	0.27	0.27	0.07	7.39
	CV, %	4.02	4.06	4.06	4.4

**Figure 1: Overview of different varieties of onion bulbs selected for the study****Table 4: Bulk density and true density of onion bulbs**

Variety	Descriptive Statistics	Bulk Density, gcc-1	True density, gcc-1
Arka Kalyan	Mean	0.66	1.01
	SD	0.02	0.08
	CV, %	3.47	7.65
Arka Bhim	Mean	0.67	0.98
	SD	0.03	0.04
	CV, %	5.1	3.88
N 53	Mean	0.71	0.92
	SD	0.05	0.03
	CV, %	6.45	3.37

Table 5 : Coefficient of static friction of onion bulbs for different surfaces

Surface	Coefficient of static friction		
	Arka Kalyan	Arka Bhim	N-53
GI Sheet	0.31	0.33	0.27
Mild steel sheet	0.34	0.29	0.31
Plastic sheet	0.22	0.3	0.29
Plywood	0.27	0.32	0.27

dimensions of length and widths of blade and soil separators. The descriptive statistics of various properties determined for three varieties of onion bulbs were computed (Table 2). The mean values of equatorial, polar and transverse diameter were 7.27 ± 0.33 , 6.42 ± 0.37 and 6.21 ± 0.30 for Arka Kalyan; 6.47 ± 0.30 , 5.49 ± 0.31 and 5.21 ± 0.29 for Arka Bhim and 7.27 ± 0.43 , 6.47 ± 0.35 and 6.37 ± 0.22 for N 53. The coefficients of variation for equatorial, polar and transverse diameter were below 10 per cent under all varieties. The Arka Kalyan and N 53 were recorded higher values as that of Arka Bhim. Shape index was 1.15 ± 0.07 for Arka Kalyan; 1.21 ± 0.07 for Arka Bhim and 1.13 ± 0.06 for N53. From the

observations, it was conceptualized that onion bulbs are spherical in shape. There were no similar studies which compared the physical properties of these varieties of onion. However, with respect to shape, similar results were observed by Bahnasawy *et al.* (2004).


The geometric mean diameter (Dgm) and arithmetic mean diameters (Dam) for all varieties were indifferent. Mean values of Dgm and Dam were 6.61 ± 0.18 cm and 6.63 ± 0.18 cm for Arka Kalyan; 5.68 ± 0.16 cm and 5.72 ± 0.16 cm for Arka Bhim and 6.67 ± 0.27 cm and 6.70 ± 0.27 cm for N 53 (Table 3). The CV values of Dgm and Dam were below 10 per cent for all the varieties. The cross sectional area was 1.74 ± 0.05 cm² for Arka Kalyan, 1.50 ± 0.04 cm² for Arka Bhim and 1.75 ± 0.07 for N 53. Marginally, the higher cross sectional area was recorded for Arka Kalyan and N 53 (1.74 ± 0.05 cm² and 1.75 ± 0.07 cm²) as compared to Arka Bhim (1.50 ± 0.04 cm²). The average mass of bulbs for three varieties Arka Kalyan, Arka Bhim and N 53 were to the tune of 90 ± 6.01 g, 131.60 ± 6.42 g and 168 ± 7.39 g, respectively. Higher mass of bulb was obtained under cultivar N 53 followed by Arka Kalyan and Arka Bhim. The lower CV (less than 5 per cent) values were observed for both mass of bulb and cross sectional area. There were no similar studies which compared the physical properties of these varieties of onion. However, the geometric dimensions of onion bulbs in general were in agreement with the results of Bahnasawy *et al.* (2004).

Bulk density and true density

The bulk and true densities are most important functional properties essential for the design of width of windrowing mechanism. In general, Arka Bhim was denser than other two varieties. However, the higher values of density were obtained for cultivars N 53 and Arka Bhim than Arka Kalyan. The bulk

Table 6 : Design values of different components of digger-cum-separator

S No.	Property	Design values	Dimensions of components to be designed
1	De	6.47 ± 0.35 cm	Length, width and thickness of blade, soil separator
2	Dp	5.49 ± 0.34 cm	
3	T	5.21 ± 0.27 cm	
4	Dgm	5.68 ± 0.20 cm	Shape of blade and length of soil separator
5	Dam	5.72 ± 0.20 cm	
6	Acs	1.75 ± 0.05 cm ²	Soil separator slits/grooves
7	SI	1.21 ± 0.07	
8	BD	0.71 ± 0.03 gcc-1	Width of blade
9	TD	1.01 ± 0.05 gcc-1	
10	W	168 ± 6.61 g	Length of soil separator
11	AR	$21 \pm 0.15^\circ$	Digger blade angle and separator angle

**Figure 2: Measurement of polar diameter and conceptual view**

density of 0.66 ± 0.02 g/cc, 0.67 ± 0.03 g/cc and 0.71 ± 0.05 g/cc was recorded under Arka Kalyan, Arka Bhim and N 53, respectively. The true density was 1.01 ± 0.08 g/cc, 0.98 ± 0.04 g/cc and 0.92 ± 0.03 g/cc for Arka Kalyan, Arka Bhim and N 53, respectively. In general, the results were in accordance with the findings of Bahnasawy *et al.* (2004).

Frictional property

The coefficient of friction is the important parameter to measure the level of friction characteristics. Hence, coefficient of friction was measured for different metal surfaces in order to finalize the material of fabrication for blade and soil separator. The Arka Bhim recorded highest coefficient for GI sheet (0.33) followed by plywood (0.32), plastic sheet (0.30) and mild steel sheet (0.29).

The cultivar N 53 had highest coefficient of friction in mild steel sheet (0.31) subsequent to plastic sheet (0.29) whereas; identical results were acquired for GI sheet and Plywood (0.27). The maximum value was obtained for the frictional surface of mild steel sheet (0.34) followed by GI sheet (0.31), plywood (0.27) and plastic sheet (0.22) for the variety Arka Kalyan (Table 5). There was marginal variation in the values of friction coefficient of bulbs of all varieties and the results were in accordance with Savitha and Naik (2011). This difference was due to variations in the unit weight of bulb and volume. In general, the results were in accordance with the findings of Bahnasawy *et al.* (2004).

Based on the measured properties of onion bulbs, the design values were finalized by considering the deviations (SD) in the values for the design of various components of digger cum separator (Shiddanagouda Yadachi *et al.*, 2013). The design values were in accordance with the relevant measured

properties of bulbs. The 3-D pro-e designing of various components was accomplished as per the design values of different components. The blade dimensions such as inclination, thickness and width were critical in attaining the good performance of digger-cum separator. All the linear dimensions were helpful for the design of blade and length of soil separators (Table 6).

ACKNOWLEDGEMENTS

The author is grateful for the financial assistance provided by University of Horticultural Sciences, Bagalkote under University Grants. The support and encouragements of co-authors and staff of College of Horticulture Engineering and Food Technology are truly acknowledged by the author.

REFERENCES

- Abdel-Ghaffar, E. A. and Hindey, F. I. 1984. Linear air flow resistance of onions. *J. Agr. Res. Tanta Univ.* **10(3)**: 721-735
- Al-Jamal, M.S., Ball, S. and Sammi, T.W. 2001. Comparison of sprinkler, trickle and furrow irrigation efficiencies for onion production. *Agric. Water Mgmt.* **46**: 253-266
- AOAC. 1990. Fruits and fruit products. In: Official Methods of Analysis of the Association of Official Analytical Chemists. Vol. II. Arlington, Virginia (USA), PP.910-928
- Arun Kumar, P. Kiran N., S. V. Patil and Palanimuthu, V. 2014. Effect of moisture on the engineering properties of Cashew Nuts, *International J. Advanced Research in Biological Sciences.* **1(7)**: 36-41
- Augusti, K.T. 1990. Therapeutic and medicinal values of onion and garlic. In onion and allied crops. CRC Press. **3**: 93-108
- Bahnasawy A. H., M.Y.El-Ansary and Sorour, H.M.2004. Physical

- and mechanical properties of some Egyptian onion cultivars. *J. Food Engineering*. **62**: 255-261
- Bahnasawy, A. H. 2007.** Some physical and mechanical properties of garlic. *International J. Food Engineering*. **3(6)**: Art. 7.1-6
- Jadhav, R.V., Turbatmath, P.A. and Gharte, L.V. 1995.** Design development and performance of onion digger windrower. *Agric. Mech. Asia, Africa and Latin America*. **26(3)**: 35-38
- Karababa, E. 2006.** Physical properties of popcorn kernels. *J. Food Eng.* **72**: 100-107.
- Kumar, R. 2014.** Biochemical characterization of *Jatropha (Jatropha curcas L.)* accession in Eastern Himalaya, India. *The Bioscan*. **9(3)**: 1131-1134
- Mohsenin N. N. 1986.** Physical properties of plant and animal materials Gordon and Breach, Sci. Pub., New York. **31(7)**:702-702
- Rani Vijaya and Srivastava, A. P. 2006.** Physical and Mechanical Properties of Onion (*Allium Cepa L.*) Crop Relevant to Mechanical De-topping. *J. Agricultural Engineering*. **43(3)**:83-86.
- Savitha, G. and Naik, G. R. 2011.** Production of biodiesel from *jatropha curcas L.* *Oil. The Bioscan*. **6**: 87-88
- Shiddanagouda Yadachi, Mani Indra, Kalra M.S., Adarshkumar and Sahoo, P. K. 2013.** Development and evaluation of inclined plate metering mechanism for carrot seed. *J. Agricultural Engineering (ISAE)*, **50(2)**:10-16
- Shiddanagouda Yadachi, Mani Indra, Kalra M.S., Lande Satish and Varghese, C.2014.** Effect coatings on physical and engineering properties of carrot (*Daucus carota L.*) seeds in relation to the planter design. *Agricultural Mechanization in Asia, Africa and Latin America*. **45(3)**:
- Somashekar and Nagesha 2010.** An overview of adoption of energy efficient technologies in indian urban households. *The Bioscan*. 1.25-34
- Tabatabaeefar, A. and Rajabipour, A. 2005.** Modeling the mass of apples by geometrical attributes. *Scientia Horticulture*. **105**:373-382



**NATIONAL ENVIRONMENTALISTS
ASSOCIATION**

The National Environmentalists Association is chartered in Ranchi as a nonprofit scientific and educational association of like minded academicians, researchers, scientists from all over the nation for the furtherance and diffusion of knowledge of Life Sciences in general and Environmental Science in particular.

The association not only honours its members but also provides FELLOWSHIP to outstanding contributors to the subject and the society.

Contact :

For Editorial Information

Prof. M. P. Sinha
Vice Chancellor
Sido Kanhu Murmu University
Dumka - 814 110
Jharkhand, INDIA

For information regarding Association :

SECRETARY,
National Environmentalists Association,
D-13, Sai Roofs, 1st Floor,
H. H. Colony,
Ranchi - 834002
Jharkhand, India

E-mails : editor.bioscan@gmail.com
dr.mp.sinha@gmail.com
nat.env.assoc@gmail.com

Cell : 94313-60645; 9572649448

Ph. : 0651-2244071

Website : www.thebioscan.com
: www.neaindia.org

NAAS Rating : 5.26

U.S.A. Office

2827 Videre Dr.,
Wilmington,
DE 19808 We, USA

Type setter Bandana Solutions Facility Management LLP
Published by Aditi Publications, Patliputra, Patna

The Bioscan

An International Quarterly Journal of Life Sciences

ISSN : 0973-7049

Volume 15(3) : 2020

Published as an official organ by

NATIONAL ENVIRONMENTALISTS ASSOCIATION

CONTENTS

Page

A. RESEARCH PAPER

1. Comparison and molecular profiling of Begomovirus infecting chilli (*Capsicum annum*) in gangetic alluvial zone of West Bengal
Uday Bikash Oraono, Lourembam Sanajaoba Singh and Jayanta Tarafdar—275 - 280
2. Assessment of probiotic characteristics of *L.plantarum*
Sravani Kandula And Rita Narayanan—281 - 286
3. Effect of rate and frequency of micronutrient on growth attributes and dry matter yield of Banana Cv. grand naine under south Gujarat condition
Narendra Singh, Sonal Tripathi, Patel V. A., Jaimin Naik and Chauhan Aditi—287 - 290
4. Character association and path analysis for seed yield and its components in Grass pea (*Lathyrus sativus* L.)
Gangishetti Ranjithkumar, Sandip Debnath and Duddukur Rajasekhar—291 - 295
5. Determination of physical and biometric properties of onion bulbs in relation to design of digger cum windrower
Shiddanagouda Yadachi and Kiran Nagajjanavar—297 - 301
6. Polygenic variation for morphological and biochemical traits of brinjal genotypes (*Solanum melongena* L.) and its wild relatives
Nisha Sharma, K. D. Bhutia, Rajesh Kumar, Sita Kumari Prasad, Ankita Debnath and Malay Marut Sharma—303 - 309
7. Effect of different sources of horizontal transmission of bacterial flacherie on Et_{50} for symptom expression and mortality of PM X CSR₂
B. L. Kavyashree., R. N. Bhaskar and C. Doreswamy—311 - 314
8. Comparative biology of *Goniozus nephantidis* (Muesbeck) on *Galleria mellonella* L. and *Corcyra cephalonica* (Stainton)
A. V. Desai, M. R. Siddhapara and N. P. Trivedi—315 - 321
9. Efficiency of ovatide on mass seed production of climbing perch (*Anabas testudineus*, Bloch, 1972) in Nalbari district, Assam
Ankur Rajbongshi, A. Ali, M. Chakravarty, M. Deka, H. Mazumdar, Pranab Kr Das and S. Baishya—323 - 326
10. Study of combining ability and gene action for yield and yield component characters in interspecific hybrids of cotton (*Gossypium hirsutum* L. X *Gossypium barbadense* L.)
S. B. Gohil., M. B. Parmar., M. P. Patel and D. A. Patel—327 - 333
11. Per oral inoculation of *Lysinibacillus sphaericus* with pathogenic microbes on rearing and cocoon parameters of silkworm, *Bombyx mori* L.
H. G. Anusha, R. N. Bhaskar and K. V. Anitharani—335 - 338
12. Effect of fungicides, plant extracts and bioagents on spore germination of *Colletotrichum lindemuthianum* causing field bean anthracnose

The Journal is Currently Abstracted / Indexed in

- Paryavarn Abstract, INDIA
- Indian Science Abstract, INDIA
- Cambridge Science Abstract, U.S.A.
- Zoological Record, U.K.
- Directory of Open Access Journal (DOAJ)
- Chemical Abstract, U. S. A.
- Research BIB
- Indian Science
- Journal Seek
- Scientific Indexing Service (SIS)
- Journal is currently rated by
- Index Copernicus
- Universal Impact Factor
- NAAS

DISCLAIMER

The Publisher and Editors cannot be held responsible for errors or any consequences arising from the use of information in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher/ Association and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher / Association and Editors of the products advertised.

- S. Narasimha Rao, S.L. Bhattiprolu, A. Vijaya Gopal and V. Sekhar— 339-344
13. Evaluation in vitro different fungicides for growth of *Rhizoctonia bataticola* A.M. Kadam, S.S. Chavan and A.H. Kendre — 345- 349
 14. Evaluation of gladiolus varieties for flowering and cut flower traits under indo-gangetic plains Girish,P. M., Anjana Sisodia and Anil K. Singh— 351- 355
 15. Effect of land configuration and different organic sources on growth, yield and quality of carrot under organic farming B. Solanki, A. R. Kaswala, P.K. Dubey and A.P. Italiya— 357- 362
 16. Verification and usability analysis of medium range weather forecast for the Kokrajhar district of lower Brahmaputra valley zone of Assam Kuldip Medhi, Kushal Sarmah, Vinod Upadhyay, Sunil Kumar Paul, Athar N. Islam and Bikash J. Gharphalia— 363 - 370
 17. Canonical root analysis and clustering for characterization and evaluation of aromatic rice germplasm based on morphological characters G. Parimala, Ch. Damodhar Raju, L.V. Subba Rao and K. Uma Maheswari— 371 - 374
 18. Genetic divergence analysis of sesame genotypes (*Sesamum indicum* L.) Dasari Rajitha, T. Srikanth, D. Padmaja and T. Kiran Babu—375 - 379
 19. Nutrient uptake and chemical properties of soil after harvest of baby corn (*Zea mays* L.) as influenced by organic manures and fertilizers D.H. Roopashree , S .Kamal Bai . Nagaraju and S.Raghavendra— 381 - 384
 20. Genotypic response on growth and yield in papaya D. K. Varu.,K. D. Patel and Sandip Makhmale— 385 - 389
 21. Studies on frequency distribution of yield and yield related traits in F₂M₂ generation of sesame (*Sesamum indicum* L.) Rajesh Kumar Kar , Tapash Kumar Mishra and Banshidhar Pradhan — 391 - 395
 22. Correlation and path analysis in cowpea (*Vigna unguiculata* (L.) Walp) R.M. Nagalakshmi, R. Usha Kumari and R. Ananda Kumar—397 - 401
 23. A simple and efficient method for DNA extraction from rabi sorghum [*Sorghum bicolor* (L.) Moench]" S. S. Gadakh, G. D. Khalekar.,U. S. Dalvi, A. A.Kale and P.L.Kulwal— 403 - 406
 24. Determination of economic injury level (EIL) of sugarcane plassey borer *Chilo tumidicostalis* hampson (Lepidoptera: pyralidae) R. K. Nath and D. K. Saikia— 407- 409
 25. Performance of different summer mung (*Vigna radiata* L.) varieties sown at different dates under Manipur valley condition Meghna Gogoi, Jamkhogin Lhungdim, Kamal Kant, Urjashi Bhattacharya and Gauri Mohan — 411 - 414

ZONAL CO-ORDINATORS OF THE ASSOCIATION

- Prof. N. Behera
School of Life Science,
Sambalpur University
- Dr. Nirmal Kumar
ISTAR, Vallabh Vidyanagar,
Anand, Gujarat
- Dr. P. N. Sudha
D. K. M. College for Women,
Vellore
- Prof. S. P. S. Dutta
Dept. of Environmental Science,
Jammu University, Jammu
- Dr. V. Salom Gnana Thanga
Dept. of Env. Scs.,
University of Kerala, Kariavattom
Tiruvananthapuram, Kerala

Publications of the Association

The Bioscan

An International Quarterly Journal of Life Sciences

The Ecoscan

An International Quarterly Journal of
Environmental Sciences

**Both the Journals are
online**

**Both the Journals are
available on Google.com**

Websites of the Journals are
www.theecoscan.in
www.thebioscan.in

FEATURES OF ASSOCIATION

- Association is registered under 80G of I.T.
- Prestigious fellowship of the Association (F. N. E. A.) to academicians of the nation.
- Regular annual conference of national and international levels organized by the Association.
- Young Scientist and Senior Scientist award during the conference of the Association.



_____ 062 & 082

_____ 000 & 000

THE BIOSCAN : SUBSCRIPTION RATES				
		India (Rs.)	SAARC Countries	Other Countries
Individuals	One Year	1,000	2,000(I:C)	US \$200
	Life Member*	10,000		
Institutions	One Year	3,000	6,000(I:C)	US \$400
	Life Member*	30,000		

*Life Member will receive the journal for 15 years while other benefits will continue whole life

THE BIOSCAN : MEMBERSHIP FORM

Please enter my subscription for the above journal for the year / life member.

Name:

Address:

E-mail:

Payment Rs. : by DD / MD in favour of
National Environmentalists Association payable at Ranchi, No. Dated
..... is enclosed.

NOTE: FOR MEMBERSHIP THE ABOVE INFORMATION CAN BE SENT ON SEPARATE SHEET