

EFFECT OF DIFFERENT SOURCES OF HORIZONTAL TRANSMISSION OF BACTERIAL FLACHERIE ON ET₅₀ FOR SYMPTOM EXPRESSION AND MORTALITY OF PM X CSR2

B. L. KAVYASHREE., R. N. BHASKAR AND C. DORESWAMY*

Department of Sericulture, University of Agricultural Sciences, Bengaluru - 560 065, INDIA.

*Agriculture college, KVK, Chamarajanagar.

e-mail: kavyabevinakatti@gmail.com

KEYWORDS

Bacterial flacherie
ET₅₀ for symptom
expression
mortality
horizontal-
transmission

Received on :

13.06.2020

Accepted on :

09.08.2020

*Corresponding
author

ABSTRACT

Flacherie disease of silkworm *Bombyx mori* L. is also called as Thatte disease (Thatte roga), Benkiroga (Dhadevu roga) and it was first observed from a village in Malavalli taluk of Mandya district during summer months of 1990 (Doreswamy *et al.*, 2001) cause flaccidity in larva, during which they become feeble, lethargic, vomit gut juice and extrude soft faeces with higher water content. Administration of 10⁻⁵ and 10⁻⁷ dilutions of horizontal sources of inoculum to fourth and fifth instar larvae exhibited significant results. It is very clearly indicated that, decreased bacterial dilution exhibited minimum ET₅₀ value for symptom expression observed for contaminated faecal pellet (9.01, 9.18; 7.01, 7.35 days) for fourth and fifth instar at 10⁻⁵ and 10⁻⁷, respectively followed by contaminated bed (9.09, 9.38; 7.18, 7.30 days) and (10.17, 10.25; 8.12, 8.30 days). The trend was found same in ET₅₀ for larval mortality (9.97, 10.12; 8.01, 8.13 days). It was also concluded that, the horizontal transmission was found minimum in contaminated rearing equipment (9.42, 9.72; 11.03, 11.65 days) and (7.51, 7.67; 9.07, 9.72 days) used for rearing compared to other sources of horizontal transmission. Which in turn recorded maximum number of days to express the disease in both the instars of PM x CSR2.

INTRODUCTION

Flacherie disease of silkworm is caused by different species of bacteria and viruses, individually or in combination. It is generally spread in rearing house by different means. Based on this hypothesis an effort has been made to know the different horizontal transmission which were found very common in the farmers rearing site. Selvakumar (2013) revealed that, the prevalence of both bacterial and viral flacherie were more during summer (5.0-20.00 %) followed by rainy (2.5- 15.00 %) and winter (0.00-7.50 %) seasons. Among several infectious diseases of silkworm, flacherie is known to cause huge cocoon crop loss of about 27 – 35 per cent with decrease in yield of cocoons to an extent of 11- 15 kg per 100DFL's (Selvakumar and Savithri, 2012).

Adverse environmental conditions such as high temperature and humidity and starvation, spacing in the rearing bed (Samson *et al.*, 1981) are considered to be the most important pre-disposing factors for the spread of the disease within the bed and also in the rearing house. Further (Swathi, 2015) has studied the effect of different types of rearing structures on incidence of flacherie disease in silkworm. According to her, the highest incidence of flacherie disease was noticed in Ramanagara district during summer season in RCC house was found to be 20.00 per cent followed by winter 12.00 and rainy 10.00 per cent. The same trend was observed in case of Chikkaballapur district. The highest incidence of flacherie disease was noticed due to unhygienic conditions followed

in and around the rearing house.

Further (Samson, 1995) made observation on age of the silkworm, overcrowding, scanty feed, contaminated leaves, cross infectivity, etc are considered to be pre-disposing factors for the spread of the flacherie in the rearing bed. These factors not only weaken the worms and prone to many microbial infections. Particularly, bacterial flacherie coupled with viral infections are very frequent in silkworm rearing houses. These pre-disposing factors will influence on the multiplication and development of diseases during silkworm rearing (Balavenkatasubbaiah *et al.*, 2006). Based on this objective different sources of horizontal transmission were administered to fourth and fifth instar larvae of PM x CSR2 to know the effective means of disease transmission in the rearing bed.

MATERIALS AND METHODS

A survey was undertaken during the month of August, 2015 in Mallur village of Sidlaghatta taluk, Chikkaballapur district. The sources of inoculum were randomly selected from five sericulture farmers' in their commercial silkworm rearing houses as per the survey conducted on bacterial septecimia by Ashok Kumar and Ramakrishna (2013). They isolated and characterized the septicaemia causing bacterial species in silkworm rearing environment (soil, phylloplane, diseased silkworm and rearing house-silkworm rearing trays and culture from walls). The same procedure was adopted in collection of

samples which are served as effective tool of transmission (contaminated food, bed, rearing equipment, body surface, faecal pellet and floor area) and used for inoculation for fourth and fifth instar silkworm larvae. All the sources of inoculum were collected with 9:1 proportion later subjected for 3000 rpm for 10 min followed by 5000 rpm for 5 min. The filtrate slowly decanted to conical flask.

All glasswares were sterilized in a hot air oven at 18° C for three hours. All growth medium and broth were sterilized in an autoclave at 15 lbs pressure for 20 min. Isolation, purification, inoculation and other microbiological works were carried out in laminar airflow chamber (Robert Pollock *et al.*, 2002).

Inoculation of silkworms

Inoculation of silkworms was done on the fourth instar first day, fifth instar first day *i.e.*, immediately after third and fourth moult, respectively. The spore dilution of 10^{-5} and 10^{-7} of different sources of inoculum were swabbed on mulberry leaf (10 x 15 sq. cm area) using sterilized cotton swab, air dried, made into small pieces and fed to the silkworms at the rate of 0.5ml per 50 worms (Anusha, 2015). Usually for all per oral infection Koch postulatue technique was adopted. The same methodology was used in inoculating fourth and fifth instar larvae. Therefore, the common mode of horizontal transmission was assessed based on sources of inoculum (contaminated food, bed, rearing equipment, body surface, faecal pellet and floor area) (Pasteur, 1870).

The results obtained in the present study are statistically analysed through complete randomized design and conclusions were drawn based on the observations recorded (Sundar Raj *et al.*, 1979).

RESULTS AND DISCUSSION

After administration of different sources of inoculum, the infected larvae (4th and 5th instar) exhibited following symptoms *viz.*, the larvae become weak, vomiting yellow body fluid followed by the anterior portion of the infected larval midgut bulged in between thorax and abdomen made the larval skin tapers, the flow of blood in the body of the silkworm was more compared to uninoculated batches and started wriggling in the bed in the form of 'C' shaped larvae. While wriggling the

entire larval midgut contents were pushed back as a result worms were unable to pass excreta due to sealing of anal flap. Before death, the corpse body become hard and after the death the larval body start become blackening posterior to anterior. After 24 hours it has become fragile, hang down oozing of brown colour fluid from the body and emits foul smell (Plate 1).

Paramasiva and Rajendra Prasad (2009) reported that, insects infected with pathogenic bacteria exhibit symptoms such as loss of appetite, diarrhoea, vomiting, larvae softening and foul odour upon death. Cocoons spun by the infected worms did not exhibit any external symptoms. The size of the cocoon was drastically reduced, compared to normal cocoons. When cocoons were cut open, the pupae were found dead or malformed. The infected cocoons were lighter in weight and moth emergence could not be seen. Infected moths became sluggish with slightly crinkled wings and showed less interest in copulation. The same type of symptoms were observed in both fourth and fifth instar larvae of PM x CSR2 in the present investigation.

The fourth instar inoculated batch of PM x CSR2 with different sources of horizontal transmission caused variation in ET₅₀ for symptom expression and larval mortality. As per the data, the lesser number of days of 9.01, 9.18 days and 9.97, 10.12 days (contaminated faecal pellet) was noticed for 10-5 and 10-7 inoculum administered batches, respectively. However, the increased trend of symptom expression and larval mortality was noticed from contaminated bed (9.09, 9.38 and 10.17, 10.25 days) followed by contaminated food (9.17, 9.59 and 10.31, 10.40 days), contaminated body surface (9.30, 9.68 and 10.97, 11.54 days), contaminated rearing equipment (9.42, 9.72 and 11.03, 11.65 days) and contaminated floor area (9.51, 9.84 and 11.25, 11.84 days) in both the inoculated batches (Table 1).

The fifth instar inoculated batch of PM x CSR2 with different sources of inoculum caused variation in ET₅₀ for symptom expression and larval mortality. As per the data, the minimum days of 7.01, 7.35 days and 8.01, 8.13 days (contaminated faecal pellet) was noticed for 10-5 and 10-7 inoculum administered batches, respectively. The remaining treatments are in increasing order followed by contaminated bed (7.18, 7.30; 8.12, 8.30days), contaminated food(7.30, 7.42; 8.25,

Table 1: Effect of different sources of horizontal transmission of bacterial flacherie on rearing parameters of silkworm, *Bombyx mori* L. (4th instar inoculated batch, PM x CSR2)

Treatments	ET ₅₀ for symptom expression(days)		ET ₅₀ for larval mortality (days)	
	10 ⁻⁵	10 ⁻⁷	10 ⁻⁵	10 ⁻⁷
T ₁ - Contaminated food	9.17	9.59	10.31	10.4
T ₂ - Contaminated bed	9.09	9.38	10.17	10.25
T ₃ - Contaminated rearing equipment	9.42	9.72	11.03	11.65
T ₄ - Contaminated body surface	9.3	9.68	10.97	11.54
T ₅ - Contaminated faecal pellet	9.01	9.18	9.97	10.12
T ₆ - Contaminated floor area	9.51	9.84	11.25	11.84
T ₇ - Distilled water	-	-	-	-
T ₈ - Uninoculated	-	-	-	-
'F' test	*	*	*	*
SEm ±	0.004	0.005	0.01	0.009
CD at 5 %	0.013	0.014	0.031	0.026

* Significant

Table 2 : Effect of different sources of horizontal transmission of bacterial flacherie on ET₅₀ for symptom expression and ET₅₀ for larval mortality of silkworm *Bombyx mori* L. (5th instar inoculated batch)

Treatments	ET ₅₀ for symptom expression (days)		ET ₅₀ for larval mortality (days)	
	10 ⁻⁵	10 ⁻⁷	10 ⁻⁵	10 ⁻⁷
T ₁ - Contaminated food	7.3	7.42	8.25	8.43
T ₂ - Contaminated bed	7.18	7.3	8.12	8.3
T ₃ - Contaminated rearing equipment	7.51	7.67	9.07	9.72
T ₄ - Contaminated body surface	7.43	7.59	8.39	8.7
T ₅ - Contaminated faecal pellet	7.01	7.35	8.01	8.13
T ₆ - Contaminated floor area	7.63	7.76	9.39	9.96
T ₇ - Distilled water	-	-	-	-
T ₈ - Uninoculated	-	-	-	-
F ₆ test	*	*	*	*
SEm ±	0.004	0.097	0.008	0.005
CD at 5 %	0.013	0.29	0.025	0.015

* Significant



Larvae showing symptoms of bacterial flacherie



After death larvae turns into black colour



Hanging dead larvae

Plate1: Symptoms of bacterial flacherie

8.43 days). The remaining sources of inoculum contaminated body surface (7.43, 7.59; 8.39, 8.70days) and contaminated floor area(7.63, 7.76; 9.39, 9.96 days) of ET₅₀ for symptom expression and larval mortality recorded in fifty instar larval batch (Table 2).

This experimental data supported by Anusha and Bhaskar (2016) reported that, when third and fourth instar larvae inoculated with *Bacillus* sp. as surface inoculation recorded decreased larval weight from 1.95 to 13.40 g/10 in PM and 6.38 to 22 g/10 in CSR2 at end of both the instars registering decrease in their development during bacillus infection as observed in the present experiment.

The same observations were confirmed with Siromani *et al.* (1994), bacterial flacherie had comparatively a long period of lethal infection of about 7 to 14 days. Death occurred within 10 to 24 hour in case of bacterial flacherie. Further, Anitha *et al.* (1994) further reported that, when the silkworm larvae were

fed with 106 cells of *Bacillus*, *Staphylococcus* and *Serratia*, no mortality was observed till 48 h. when 107 and 108 cells were fed, mortality was observed within 48 h, which was 100 per cent in *Bacillus* and *Staphylococcus* infections and 80 per cent in *Serratia* infection. The LD50 was 5.9 x 10⁷ for *Bacillus* and *Staphylococcus* infections and 2.49 x 10⁷ in *Serratia* infection as revealed in ET₅₀ values of fourth and fifth instar larvae of PM x CSR2 in the present study.

ET₅₀ for symptom expression value reduced significantly in *Streptococcus faecalis* (166.33h), *Staphylococcus aureus* (171.25h) in the mean of six dilutions (stock, 10⁻², 10⁻⁴, 10⁻⁸, 10⁻¹⁶ and 10⁻³²). The values increased with increase in bacterial dilutions. The ET₅₀ value for larval mortality was observed in *Bacillus* sp. (186.00h) and *Staphylococcus aureus* for stock, 10⁻², 10⁻⁴, 10⁻⁸ and mean of dilutions. Death occurred within 10 to 24 h during bacterial flacherie (Doreswamy *et al.*, 2001).

It was further supported by Chitra *et al.* (1973) the infection

with several bacteria caused severe mortality and majority of the deaths occurred during moult and when the worms were about to spin. Mortality was maximum in fifth instar irrespective of the age of the silkworm. The per cent mortality due to different bacterial species was found to be 37.50 (*Achromobacter superficialis*) to 65.00 (*Achromobacter delmarvae*, *Pseudomonas ovalis*, *P. boreopolis*) per cent on eighth day of fifth instar. As it is revealed in the present experiment.

REFERENCES

- Anitha., T. Shiromani, P. Meena and Nitha Rani, R. 1994. Isolation and characterization of pathogenic bacterial species in the silkworm, *Bombyx mori* L. *Sericologia*. **34**: 97-102.
- Anusha, H. G. and Bhaskar , R. N. 2015. Effect of bed disinfectants and seasonal incidence of silkworm diseases in stone and RCC rearing houses. *The Bioscan*. **10(3)**: 1027-1029.
- Anusha , H. G. and Bhaskar, R. N., 2016. Per oral inoculation of Bacillus species(Surface and Midgut flora) on larval weight of PM and CSR2. *The Bioscan*. **11(1)**: 193-195.
- Aruga, H. 1994. Diseases of silkworm. In: Principles of Sericulture (Translated from Japanese). Oxford and IBH Publishing Company Ltd., New Delhi. pp. 207-215.
- Ashok kumar, H. K. and Ramakrishna, S. 2013. Isolation and Characterization of Serratia species in silkworm rearing environment. *International J. Advanced Research*. **1(10)**: 465-472.
- Bala venkatasubbaiah, M. Vijayakumari, K. M. Nataraju, B. Rajan, R. K. and Thiagarajan, V. 2006. Effect of different stress factors on the spread of grasserie and flacherie diseases in silkworm rearing. *Bull. Ind. Acad. Seric.* **10(1)**: 27-32.
- Chitra, C. Bhandarkar, A. Karanth, N. G. K. and Vasantharajan, V. N. 1973. Studies on 'Sappe' disease of the silkworm, *Bombyx mori* L. II. Isolation and characterization of pathogenic bacteria from diseased silkworms. *Curr. Sci.* **42**: 373-376.
- Doreswamy, C. Govindan, R. and Devaiah, M. C. 2001. A field survey on the incidence of Thatte disease of silkworm, *Bombyx mori* L. Abstracts of National Seminar on Mulberry Sericulture Research in India, Thalaghattapura, Bangalore, pp.185-186.
- Govindan, R. Narayanaswamy, T. K. and Devaiah, M. C. 1998. Principles of Silkworm Pathology. Seri. Scientific Publishers, Bangalore. p. 420.
- Paramasiva , I. and Rajendra Prasad, P. 2009. Morphological, histopathological and biochemical changes in silkworm, *Bombyx mori* L. due to incidence of bacterial flacherie caused by *Bacillus thuringiensis*. *Indian J. Seric.* **48(1)**: 81-83.
- Pasteur, L. 1870. Etudes sur la maladie des vers de soie. Ganthiervillars, Paris. **2**: 337.
- Robert , A. Pollack, Lorraine Findlay, Water Mondschein and R. Ronald Modesto. 2002. Laboratory Exercises in Microbiology, John Wiley and Sons, INC. p. 120.
- Samson, M. V. 1995. Flacherie in *Bombyx mori* L. *Indian Silk*, **33(11)**: 31-32.
- Samson, M. V. Nataraju, B. Baig, M. and Krishna swami, S. 1981. Starvation of *Bombyx mori* L. on cocoon crop and incidence of loss due to diseases. *Indian J. Seric.* **20**: 42-47.
- Selvakumar, T. 2013. Prevalence of flacherie diseases and pathogenicity of isolated pathogens in silkworm, *Bombyx mori* L. under different environmental conditions. *Agric. Sci. Digest*. **33(4)**: 253-258.
- Selvakumar, T. and Savithri, M. 2012. Thermotherapy of infectious flacherie in young-age silkworm larvae of *Bombyx mori* L., *J. Sericulture and Technology*. **3(1)**: 62-64.
- Sirmani, A. T. Meena, P. and Vanitha Rani, R. 1994. Isolation and characterization of pathogenic bacterial species in the silkworm, *Bombyx mori* L. *Sericologia*. **34**: 97-102.
- Sundarraj, N., Nagaraju, S., Venkataramu, M. N. and Jagannath, M. K. 1972. Design and Analysis of field experiments. Directorate of research, UAs, Bangalore, p. 419.
- Sugun, R. 2000. Status and management of Thatte disease of silkworm. Advances in Tropical Sericulture. Proc. Natl. Sem. Tropic. Seric., UAS, GKVK, Bengaluru, pp. 84-86.
- Swathi, H. C., 2015. Isolation and molecular characterization of bacteria associated with flacherie disease of silkworm, *Bombyx mori* L. Ph.D. (Seri.) Thesis, UAS, GKVK, Bengaluru. pp. 61-69.



**NATIONAL ENVIRONMENTALISTS
ASSOCIATION**

The National Environmentalists Association is chartered in Ranchi as a nonprofit scientific and educational association of like minded academicians, researchers, scientists from all over the nation for the furtherance and diffusion of knowledge of Life Sciences in general and Environmental Science in particular.

The association not only honours its members but also provides FELLOWSHIP to outstanding contributors to the subject and the society.

Contact :

For Editorial Information

Prof. M. P. Sinha
Vice Chancellor
Sido Kanhu Murmu University
Dumka - 814 110
Jharkhand, INDIA

For information regarding Association :

SECRETARY,
National Environmentalists Association,
D-13, Sai Roofs, 1st Floor,
H. H. Colony,
Ranchi - 834002
Jharkhand, India

E-mails : editor.bioscan@gmail.com
dr.mp.sinha@gmail.com
nat.env.assoc@gmail.com

Cell : 94313-60645; 9572649448

Ph. : 0651-2244071

Website : www.thebioscan.com
: www.neaindia.org

NAAS Rating : 5.26

U.S.A. Office

2827 Videre Dr.,
Wilmington,
DE 19808 We, USA

Type setter Bandana Solutions Facility Management LLP
Published by Aditi Publications, Patliputra, Patna

The Bioscan

An International Quarterly Journal of Life Sciences

ISSN : 0973-7049

Volume 15(3) : 2020

Published as an official organ by

NATIONAL ENVIRONMENTALISTS ASSOCIATION

CONTENTS

Page

A. RESEARCH PAPER

1. Comparison and molecular profiling of Begomovirus infecting chilli (*Capsicum annum*) in gangetic alluvial zone of West Bengal
Uday Bikash Oraono, Lourembam Sanajaoba Singh and Jayanta Tarafdar—275 - 280
2. Assessment of probiotic characteristics of *L.plantarum*
Sravani Kandula And Rita Narayanan—281 - 286
3. Effect of rate and frequency of micronutrient on growth attributes and dry matter yield of Banana Cv. grand naine under south Gujarat condition
Narendra Singh, Sonal Tripathi, Patel V. A., Jaimin Naik and Chauhan Aditi—287 - 290
4. Character association and path analysis for seed yield and its components in Grass pea (*Lathyrus sativus* L.)
Gangishetti Ranjithkumar, Sandip Debnath and Duddukur Rajasekhar—291 - 295
5. Determination of physical and biometric properties of onion bulbs in relation to design of digger cum windrower
Shiddanagouda Yadachi and Kiran Nagajjanavar—297 - 301
6. Polygenic variation for morphological and biochemical traits of brinjal genotypes (*Solanum melongena* L.) and its wild relatives
Nisha Sharma, K. D. Bhutia, Rajesh Kumar, Sita Kumari Prasad, Ankita Debnath and Malay Marut Sharma—303 - 309
7. Effect of different sources of horizontal transmission of bacterial flacherie on Et_{50} for symptom expression and mortality of PM X CSR₂
B. L. Kavyashree., R. N. Bhaskar and C. Doreswamy—311 - 314
8. Comparative biology of *Goniozus nephantidis* (Muesbeck) on *Galleria mellonella* L. and *Corcyra cephalonica* (Stainton)
A. V. Desai, M. R. Siddhapara and N. P. Trivedi—315 - 321
9. Efficiency of ovatide on mass seed production of climbing perch (*Anabas testudineus*, Bloch, 1972) in Nalbari district, Assam
Ankur Rajbongshi, A. Ali, M. Chakravarty, M. Deka, H. Mazumdar, Pranab Kr Das and S. Baishya—323 - 326
10. Study of combining ability and gene action for yield and yield component characters in interspecific hybrids of cotton (*Gossypium hirsutum* L. X *Gossypium barbadense* L.)
S. B. Gohil., M. B. Parmar., M. P. Patel and D. A. Patel—327 - 333
11. Per oral inoculation of *Lysinibacillus sphaericus* with pathogenic microbes on rearing and cocoon parameters of silkworm, *Bombyx mori* L.
H. G. Anusha, R. N. Bhaskar and K. V. Anitharani—335 - 338
12. Effect of fungicides, plant extracts and bioagents on spore germination of *Colletotrichum lindemuthianum* causing field bean anthracnose

The Journal is Currently Abstracted / Indexed in

- Paryavarn Abstract, INDIA
- Indian Science Abstract, INDIA
- Cambridge Science Abstract, U.S.A.
- Zoological Record, U.K.
- Directory of Open Access Journal (DOAJ)
- Chemical Abstract, U. S. A.
- Research BIB
- Indian Science
- Journal Seek
- Scientific Indexing Service (SIS)
- Journal is currently rated by
- Index Copernicus
- Universal Impact Factor
- NAAS

DISCLAIMER

The Publisher and Editors cannot be held responsible for errors or any consequences arising from the use of information in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher/ Association and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher / Association and Editors of the products advertised.

- S. Narasimha Rao, S.L. Bhattiprolu, A. Vijaya Gopal and V. Sekhar— 339-344
13. Evaluation in vitro different fungicides for growth of *Rhizoctonia bataticola*
A.M. Kadam, S.S. Chavan and A.H. Kendre — 345- 349
 14. Evaluation of gladiolus varieties for flowering and cut flower traits under indo-gangetic plains
Girish, P. M., Anjana Sisodia and Anil K. Singh — 351- 355
 15. Effect of land configuration and different organic sources on growth, yield and quality of carrot under organic farming
B. Solanki, A. R. Kaswala, P.K. Dubey and A.P. Italiya — 357- 362
 16. Verification and usability analysis of medium range weather forecast for the Kokrajhar district of lower Brahmaputra valley zone of Assam
Kuldip Medhi, Kushal Sarmah, Vinod Upadhyay, Sunil Kumar Paul, Athar N. Islam and Bikash J. Gharphalia — 363 - 370
 17. Canonical root analysis and clustering for characterization and evaluation of aromatic rice germplasm based on morphological characters
G. Parimala, Ch. Damodhar Raju, L.V. Subba Rao and K. Uma Maheswari — 371 - 374
 18. Genetic divergence analysis of sesame genotypes (*Sesamum indicum* L.)
Dasari Rajitha, T. Srikanth, D. Padmaja and T. Kiran Babu — 375 - 379
 19. Nutrient uptake and chemical properties of soil after harvest of baby corn (*Zea mays* L.) as influenced by organic manures and fertilizers
D.H. Roopashree, S. Kamal Bai, Nagaraju and S. Raghavendra — 381 - 384
 20. Genotypic response on growth and yield in papaya
D. K. Varu., K. D. Patel and Sandip Makhmale — 385 - 389
 21. Studies on frequency distribution of yield and yield related traits in F_2M_2 generation of sesame (*Sesamum indicum* L.)
Rajesh Kumar Kar, Tapash Kumar Mishra and Banshidhar Pradhan — 391 - 395
 22. Correlation and path analysis in cowpea (*Vigna unguiculata* (L.) Walp)
R.M. Nagalakshmi, R. Usha Kumari and R. Ananda Kumar — 397 - 401
 23. A simple and efficient method for DNA extraction from rabi sorghum [*Sorghum bicolor* (L.) Moench]"
S. S. Gadakh, G. D. Khalekar, U. S. Dalvi, A. A. Kale and P.L. Kulwal — 403 - 406
 24. Determination of economic injury level (EIL) of sugarcane planthopper borer *Chilo tumidicostalis* Hampson (Lepidoptera: pyralidae)
R. K. Nath and D. K. Saikia — 407- 409
 25. Performance of different summer mung (*Vigna radiata* L.) varieties sown at different dates under Manipur valley condition
Meghna Gogoi, Jamkhogin Lhungdim, Kamal Kant, Urjashi Bhattacharya and Gauri Mohan — 411 - 414



ZONAL CO-ORDINATORS OF THE ASSOCIATION

- Prof. N. Behera
School of Life Science,
Sambalpur University
- Dr. Nirmal Kumar
ISTAR, Vallabh Vidyanagar,
Anand, Gujarat
- Dr. P. N. Sudha
D. K. M. College for Women,
Vellore
- Prof. S. P. S. Dutta
Dept. of Environmental Science,
Jammu University, Jammu
- Dr. V. Salom Gnana Thanga
Dept. of Env. Scs.,
University of Kerala, Kariavattom
Tiruvananthapuram, Kerala

Publications of the Association

The Bioscan

An International Quarterly Journal of Life Sciences

The Ecoscan

An International Quarterly Journal of
Environmental Sciences

**Both the Journals are
online**

**Both the Journals are
available on Google.com**

Websites of the Journals are
www.theecoscan.in
www.thebioscan.in

FEATURES OF ASSOCIATION

- Association is registered under 80G of I.T.
- Prestigious fellowship of the Association (F. N. E. A.) to academicians of the nation.
- Regular annual conference of national and international levels organized by the Association.
- Young Scientist and Senior Scientist award during the conference of the Association.



_____ 062 & 082

_____ 000 & 000

THE BIOSCAN : SUBSCRIPTION RATES				
		India (Rs.)	SAARC Countries	Other Countries
Individuals	One Year	1,000	2,000(I:C)	US \$200
	Life Member*	10,000		
Institutions	One Year	3,000	6,000(I:C)	US \$400
	Life Member*	30,000		

*Life Member will receive the journal for 15 years while other benefits will continue whole life

THE BIOSCAN : MEMBERSHIP FORM

Please enter my subscription for the above journal for the year / life member.

Name:

Address:

E-mail:

Payment Rs. : by DD / MD in favour of
National Environmentalists Association payable at Ranchi, No. Dated
..... is enclosed.

NOTE: FOR MEMBERSHIP THE ABOVE INFORMATION CAN BE SENT ON SEPARATE SHEET