

EVALUATION IN VITRO DIFFERENT FUNGICIDES FOR GROWTH OF RHIZOCTONIA BATATICOLA

A.M. KADAM, S.S. CHAVAN AND A.H. KENDRE

Department of Plant Pathology, Sau. K.S.K. college of Agriculture, Beed,
Vasantrao Naik Marathwada Krishi Vidhyapeeth, Parbhani - 431 402.

e-mail: kadamamruta14@gmail.com

KEYWORDS

Chickpea
Dry root rot
Fungicides.

Received on :
22.05.2020

Accepted on :
07.08.2020

***Corresponding
author**

ABSTRACT

The present study was conducted to test the efficacy of some systemic, non-systemic and combine fungicides like Propiconazole, carbendazim, captan, carboxin, benomyl, mancozeb, thiram, carbendazim + mancozeb against dry root rot of chickpea. A variant with no application of fungicide was used as control. In laboratory condition result revealed that fungicides evaluated against *R.bataticola* exhibited a wide range of mycelial growth of *R. bataticola* and was found to be decreased drastically and showed inhibition with increased concentration of fungicides tested.

The suppression of growth of pathogen was maximum with Carbendazim (Bavistin 50%WP) (100.00 per cent) followed by Benomyl (Benlate 50 WP) (93.61 (75.35)) per cent and the least effective fungicides was Carboxin (Vitavax 75 WP) (61.17 (51.45) per cent). Carbendazim (Bavistin 50%WP) showed best performance against the pathogen.

INTRODUCTION

The cultivated chickpea (*Cicer arietinum* L.) was one of the first grain legume to be domesticated in the old world. The genus *Cicer* belongs to family Leguminosae and sub family Papilionoidae. Chickpea has been well recognized as a valuable source of protein particularly in the developing countries, where majority of the populations depends on the low priced food for meeting its dietary requirements nutritionally, chickpea is low in sodium, contains (21.1%), fats (4.5%), no cholesterol and overall an excellent source of both soluble and insoluble fiber, complex carbohydrates (61.5%), vitamins (especially B vitamins) and minerals (especially potassium, phosphorous, calcium, magnesium, copper, iron and zinc). Therefore, chickpea is an excellent healthy food that may be beneficial to the prevention of coronary and cardiovascular diseases and by reducing blood lipids also help some serious complications of diabetes.

In India chickpea is primarily grown as Rabi (post rainy) season crop on residual soil moisture. It ensures nutritional security besides being a rich source of protein and is also important in substantial agriculture as it improves physical, chemical and biological properties of soil by mixing atmospheric nitrogen symbiotically. Deep roots of pulse crop also open up the soil by increasing soil aeration and fit well in various cropping ecosystem and hence have got unique position in rainfed agriculture.

Chickpea is grown throughout the world. In India on large scale in Punjab, Hariyana, Uttar Pradesh, Madhya Pradesh, Rajasthan, Andhra Pradesh, Karnataka, and Maharashtra it is mostly cultivated under Rainfed condition in a variety of soil,

varying in a residual moisture. In India area under chickpea was 10.57 M/ha with production 11.16 MT and productivity 1056 kg/ha during 2018. susceptibility of the crop to different biotic and abiotic stress. Abiotic stress is basically due to insufficient moisture. regarding biotic stresses, diseases, insect pest, nematodes and parasitic weeds account major losses for example, extend of yield loss due to wilt and root rot disease is far more in the event of drought high temperature in the country. The chick pea crop is attacked by 172 pathogen viz as 67 fungi, 22 viruses, 3 bacteria, 80 nematodes and mycoplasma from all over the world (Nene et al., 1996).

Being soil borne in a nature it is very difficult to control this pathogen. In primary use of resistant cultivar is more effective strategy for sustainable chickpea cultivation, and cultural scouting, sanitation and development of the host plant resistance with the application of fungicides. Regarding the management of dry root rot of chickpea many workers had done lots of works based on chemical control. Earlier workers reported application of fungicides is the most effective method (Sangeetha and Jahagirdar 2013), reported that dry root rot causing pathogen *Rhizoctonia bataticola*, In vitro screening of fungicides was undertaken to identify an effective compound against *R. bataticola*. Mancozeb, carbendazim, thiophanate methyl, hexaconazole, propiconazole, were found effective in inhibiting mycelia growth of pathogen. Therefore, the present investigation was carried out to evaluate with some systemic, non-systemic and combine fungicides for management of dry root rot of chickpea.

MATERIALS AND METHODS

The effect of nine fungicides on *R.bataticola* (Taub) Butler was

evaluated in-vitro by employing "Poisoned food technique" (Kumari *et al.*, 2012). The requisite amount of each fungicide @ 500, 1000 and 2000 ppm based on active ingredient was added in an autoclaved and cooled (450C) potato dextrose agar medium to obtain desired concentrations (Singh and Chohan), (Ramadoss and Sivaprakasam). The fungicides used with their concentrations were propiconazole, carbendazim, captan, carboxin, benomyl, mancozeb, thiram, carbendazim + mancozeb, Plain PDA without fungicide served as control. The PDA was then poured into sterile glass petriplates (90 mm diameter) and allowed to solidify. These plates were inoculated with a 5 mm mycelial disc of the test fungus and incubated for 7 days at 28 ± 20 C. All the treatments were replicated thrice. Observations on radial mycelial growth in treatment plates were recorded after 7-8 days. Per cent inhibition of the test fungus over control was calculated by applying the formula given below, (Vincent 1947).

Details of the experiment :-

Design	- Complete Randomized Design (CRD)
Replications	- Three
Treatments	- Nine
T1	- Propiconazole
T2	- Carbendazim
T3	- Captan
T4	- Carboxin
T5	- Benomyl
T6	- Mancozeb
T7	- Thiram
T8	- Carbendazim + mancozeb
T9	- Control

$$\text{Per cent inhibition (I)} = \frac{C - T}{C} \times 100$$

Where,

C = Growth (mm) of test fungus in control plate.

T = Growth (mm) of test fungus in treatment plate.

RESULTS AND DISCUSSION

Results (Table and Fig.) revealed that fungicides evaluated in-vitro against *R. bataticola* exhibited a wide range of mycelial growth of *R. bataticola* and was found to be decreased drastically and showed inhibition with increased concentration of fungicides tested.

Mycelial growth


At 500 ppm, radial mycelial growth of the test pathogen was ranged from 00.00 mm (carbendazim) to 37.94 mm (carboxin), as against 90.00 mm in untreated control. However, significantly least mycelium growth was recorded in carbendazim (0.00mm). This was followed by the fungicides viz., benomyl (07.22mm), Saff (08.19mm) and mancozeb (09.83mm) all these three were statistically at par. These were followed by thiram (16.07mm), propiconazole (26.95mm) and captan (30.12mm). Fungicide carboxin (37.94mm) was found comparatively less effective with maximum mycelial growth (Singh and Chohan).

At 1000 ppm, all the 09 fungicides tested exhibited similar trend of mycelial growth as that of observed at 500 ppm, but growth was comparatively reduced and was ranged from 00.00 mm (carbendazim) to 35.58 mm (carboxin), as against 90.00 mm in untreated control. However, significantly least mycelial growth was recorded in carbendazim (00.00mm). This was followed by the fungicides viz., benomyl (05.08mm), Saff

Table1: In vitro efficacy of fungicides against mycelial growth and inhibition of *R. bataticola*.

Tr. No.	Treatments	Col. dia. *(mm) at conc.			Av. (mm)	% Inhibition*			Av. (%) inhibition
		500 ppm	1000 ppm	2000 ppm		500 ppm	1000 ppm	2000 ppm	
T ₁	Propiconazole (Tilt 25% EC)	26.95	22.39	13.06	20.07	70.06 (56.82)	75.12 (60.07)	78.58 (62.43)	74.58 (59.72)
T ₂	Carbendazim (Bavistin 50% WP)	0	0	0	0	100 (90)	100 (90)	100 (90)	100.00 (90.00)
T ₃	Captan (Captaf 75% WP)	30.12	26.17	19.19	25.16	66.52 (54.64)	70.91 (57.36)	78.83 (62.6)	72.08 (58.1)
T ₄	Carboxin (Vitavax 75 WP)	37.94	35.58	31.28	26.02	57.83 (49.5)	60.45 (51.03)	65.24 (53.87)	61.17 (51.45)
T ₅	Benomyl (Benlate 50 WP)	7.22	5.08	5.06	6.2	92.13 (73.7)	94.34 (76.23)	94.37 (76.27)	93.61 (75.35)
T ₆	Mancozeb (Indofil M-45 75% WP)	9.83	6.76	5.65	7.41	89.13 (70.74)	92.48 (74.08)	93.88 (75.67)	91.83 (73.39)
T ₇	Thiram (Thiram 75% WP)	16.07	16.07	11.94	14.69	82.13 (64.99)	82.13 (64.99)	86.72 (68.62)	83.66 (66.15)
T ₈	Carbendazim + mancozeb (Saff 75% WP)	8.19	6.09	5	6.69	90.89 (72.43)	93.25 (74.94)	94.59 (76.55)	92.91 (74.55)
T ₉	Control (Untreated)	90	90	90	90	0 0	0 0	0 0	000.00 (00.00)
	S.E. ±	1.2	0.62	2.68	1.5	1.34	0.69	0.67	0.9
	C.D. (P=0.01)	3.27	1.68	7.29	4.08	3.64	1.87	1.82	2.44

*-Mean of three replications, Col. = Colony, Dia. = Diameter, Conc. = Concentration; Av. = Average, Figures in parenthesis are arc sine transformed value.



Effect of fungicide on mycelial growth of *R.bataticla*
C = control, 1 = 500 ppm, 2 = 1000 ppm, 3 = 2000 ppm

Plate V

(6.09mm) and mancozeb (06.76mm) all these three were statistically at par and was followed by thiram (16.07mm), propiconazole (22.39mm) and captan (26.17mm). In fungicides, fungicide carboxine was found least effective with maximum colony diameter (35.58mm).

At 2000 ppm, all the fungicides tested exhibited somewhat similar trend of mycelial growth as that of observed at 500 ppm and 1000 ppm and it was ranged from 00.00 mm (carbendazim) to 31.28 mm (carboxin), as against 90.00 mm untreated control. However, with the fungicide carbendazim significantly least mycelium growth (00.00mm) was observed. This was followed by the fungicides viz., Saff (05.00mm), benomyl (05.06mm), mancozeb (5.65mm) and thiram which at par. Followed by other fungicides propiconazole (13.06mm) and captan (26.17mm), while the fungicide carboxin was recorded comparatively less effective with maximum mycelia growth of (31.28mm), (Ramdoss and Sivaprakasam).

Average radial mycelial growth recorded in all the 09 fungicides tested was ranged from 00.00 mm (carbendazim) to 26.02 mm (carboxin), as against 90.00 mm in untreated control. However, the least average mycelia growth was recorded with carbendazim (00.00mm), (Taneja and Grover). This was followed by the fungicides viz. benomyl (06.20mm), Saff (06.69mm) and mancozeb (07.41mm) all these three were at par and was followed by thiram (14.69mm), propiconazole (20.07mm) and captan (25.16mm). The highest average colony diameter was recorded in carboxin (26.02mm).


Mycelial growth inhibition

Results (Table and Fig.) revealed that all the fungicides tested (@ 500, 1000 and 2000 ppm each) significantly inhibited mycelial growth of *R.bataticola* over untreated control (00.00%). Further, the percentage mycelial growth inhibition of test pathogen was increased with increase in concentrations


of the fungicides tested (PLATE 5 and 6).

At 500 ppm, percentage mycelial growth inhibition of the test pathogen was ranged from 57.83 (carboxin) to 100.00 per cent (carbendazim). However, significantly higher mycelia inhibition was recorded with fungicide carbendazim (100.00%). The second, third and fourth best fungicides found were benomyl (92.13%) and Saff (90.89%) and mancozeb (89.13%) which were at par followed by thiram (82.13%), propiconazole (70.06%), captan (66.52%). The fungicide carboxin was found less effective with 57.83 per cent inhibition of the test pathogen.

At 1000 ppm, mycelial growth inhibition was increased compared to 500 ppm and similar trends of mycelial growth inhibition were observed. It was ranged from 60.45 (carboxin) to 100.00 (carbendazim) per cent. However, significantly highest mycelia inhibition was found with fungicide carbendazim (100.00%). The second, third best fungicides,



In vitro efficacy of fungicides against mycelial growth and inhibition of *R.bataticola*.



Effect of fungicide on mycelial growth of *R. bataticola*
C = control, 1 = 500 ppm, 2 = 1000 ppm, 3 = 2000 ppm

Plate vi

viz. benomyl (94.34%), Saff (93.25%) both were at par. These were followed by fungicides *viz.* mancozeb (92.48%), thiram (82.13%), propiconazole (75.12%), and captan (70.91%). The fungicide carboxin was found less effective with 60.45mm inhibition of the test pathogen.

At 2000 ppm, all the 09 fungicides tested exhibited comparatively increased mycelial growth inhibition than that of at 500 and 1000 ppm and it was ranged from 65.24 (carboxin) to 100 (carbendazim) per cent. However, significantly highest mycelia inhibition was recorded with fungicide carbendazim (100.00%). This was followed by the fungicides, *viz.* saff (94.59%), benomyl (94.37%) and mancozeb (93.88%) were at par followed by thiram (86.72%). The fungicides captan (78.83%), propiconazole (78.58%) were at par the fungicide carboxin was reported comparatively less effective with growth inhibition of 65.24%.

Average mycelial growth inhibition recorded in the test fungicides was ranged from 61.17 (carboxin) to 100.00 (carbendazim) per cent. However, significantly by highest average mycelial growth inhibition was recorded in carbendazim (100.00%). The second and third best fungicides found were benomyl (93.61%) and Saff (92.91%), both were at par. These were followed by the fungicides *viz.*, mancozeb (91.83%), thiram (83.66%), propiconazole (74.58%), captan (72.08%). The fungicide carboxin was comparatively less effective with minimum growth inhibition of 61.17 per cent. Thus, all the fungicides tested were found fungistatic against *R. bataticola* and significantly inhibited its mycelial growth over untreated control. However, fungicides found most effective in the order of merit were carbendazim, benomyl, Saff, mancozeb, thiram, propiconazole and captan.

Similar fungistatic effects of the test fungicides against *R. bataticola* infecting chickpea and many other crops were

reported earlier by several workers, Goyal and Mehrotra 1981, Taneja and Grover 1982, Sarwar and Gopal Raju 1985, Ramadoss and Shivprakasham 1987, Singh and Sindan 1988, Peshney *et al.*, 1992, Prajapati *et al.*, 2004, Sandipan *et al.*, 2007, Date 2015.

REFERENCES

- Bhaliya, C.M. and Jadeja, K. B. 2014.** Efficacy of different fungicides against coriander root rot. *The Bioscan*. **9(3)**:1225-1227.
- Bansal, R. K. and Asha Shivpuri. 1988.** Evaluation of some systemic and non-systemic Fungicides in vitro against *R. bataticola* (Taub) Butler. *Pesticide*. **22(11)**:48-49
- Goyal, M. K. and Mehrotra, R. S. 1981.** Chemical control of dry root-rot of gram caused by *R. bataticola*. *Acta- Botanica- India*. **(9)**:228-232.
- Kumari, R. Shekhawat, K. S., Gupta Renu and Khokar, M.K. 2012.** Integrated Management against Root-rot of Mungbean [*Vignaradiata* (L.) Wilczek] incited by *Macrophomina phaseolina*. *J. Plant Pathol. Microb.* **3**:136.
- Nene, Y. L. 1985.** Opportunities for research on diseases of pulses crops. *Indian Phytopath.* **38**: 1-10.
- Peshney, N. L. Gade, R. M. and Thakre K. G. 1992.** Sensitivity and adaptability of *Rhizoctonia bataticola* to different fungicides. *J. Soils crop*. **2(1)**:35-38.
- Prajapati, R. K. Gangwar, R. K. and Shrivastava, S.L. 2003.** Resistance source of chickpea against dry root rot. *J. Farm Sci.* **12(1)**: 86.
- Ramadoss, S. and Sivaprakasam, K. 1987.** Effect of fungicides and insecticides on the linear growth of *Macrophomina phaseolina* (Tassi.) *Goid. Madras Agril. J.* **74(1)**:29-33.
- Sangeetha, T.V. and Jahagirdar, S. 2013.** Survey for the assessment of disease incidence of root rot/ wilt of soyabean in Northern Karnataka. *J. Agric. Sci.* **26(4)**:563-564.
- Sarwar, H.A.K. and Gopala Raju D. 1985.** Topsin M-70 the most effective fungicide for the control of *Rhizoctonia bataticola*, the root

rot and stem rot disease causing pathogen in castor. *Pesticides*.**8(1)**:25-29.

Singh R.S. and chohan,J.S. 1981.Effect of benlate on growth and spore germination of *Macrophomina phaseolina*.*Pesticides*.15

Sandipan, P.B., Singh, A., Patel, J.G., Patel R.L. and Sharma K. 2007.Efficacy of fungicides against dry root rot and charcoal rot of potato. *J. Mycol. Pl Pathol*.**37(1)**:183.

Singh, R. and Sindhan, G.S. 1998. Effect of fungicides on the incidence

of dry root and biochemical status of Chickpea plants. *Indian J .Mycol. Pl. Pathol.* **13(1)**: 14-17.

Taneja,M. and Grover, R. K. 1982. Efficacy of benzimidazole and related fungicides against *Rhizoctonia solani* and *R. bataticola*. *Annals of Applied Biology.* **100**: 425-432.

Vincent, J.M1947. Distortion of fungal hyphae in the presence of certain inhibitor.*Nature*.**159**: 239-241.



**NATIONAL ENVIRONMENTALISTS
ASSOCIATION**

The National Environmentalists Association is chartered in Ranchi as a nonprofit scientific and educational association of like minded academician, researchers, scientists from all over the nation for the furtherance and diffusion of knowledge of Life Sciences in general and Environmental Science in particular.

The association not only honours its members but also provides FELLOWSHIP to outstanding contributors to the subject and the society.

Contact :

For Editorial Information

Prof. M. P. Sinha
Vice Chancellor
Sido Kanhu Murmu University
Dumka - 814 110
Jharkhand, INDIA

For information regarding Association :

SECRETARY,
National Environmentalists Association,
D-13, Sai Roofs, 1st Floor,
H. H. Colony,
Ranchi - 834002
Jharkhand, India

E-mails : editor.bioscan@gmail.com
dr.mp.sinha@gmail.com
nat.env.assoc@gmail.com

Cell : 94313-60645; 9572649448

Ph. : 0651-2244071

Website : www.thebioscan.com
: www.neaindia.org

NAAS Rating : 5.26

U.S.A. Office

2827 Videre Dr.,
Wilmington,
DE 19808 We, USA

Type setter Bandana Solutions Facility Management LLP
Published by Aditi Publications, Patliputra, Patna

The Bioscan

An International Quarterly Journal of Life Sciences

ISSN : 0973-7049

Volume 15(3) : 2020

Published as an official organ by

NATIONAL ENVIRONMENTALISTS ASSOCIATION

CONTENTS

Page

A. RESEARCH PAPER

1. Comparison and molecular profiling of Begomovirus infecting chilli (*Capsicum annum*) in gangetic alluvial zone of West Bengal
Uday Bikash Oraono, Lourembam Sanajaoba Singh and Jayanta Tarafdar—275 - 280
2. Assessment of probiotic characteristics of *L.plantarum*
Sravani Kandula And Rita Narayanan—281 - 286
3. Effect of rate and frequency of micronutrient on growth attributes and dry matter yield of Banana Cv. grand naine under south Gujarat condition
Narendra Singh, Sonal Tripathi, Patel V. A., Jaimin Naik and Chauhan Aditi—287 - 290
4. Character association and path analysis for seed yield and its components in Grass pea (*Lathyrus sativus* L.)
Gangishetti Ranjithkumar, Sandip Debnath and Duddukur Rajasekhar—291 - 295
5. Determination of physical and biometric properties of onion bulbs in relation to design of digger cum windrower
Shiddanagouda Yadachi and Kiran Nagajjanavar—297 - 301
6. Polygenic variation for morphological and biochemical traits of brinjal genotypes (*Solanum melongena* L.) and its wild relatives
Nisha Sharma, K. D. Bhutia, Rajesh Kumar, Sita Kumari Prasad, Ankita Debnath and Malay Marut Sharma—303 - 309
7. Effect of different sources of horizontal transmission of bacterial flacherie on Et_{50} for symptom expression and mortality of PM X CSR₂
B. L. Kavyashree, R. N. Bhaskar and C. Doreswamy—311 - 314
8. Comparative biology of *Goniozus nephantidis* (Muesbeck) on *Galleria mellonella* L. and *Corcyra cephalonica* (Stainton)
A. V. Desai, M. R. Siddhapara and N. P. Trivedi—315 - 321
9. Efficiency of ovatide on mass seed production of climbing perch (*Anabas testudineus*, Bloch, 1972) in Nalbari district, Assam
Ankur Rajbongshi, A. Ali, M. Chakravarty, M. Deka, H. Mazumdar, Pranab Kr Das and S. Baishya—323 - 326
10. Study of combining ability and gene action for yield and yield component characters in interspecific hybrids of cotton (*Gossypium hirsutum* L. X *Gossypium barbadense* L.)
S. B. Gohil, M. B. Parmar, M. P. Patel and D. A. Patel—327 - 333
11. Per oral inoculation of *Lysinibacillus sphaericus* with pathogenic microbes on rearing and cocoon parameters of silkworm, *Bombyx mori* L.
H. G. Anusha, R. N. Bhaskar and K. V. Anitharani—335 - 338
12. Effect of fungicides, plant extracts and bioagents on spore germination of *Colletotrichum lindemuthianum* causing field bean anthracnose

The Journal is Currently Abstracted / Indexed in

- Paryavarn Abstract, INDIA
- Indian Science Abstract, INDIA
- Cambridge Science Abstract, U.S.A.
- Zoological Record, U.K.
- Directory of Open Access Journal (DOAJ)
- Chemical Abstract, U. S. A.
- Research BIB
- Indian Science
- Journal Seek
- Scientific Indexing Service (SIS)
- Journal is currently rated by
- Index Copernicus
- Universal Impact Factor
- NAAS

DISCLAIMER

The Publisher and Editors cannot be held responsible for errors or any consequences arising from the use of information in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher/ Association and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher / Association and Editors of the products advertised.

- S. Narasimha Rao, S.L. Bhattiprolu, A. Vijaya Gopal and V. Sekhar— 339-344
13. Evaluation in vitro different fungicides for growth of *Rhizoctonia bataticola*
A.M. Kadam, S.S. Chavan and A.H. Kendre — 345- 349
 14. Evaluation of gladiolus varieties for flowering and cut flower traits under indo-gangetic plains
Girish, P. M., Anjana Sisodia and Anil K. Singh — 351- 355
 15. Effect of land configuration and different organic sources on growth, yield and quality of carrot under organic farming
B. Solanki, A. R. Kaswala, P.K. Dubey and A.P. Italiya — 357- 362
 16. Verification and usability analysis of medium range weather forecast for the Kokrajhar district of lower Brahmaputra valley zone of Assam
Kuldip Medhi, Kushal Sarmah, Vinod Upadhyay, Sunil Kumar Paul, Athar N. Islam and Bikash J. Gharphalia — 363 - 370
 17. Canonical root analysis and clustering for characterization and evaluation of aromatic rice germplasm based on morphological characters
G. Parimala, Ch. Damodhar Raju, L.V. Subba Rao and K. Uma Maheswari — 371 - 374
 18. Genetic divergence analysis of sesame genotypes (*Sesamum indicum* L.)
Dasari Rajitha, T. Srikanth, D. Padmaja and T. Kiran Babu — 375 - 379
 19. Nutrient uptake and chemical properties of soil after harvest of baby corn (*Zea mays* L.) as influenced by organic manures and fertilizers
D.H. Roopashree, S. Kamal Bai, Nagaraju and S. Raghavendra — 381 - 384
 20. Genotypic response on growth and yield in papaya
D. K. Varu., K. D. Patel and Sandip Makhmale — 385 - 389
 21. Studies on frequency distribution of yield and yield related traits in F_2M_2 generation of sesame (*Sesamum indicum* L.)
Rajesh Kumar Kar, Tapash Kumar Mishra and Banshidhar Pradhan — 391 - 395
 22. Correlation and path analysis in cowpea (*Vigna unguiculata* (L.) Walp)
R.M. Nagalakshmi, R. Usha Kumari and R. Ananda Kumar — 397 - 401
 23. A simple and efficient method for DNA extraction from rabi sorghum [*Sorghum bicolor* (L.) Moench]"
S. S. Gadakh, G. D. Khalekar, U. S. Dalvi, A. A. Kale and P.L. Kulwal — 403 - 406
 24. Determination of economic injury level (EIL) of sugarcane planthopper borer *Chilo tumidicostalis* Hampson (Lepidoptera: pyralidae)
R. K. Nath and D. K. Saikia — 407- 409
 25. Performance of different summer mung (*Vigna radiata* L.) varieties sown at different dates under Manipur valley condition
Meghna Gogoi, Jamkhogin Lhungdim, Kamal Kant, Urjashi Bhattacharya and Gauri Mohan — 411 - 414



ZONAL CO-ORDINATORS OF THE ASSOCIATION

- Prof. N. Behera
School of Life Science,
Sambalpur University
- Dr. Nirmal Kumar
ISTAR, Vallabh Vidyanagar,
Anand, Gujarat
- Dr. P. N. Sudha
D. K. M. College for Women,
Vellore
- Prof. S. P. S. Dutta
Dept. of Environmental Science,
Jammu University, Jammu
- Dr. V. Salom Gnana Thanga
Dept. of Env. Scs.,
University of Kerala, Kariavattom
Tiruvananthapuram, Kerala

Publications of the Association

The Bioscan

An International Quarterly Journal of Life Sciences

The Ecoscan

An International Quarterly Journal of
Environmental Sciences

**Both the Journals are
online**

**Both the Journals are
available on Google.com**

Websites of the Journals are
www.theecoscan.in
www.thebioscan.in

FEATURES OF ASSOCIATION

- Association is registered under 80G of I.T.
- Prestigious fellowship of the Association (F. N. E. A.) to academicians of the nation.
- Regular annual conference of national and international levels organized by the Association.
- Young Scientist and Senior Scientist award during the conference of the Association.



_____ 062 & 082

_____ 000 & 000

THE BIOSCAN : SUBSCRIPTION RATES				
		India (Rs.)	SAARC Countries	Other Countries
Individuals	One Year	1,000	2,000(I:C)	US \$200
	Life Member*	10,000		
Institutions	One Year	3,000	6,000(I:C)	US \$400
	Life Member*	30,000		

*Life Member will receive the journal for 15 years while other benefits will continue whole life

THE BIOSCAN : MEMBERSHIP FORM

Please enter my subscription for the above journal for the year / life member.

Name:

Address:

E-mail:

Payment Rs. : by DD / MD in favour of
National Environmentalists Association payable at Ranchi, No. Dated
..... is enclosed.

NOTE: FOR MEMBERSHIP THE ABOVE INFORMATION CAN BE SENT ON SEPARATE SHEET