

# EVALUATION OF GLADIOLUS VARIETIES FOR FLOWERING AND CUT FLOWER TRAITS UNDER INDO-GANGETIC PLAINS

GIRISH, P. M., ANJANA SISODIA\* AND ANIL K. SINGH

Department of Horticulture, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi-221 005, U.P., INDIA.

e-mail: anjana.floriculture@gmail.com

## KEYWORDS

Evaluation  
Gladiolus  
Varieties  
Flowering

**Received on :**  
20.05.2020

**Accepted on :**  
30.07.2020

**\*Corresponding author**

## ABSTRACT

An experiment was conducted at Horticultural Research Farm of the Department of Horticulture, Banaras Hindu University, Varanasi, Uttar Pradesh, India during 2018-19. In this experiment 29 varieties of gladiolus were taken as treatments in a Randomized Block Design with three replications. The variety Chemistry showed early spike emergence (70.88 days), early opening of 1<sup>st</sup>, 3<sup>rd</sup> and 5<sup>th</sup> florets (88.35 days, 88.92 days and 90.40 days, respectively) while, variety Anjali showed late withering of 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup> and last florets (109.30, 110.89, 111.75 and 117.70 days, respectively). The variety American Beauty reported maximum length of 1<sup>st</sup>, 3<sup>rd</sup> and 5<sup>th</sup> florets (10.90 cm, 10.75 cm and 10.57 cm) and maximum diameter of 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup> and last florets (10.55 cm, 10.25 cm, 9.71 cm and 9.18 cm, respectively). The variety Pink Friendship reported maximum flowering duration (14.17 days), maximum number of florets open (15.00) and maximum florets per spike (16.00). Longest spike and inter-node (1<sup>st</sup> to 4<sup>th</sup> floret) was observed in the variety IIHR (74.06 cm and 15.13 cm, respectively) while, longest rachis was in American Beauty (46.35 cm). The varieties Pink Friendship and American Beauty were found superior to others in this study.

## INTRODUCTION

Cut flowers are mainly used for floral decorations and bouquets. Gladiolus is one of the leading cut flower in domestic as well as international flower market (Singh and Sisodia, 2017). Gladiolus (*Gladiolus* sp) is bulbous flower crop belonging to family Iridaceae with a chromosome number of  $2n = 30-120$  and originating from South Africa. The word gladiolus is derived from the Latin word Gladius which means sword, as the leaves of the plant are sword shaped it is also known as Sword lily (Singh, 2014; Mushtaq *et al.*, 2018). Gladiolus is mainly used as cut flower, flower beds, pots, etc. Gladiolus inflorescence is known as spike which is commercially and ornamentally valued part. The spike consists of number of florets arranged alternately on the axis which open acropetally. Gladiolus has gained importance as cut flower since its beautiful spikes contain elegant coloured florets, long flowering duration and good shelf life, the character which are required for a better cut flower (Singh, 2006). Gladiolus has quite a large number of varieties with varying colour (mono or multi), size and texture (plain, frilled or ruffled) (Rashmi and Chandrashekar, 2016). In India, it is mainly cultivated in states like West Bengal, Uttar Pradesh, Karnataka, Himachal Pradesh, Sikkim, Tamil Nadu, Punjab and Delhi (Bhaskar *et al.*, 2019). Indo-Gangetic plain has subtropical climate with sandy loam soil which is suitable for the cultivation of gladiolus, it is even a major market for cut flowers like gladiolus. Gladiolus consists of more than 30,000 established varieties and every year new varieties are being added to the list which makes it necessary for evaluation (Rashmi and Chandrashekar, 2016). Most of the varieties are

purpose specific and agro-climate specific, hence varietal evaluation helps find region specific varieties. Several experiments on varietal evaluation were previously carried out in Indo-Gangetic plains by Singh *et al.* (2013), Singh *et al.* (2017) and Sisodia *et al.* (2018) which reported significant differences between the varieties. Therefore, for further evaluation an experiment was carried out to find the suitable gladiolus varieties under Indo-Gangetic plains of Uttar Pradesh for flowering traits and cut flower production.

## MATERIALS AND METHODS

A field trial was carried out to see the performance of various Indian and exotic varieties of gladiolus for different flowering traits. The experiment was conducted at Horticultural Research Farm of the Department of Horticulture, Banaras Hindu University, Varanasi, Uttar Pradesh, India during the year 2018-19. In this experiment 29 varieties of gladiolus viz. Priscilla, American Beauty, Sunayna, Aldebaran, Anjali, Nova Lux, Snow Princess, Pink Friendship, Jester Gold, IIHR, Chandini, Subhangini, Tiger Flame, Punjab Dawn, Punjab Morning, Dhanvantari, Flevo Laguna, Green Star, Arti, Chemistry, Jyotsna, Pusa Shubham, Pusa Srijan, Surya Kiran, Yellow Star, Princess Margaret Rose, White Prosperity, Darshan and Sabnam were used as treatments. The healthy corms of these varieties were planted at a spacing of 30 cm between the rows and 20 cm between the plants in a Randomized Block Design with three replications on 17<sup>th</sup> of November, 2018. All cultural practices were followed uniformly. Flowering parameters were observed from randomly selected plants. The required measurements were done using standard

meter scale and mean value of the selected plants in each treatment was taken to represent a particular variety with respect to a character and noted observations were later subjected to statistical analysis as suggested by Panse and Sukhatme (1985).

## RESULTS AND DISCUSSION

### Flowering parameters

Significant differences were observed in flowering parameters due to the varieties (Table 1). Early spike emergence was observed in variety the Chemistry (70.88 days) which was at par with IIHR (72.12 days) and Chandini (72.83 days), whereas, late spike emergence was noticed in Anjali (94.38 days). The early colour show was exhibited in variety Jyotsna (83.25 days) which was at par with the varieties Chandini (84.78 days) and Chemistry (85.02 days), whereas, it was last in Anjali (103.83 days). Early opening of 1<sup>st</sup> floret was noticed in the variety Chemistry (88.35 days) which was statistically at par with Jyotsna (88.89 days) and Chandini (89.11 days). Early opening of 3<sup>rd</sup> floret was recorded in the variety Chemistry (88.92 days) which was at par with Jyotsna (90.00 days) and Princess Margaret Rose (91.44 days). Early opening of 5<sup>th</sup> floret was observed in the variety Chemistry (90.40 days) which was at par with Jyotsna (91.92 days), Princess Margaret Rose (92.55 days) and Chandini (93.67 days). Early opening of last floret was in the variety Jyotsna (95.48 days) which was at par with Chandini (96.55 days), Chemistry (97.00 days) and Princess Margaret Rose (98.25 days). The variety Anjali (106.93 days, 108.56 days, 109.56 days and 114.75 days) showed late opening of 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup> and last florets, respectively. The spike

emergence and floret opening depends on the maturity period of the variety and environmental factors, which can be further used for planned cultivation. Differences in days to floret opening were earlier reported by Singh *et al.* (2013), Kadam *et al.* (2014), Singh *et al.* (2017) and Kaur and Bajpay (2019). Late withering of 1<sup>st</sup> floret was in the variety Anjali (109.30 days) which was statistically at par with the varieties Tiger Flame (108.26 days), White Prosperity (106.53 days) and Yellow Star (106.00 days). Late withering of 3<sup>rd</sup> floret was recorded in the variety Anjali (110.89 days) which was at par with the varieties Tiger Flame (108.97 days), White Prosperity (107.58 days) and Flevo Laguna (107.43 days). Late withering of 5<sup>th</sup> floret was observed in the variety Anjali (111.75 days) which was statistically at par with the varieties Tiger Flame (110.06 days), White Prosperity (109.55 days), Flevo Laguna (108.88 days) and Yellow Star (108.00 days). Late withering of last floret was in the variety Anjali (117.70 days) which was at par with the varieties Flevo Laguna (108.88 days) and Tiger Flame (116.07 days). Whereas, early withering of 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup> and last florets was observed in the variety Chemistry (91.03 days, 91.67 days and 92.97 days) and Jyotsna (98.19 days), respectively. The variation in flowering parameters of various cultivars may be due to the genetic makeup of the cultivars which was similar to the findings of Chaudhary and Dhatt (2017), Mushtaq *et al.* (2018), Singh *et al.* (2018) and Kumar *et al.* (2019).

The diameter of 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup> and last floret was maximum in the variety American Beauty (10.55 cm, 10.25 cm, 9.71 cm and 9.18 cm, respectively) which was at par with the variety Priscilla (9.56 cm and 9.40 cm) for 3<sup>rd</sup> and 5<sup>th</sup> florets respectively and with Priscilla (8.93 cm), IIHR (8.90 cm) and Nova Lux (8.68

**Table 1: Performance of gladiolus varieties for flowering parameters**

	Days to spike emergence	Days to colour show	Days to 1 <sup>st</sup> floret opening	Days to 3 <sup>rd</sup> floret opening	Days to 5 <sup>th</sup> floret opening	Days to last floret opening	Days to 1 <sup>st</sup> floret withering	Days to 3 <sup>rd</sup> floret withering	Days to 5 <sup>th</sup> floret withering	Days to last floret withering
Priscilla	74.39	87.31	92.15	93.27	94.6	100.52	95.21	96.4	97.61	103.16
American Beauty	76.62	89.84	94.24	95.76	96.96	104.71	97.09	98.7	99.9	107.14
Sunayna	74.05	87	91.78	92.78	94.39	100.22	94.61	95.89	97.39	103.28
Aldebaran	83.97	94.08	97.58	98.77	100.2	102.38	100.29	101.63	103	105.05
Anjali	94.38	103.83	106.93	108.56	109.56	114.75	109.3	110.89	111.75	117
NovaLux	84	96.17	100.67	101.17	101.33	106.5	103.17	104.33	104.17	108.5
Snow Princess	78.73	91.25	95.18	96.03	97.18	103.17	97.61	98.62	99.83	106.08
Pink Friendship	82.83	94.83	97.56	99	100.25	109.58	100.36	102	103.33	112.33
Jester Gold	84	96	98.83	99.67	100.83	105.33	101.67	102.83	103.83	108.17
IIHR	72.12	87.65	92.35	93.92	95.38	100.31	95.57	96.58	97.83	102.98
Chandini	72.83	84.78	89.11	92.5	93.67	96.55	94	94.72	95.76	98.89
Subhangini	82.33	95	98.44	99.29	100.39	106.06	100.6	101.48	102.25	108.2
Tiger Flame	89.51	101.31	105.13	105.94	106.84	113.4	108.26	108.97	110.06	116.07
Punjab Dawn	81.16	91.9	95.75	96.7	98.06	101.87	98.39	99.18	100.53	104.95
Punjab Morning	82.95	94.33	98.05	99.19	100.59	103.39	100.33	101.48	102.79	106.23
Danavantri	86.67	96.89	100.15	101.36	102.53	108.11	102.68	103.79	105.07	110.63
Flevo Laguna	90.55	100.58	103.25	104.7	106.03	114.08	105.62	107.43	108.88	116.72
Green Star	83.83	95.17	97.83	98.83	99.83	105.33	101	102	102.33	107.67
Arti	86.11	97.86	101.95	103.25	104.99	110.51	104.26	105.66	107.53	112.96
Chemistry	70.88	85.02	88.35	88.92	90.4	97	91.03	91.67	92.97	99.36
Jyotsna	73.03	83.25	88.89	90	91.92	95.48	92.35	93.13	94.5	98.19
Pusa Shubham	83.44	93.33	96.83	97.83	97.83	104	99.17	100.89	100.89	107
Pusa Srijan	82.13	93.42	95.71	96.67	97.31	101.27	97.92	99.05	99.76	104.13
Surya Kiran	82.4	91.47	94.92	95.7	96.97	102.9	96.92	97.83	99.38	105.62
Yellow Star	91.67	99	102.67	103.33	105	106.67	106	107	108	109.67
Princess Margaret Rose	74.5	86.94	90.61	91.44	92.55	98.25	93.47	94.5	95.36	100.72
White Prosperity	89.28	100.19	103.61	104.89	106.75	110.11	106.53	107.58	109.55	112.78
Darshan	84.25	96.33	100.08	101.25	103.17	109.83	102.33	104.08	105.33	112.25
Sabnam	84.33	94.5	99.17	100.33	102.67	108.33	101.83	103.17	105.5	110.67
S.E.	1.46	1.25	1.3	1.3	1.35	1.25	1.27	1.31	1.36	1.23
C.D. 5%	4.12	3.53	3.69	3.69	3.83	3.55	3.61	3.7	3.84	3.48

**Table 2: Performance of gladiolus varieties for flowering parameters**

	Diameter of 1 <sup>st</sup> floret (cm)	Diameter of 3 <sup>rd</sup> floret (cm)	Diameter of 5 <sup>th</sup> floret (cm)	Diameter of last floret (cm)	Length of 1 <sup>st</sup> floret (cm)	Length of 3 <sup>rd</sup> floret (cm)	Length of 5 <sup>th</sup> floret (cm)	Length of last floret (cm)
Priscilla	9.64	9.56	9.4	8.93	10.34	10.15	10.03	9.81
American Beauty	10.55	10.25	9.71	9.18	10.9	10.75	10.57	9.84
Sunayna	8.62	8.2	7.91	6.59	9.91	9.74	9.4	8.87
Aldebaran	8.98	8.21	8.29	8.13	9.43	8.99	8.98	8.85
Anjali	5.01	4.93	5.21	4.47	8.3	8.09	8.52	7.79
Nova Lux	8.46	7.66	7.81	8.68	10.08	9.95	9.97	10.08
Snow Princess	6.8	7	6.7	6	9.99	9.95	9.95	8.79
Pink Friendship	8.79	8.97	8.79	8.07	9.63	9.95	9.73	9.07
Jester Gold	7.85	8.48	7.73	7.43	9.83	10.07	9.97	9.82
IIHR	9.44	9.36	8.74	8.9	9.9	10.37	9.83	9.44
Chandini	7.37	7.13	7.12	6.11	9.03	9.01	8.83	8.75
Subhangini	5.38	5.47	5.54	5.05	9.38	9.54	9.38	8.83
Tiger Flame	8.35	8.2	8.18	6.87	9.75	9.71	8.86	8.39
Punjab Dawn	7.42	6.97	7.1	6.77	8.76	8.66	8.67	8.4
Punjab Morning	7.13	6.94	6.73	6.36	8.8	8.75	8.54	8.46
Dhanvantari	7.86	7.79	7.67	7.64	10.67	10.58	10.47	9.85
Flevo Laguna	5.02	5.29	5.15	4.98	6.92	7.44	6.8	6.65
Green Star	7.83	8.22	7.56	6.34	8.73	8.87	8.37	7
Arti	6.67	6.6	6.53	6.72	9.6	9.6	9.41	9.03
Chemistry	8.29	7.92	7.68	7.09	9.69	9.75	9.72	8.86
Jyotsna	8.3	8.21	7.63	6.39	9.99	10.18	9.65	9.24
Pusa Shubham	8.21	8.24	7.73	7.47	10.1	9.85	9.82	9.58
Pusa Srijan	8	7.83	7.76	7.58	8.75	8.82	8.7	8.91
Surya Kiran	8.23	8.01	7.59	7.03	10.1	9.57	9.65	9.15
Yellow Star	8.42	8.18	7.57	8.5	10.33	10.17	10.17	10.17
Princess Margaret Rose	7.43	6.88	6.52	6.54	9.67	9.66	9.36	8.52
White Prosperity	7.78	7.43	7.99	6.76	10.01	9.79	9.8	9.11
Darshan	6.77	6.68	6.44	6.6	8.99	8.87	8.79	8.68
Sabnam	7.54	7.65	7.47	6.64	9.37	10.05	9.82	9.23
S.E.	0.31	0.3	0.3	0.35	0.22	0.23	0.27	0.27
C.D. 5%	0.88	0.86	0.86	0.98	0.63	0.65	0.77	0.78

cm) for last floret (Table 2). Whereas, it was minimum for 1<sup>st</sup>, 3<sup>rd</sup> and last florets in variety Anjali (5.01 cm, 4.93 cm and 4.47 cm, respectively) and for 5<sup>th</sup> floret in the variety Flevo Laguna (5.15 cm). The length of 1<sup>st</sup> floret was highest in the variety American Beauty (10.90 cm) which was at par with the variety Dhanvantari (10.67 cm) and Yellow Star (10.33 cm). The length of 3<sup>rd</sup> floret was highest in the variety American Beauty (10.75 cm) which was at par with the variety Dhanvantari (10.58 cm), IIHR (10.37 cm), Jyotsna (10.18 cm), Yellow Star (10.33 cm) and Priscilla (10.15 cm). The length of 5<sup>th</sup> floret was highest in the variety American Beauty (10.57 cm) which was at par with the variety Dhanvantari (10.47 cm), Priscilla (10.03 cm), Nova Lux (9.97 cm), Jester Gold (9.97 cm), Snow Princess (9.95 cm), IIHR (9.83 cm), Pusa Shubham (9.23 cm), Sabnam (9.82 cm) and White Prosperity (9.80 cm). However, the length of last floret was highest in the variety Yellow Star (10.17 cm) which was at par with the variety Nova Lux (10.08 cm), Dhanvantari (9.85 cm), American Beauty (9.84 cm), Jester Gold (9.82 cm), Priscilla (9.81 cm) and Pusa Shubham (9.58 cm). Whereas, it was least in the variety Flevo Laguna (6.92 cm, 7.44 cm, 6.80 cm and 6.65 cm) for 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup> and last florets respectively. Variations in size of the florets were also reported by Chourasia *et al.* (2015), Chaudhary and Dhatt (2017), Singh *et al.* (2018), Sisodia *et al.* (2018) and Nalage *et al.* (2019). The difference in size of the floret in different gladiolus cultivars may be due to their corm vigour which is inherent genetic variability (Nalage *et al.*, 2019).

The maximum number of florets per spike was found in the variety Pink Friendship (16.00) which was at par with varieties Flevo Laguna (14.70), American Beauty (14.29), Snow Princess (14.11) and Priscilla (14.03) whereas, minimum was in the variety Yellow Star (8.33). Number of florets depends on the rachis and inter-nodal length, higher number of florets per spike is preferred for commercial purpose. The maximum number of florets opened at a time was in recorded the variety Princess Margaret Rose (5.95) which was statistically at par with the varieties Yellow Star (5.67), Green Star (5.50), Priscilla (5.34), Pink Friendship (5.7), Jester Gold (5.17), Chemistry (5.17), Snow Princess (5.06), Tiger Flame (4.94), American Beauty (4.90), Sunayna (4.89), Chandini (4.89), Nova Lux (4.83) and White Prosperity (4.83), whereas, the minimum was in the variety Anjali (3.32). The maximum number of florets open per spike was in the variety Pink Friendship (15.00) which was statistically at par with the varieties Flevo Laguna (13.47), American Beauty (12.40) and Darshan (12.17), whereas, minimum was in the variety Yellow Star (7.00). The maximum flowering duration was in the variety Pink Friendship (14.17 days) which was at par with the varieties Flevo Laguna (13.47 days), American Beauty (12.40 days) and Darshan (12.17 days), whereas, the minimum was in Yellow Star (7.00 days). The longest spike was found in the variety IIHR (74.06 cm) which was at par with the variety American Beauty (65.97 cm), whereas, the shortest was in the variety Anjali (36.19 cm). The maximum rachis length was observed in the variety

**Table 3: Performance of gladiolus varieties for flowering parameters**

	Spike length (cm)	Rachis length(cm)	Inter-nodal length (cm) (upto 4 <sup>th</sup> floret)	No. of florets / spike	No. of florets open at a time	No. of florets open / spike	Flowering duration
Priscilla	63.73	44.12	12.13	14.03	5.34	13.03	11.01
American Beauty	65.97	46.35	13.22	14.29	4.9	12.63	12.4
Sunayna	62.85	39.52	12.67	12.94	4.89	12	11.5
Aldebaran	48.73	27.41	11.12	8.44	4.44	7.32	7.4
Anjali	36.19	24.02	9.56	10.25	3.32	9.12	9.83
Nova Lux	47.67	26.92	10.83	10.17	4.83	8.83	7.83
Snow Princess	57.25	43.88	12.49	14.11	5.06	13.01	10.9
Pink Friendship	59.8	44.6	12.04	16	5.17	15	14.17
Jester Gold	61.12	34.82	11.43	11.17	5.17	9.83	9.33
IIHR	74.06	43.73	15.13	12.31	4.42	11.36	10.63
Chandini	58.58	25.96	11.07	9.28	4.89	8.28	9.22
Subhangini	59.01	41.53	12.46	13.37	4.46	12.01	9.84
Tiger Flame	55.71	34.89	11.73	11.9	4.94	10.95	10.94
Punjab Dawn	48.56	28.39	10.73	12.12	4.76	10.99	9.2
Punjab Morning	44.52	22.46	9.86	10	4.1	8.95	8.24
Dhanvantari	57.39	36.81	11.3	12.57	4.31	11.57	10.49
Flevo Laguna	50.6	32.5	9.08	14.7	3.45	13.8	13.47
Green Star	60.67	37.48	13.5	13	5.5	11.83	9.83
Arti	48.85	30.34	9.66	10.66	3.58	9.57	11.01
Chemistry	59.69	39.88	13.06	13.28	5.17	12.28	10.64
Jyotsna	61.8	31.57	11.11	10.92	4.18	9.7	9.52
Pusa Shubham	43.39	27.36	11.72	9.67	4.11	8.67	10.17
Pusa Srijan	60.26	30.78	11.69	10.2	4.42	9.27	8.42
Surya Kiran	50.33	30.38	9.88	12.68	4.15	11.47	10.7
Yellow Star	45.5	23.83	10.33	8.33	5.67	7.33	7
Princess Margaret Rose	53.38	34.97	11.55	12.28	5.95	11.39	10.11
White Prosperity	54.97	32.44	12.76	10.61	4.83	9.39	9.17
Darshan	42.8	27.99	9.49	12.42	3.92	11.42	12.17
Sabnam	57.52	32.83	12.53	11.17	3.67	10.17	11.5
S.E.	3.46	2.72	0.51	0.89	0.3	0.92	0.77
C.D. 5%	9.81	7.71	1.45	2.53	0.86	2.61	2.17

American Beauty (46.35 cm) which was statistically at par with the varieties Pink Friendship (44.60 cm), Priscilla (44.12 cm), Snow Princess (43.88 cm), IIHR (43.73 cm), Chemistry (39.88 cm) and Sunayna (39.52 cm), whereas, the minimum was found in Punjab Morning (22.46 cm). The maximum inter-nodal length (1<sup>st</sup> to 4<sup>th</sup> floret) was observed in the variety IIHR (15.13 cm), whereas, the minimum inter-nodal length was observed in the variety Flevo Laguna (9.08 cm). Variations in the parameters were due to the genetic makeup of the varieties and size of the plants (Sisodia *et al.*, 2018), better vegetative growth increases photosynthetic area thereby leading to better food accumulation and better flowering (Bhaskar *et al.*, 2019). Similar findings for flowering parameters were also found by Singh *et al.* (2013), Reshma *et al.* (2016), Singh *et al.* (2017) and Swaroop *et al.* (2019) in gladiolus, Maurya *et al.* (2015) and Pal *et al.* (2018) in balsam, Singh *et al.* (2013) in snapdragon and Kumar *et al.* (2014) in tuberose.

## REFERENCES

**Bhaskar, V.V., Dorajeero, A.V.D., Suseela, T. and Salma, Z. 2019.** Performance of gladiolus cultivars under the coastal tropical humid climatic conditions of Andhra Pradesh. *J. Pharmacognosy and Phytochemistry*. **8(4)**: 1882-1885.

**Chaudhary, N. and Dhatt, K.K. 2017.** Varietal response of gladiolus to planting time for off season flower production. *The Bioscan*. **12(4)**: 1975-1980.

**Chourasia, A., Viradia, R.R., Ansar, H. and Madle, S.N. 2015.** Evaluation of different gladiolus cultivars for growth, flowering, spike yield and corm yield under Saurashtra region of Gujarat. *The Bioscan*. **10(1)**:131-134.

**Kadam, G.B., Kumar, G., Saha, T.N., Tiwari, A.K. and Kumar, R. 2014.** Varietal evaluation and genetic variability studies on gladiolus. *Indian J. Horticulture*. **71(3)**: 379-384.

**Kaur, H. and Bajpay, A. 2019.** Performance of various Gladiolus cultivars under Punjab conditions. *J. Pharmacognosy and Phytochemistry*. **8(4)**: 875-878.

**Kumar, A., Sisodia, A. and Singh, A.K. 2014.** Evaluation of tuberose cultivars for growth, flowering and post-harvest life. *Indian Perfumer*. **58(1)**: 29-32.

**Kumar, K.P., Sarkar, A., Chowdhuri, T.K. and Sadhukan, R. 2019.** Performance of new gladiolus cultivars under the Gangetic plateau of West Bengal conditions. *J. Pharmacognosy and Phytochemistry*. **8(3)**: 444-447.

**Maurya, J.K., Pal, A.K., Singh, B.K., Singh, S.V. and Singh, A.K. 2015.** Evaluation of different type single whorled germplasm of balsam (*Impatiens balsamina* L.). *Indian J. Agriculture and Allied Sciences*. **1(3)**: 44-47.

**Mushtaq, S., Hafiz, I.A., Arif, M. and Anwar, A. 2018.** Performance evaluation of elite gladiolus cultivars under agro-climatic conditions of Rawalpindi. *Asian J. Advances in Agricultural Research*. **5(3)**: 1-6.

**Pal, S., Singh, A.K., Sisodia, A., Pal, A.K. and Tiwari, A. 2018.** Evaluation of double whorled balsam (*Impatiens balsamina* L.) genotypes for growth, flowering and seed attributes. *J. Pharmacognosy and Phytochemistry*. **7(2)**: 2901-2904.

- Panse, V.G. and Sukhatme, P.V. 1985.** Statistical Methods for Agricultural Workers. Indian Council of Agricultural Research, New Delhi. p. 158.
- Rashmi, R. and Chandrashekar, S.Y. 2016.** Evaluation of genetic variability, heritability and genetic advances in gladiolus (*Gladiolus hybridus* L.) genotypes. *The Bioscan*. **11(3)**: 1829-1832.
- Reshma, V.S., Panchbhai, D.M. and Nagar, D. 2016.** Performance of five different gladiolus varieties with respect to growth, flowering and yield attributes under Nagpur condition. *The Ecoscan*. **9**: 561-565.
- Singh, A.K. 2006.** Flower crops: Cultivation and management. Vol-1. New India Publishing Agency, New Delhi. p. 147.
- Singh, A.K. 2014.** Breeding and Biotechnology of Flowers: Commercial Flowers. Vol-1. New India Publishing Agency, New Delhi. p. 705.
- Singh, A.K. and Sisodia, A. 2017.** Textbook of Floriculture and Landscaping. New India Publishing Agency, New Delhi. p. 432.
- Singh, A.K., Jauhari, S. and Sisodia, A. 2013.** Development of snapdragon hybrids and their evaluation for vegetative, flowering and seed attributes. *Progressive Horticulture*. **46(1)**: 133-142.
- Singh, A.K., Kumar, A. and Ghimire, N.R. 2013.** Performance of Indian and exotic varieties of gladiolus under eastern Uttar Pradesh conditions. *Asian J. Horticulture*. **8(1)**: 191-194.
- Singh, A.K., Kumar, D., Sisodia, A. and Padhi, M. 2018.** Varietal evaluation of gladiolus for flowering attributes under Indo-Gangetic plains. *International J. Current Microbiology and Applied Sciences*. **7(7)**: 3627-3669.
- Singh, A.K., Sisodia, A., Sisodia, V. and Ray, P. 2017.** Performance of Indian and exotic varieties of gladiolus under eastern U.P. conditions. *J. Ornamental Horticulture*. **20(3-4)**: 153-157.
- Sisodia, A., Kumawat, P. and Singh, A.K. 2018.** Performance of gladiolus varieties for flowering traits. *J. Pharmacognosy and Phytochemistry*. **7(3)**: 3383-3386.
- Sisodia, A., Pal, A. and Singh, A.K. 2018.** Varietal response and effect of *Trichoderma* on flowering in gladiolus. *J. Pharmacognosy and Photochemistry*. **7(3)**: 3658-3660.
- Swaroop, K., Singh, K.P. and Kumar, P. 2019.** Evaluation of gladiolus (*Gladiolus grandiflora*) genotypes for morphological diversity and corm yield. *Current Horticulture*. **7(2)**: 48-51.
- Nalage, N.A., Haldankar, P.M., Gawankar, M.S. and Rathod, N.G. 2019.** Evaluation of different gladiolus varieties (*Gladiolus hybridus* Hort.) under Konkan conditions of Maharashtra. *International J. Chemical Studies*. **7(2)**: 2018-2021.

---



**NATIONAL ENVIRONMENTALISTS  
ASSOCIATION**

The National Environmentalists Association is chartered in Ranchi as a nonprofit scientific and educational association of like minded academicians, researchers, scientists from all over the nation for the furtherance and diffusion of knowledge of Life Sciences in general and Environmental Science in particular.

The association not only honours its members but also provides FELLOWSHIP to outstanding contributors to the subject and the society.

**Contact :**

**For Editorial Information**

Prof. M. P. Sinha  
Vice Chancellor  
Sido Kanhu Murmu University  
Dumka - 814 110  
Jharkhand, INDIA

**For information regarding Association :**

SECRETARY,  
National Environmentalists Association,  
D-13, Sai Roofs, 1<sup>st</sup> Floor,  
H. H. Colony,  
Ranchi - 834002  
Jharkhand, India

E-mails : editor.bioscan@gmail.com  
dr.mp.sinha@gmail.com  
nat.env.assoc@gmail.com

Cell : 94313-60645; 9572649448

Ph. : 0651-2244071

Website : www.thebioscan.com  
: www.neaindia.org

**NAAS Rating : 5.26**

**U.S.A. Office**

2827 Videre Dr.,  
Wilmington,  
DE 19808 We, USA

Type setter Bandana Solutions Facility Management LLP  
Published by Aditi Publications, Patliputra, Patna

# The Bioscan

An International Quarterly Journal of Life Sciences

ISSN : 0973-7049

Volume 15(3) : 2020

Published as an official organ by

**NATIONAL ENVIRONMENTALISTS ASSOCIATION**

## CONTENTS

	Page
<b>A. RESEARCH PAPER</b>	
1. Comparison and molecular profiling of Begomovirus infecting chilli ( <i>Capsicum annum</i> ) in gangetic alluvial zone of West Bengal Uday Bikash Oraono, Lourembam Sanajaoba Singh and Jayanta Tarafdar	275 - 280
2. Assessment of probiotic characteristics of <i>L.plantarum</i> Sravani Kandula And Rita Narayanan	281 - 286
3. Effect of rate and frequency of micronutrient on growth attributes and dry matter yield of Banana Cv. grand naine under south Gujarat condition Narendra Singh, Sonal Tripathi, Patel V. A., Jaimin Naik and Chauhan Aditi	287 - 290
4. Character association and path analysis for seed yield and its components in Grass pea ( <i>Lathyrus sativus</i> L.) Gangishetti Ranjithkumar, Sandip Debnath and Duddukur Rajasekhar	291 - 295
5. Determination of physical and biometric properties of onion bulbs in relation to design of digger cum windrower Shiddanagouda Yadachi and Kiran Nagajjanavar	297 - 301
6. Polygenic variation for morphological and biochemical traits of brinjal genotypes ( <i>Solanum melongena</i> L.) and its wild relatives Nisha Sharma, K. D. Bhutia, Rajesh Kumar, Sita Kumari Prasad, Ankita Debnath and Malay Marut Sharma	303 - 309
7. Effect of different sources of horizontal transmission of bacterial flacherie on Et <sub>50</sub> for symptom expression and mortality of PM X CSR <sub>2</sub> B. L. Kavyashree., R. N. Bhaskar and C. Doreswamy	311 - 314
8. Comparative biology of <i>Goniozus nephantidis</i> (Muesbeck) on <i>Galleria mellonella</i> L. and <i>Corcyra cephalonica</i> (Stainton) A. V. Desai, M. R. Siddhapara and N. P. Trivedi	315 - 321
9. Efficiency of ovatide on mass seed production of climbing perch ( <i>Anabas testudineus</i> , Bloch, 1972) in Nalbari district, Assam Ankur Rajbongshi, A. Ali, M. Chakravarty, M. Deka, H. Mazumdar, Pranab Kr Das and S. Baishya	323 - 326
10. Study of combining ability and gene action for yield and yield component characters in interspecific hybrids of cotton ( <i>Gossypium hirsutum</i> L. X <i>Gossypium barbadense</i> L.) S. B. Gohil., M. B. Parmar., M. P. Patel and D. A. Patel	327 - 333
11. Per oral inoculation of <i>Lysinibacillus sphaericus</i> with pathogenic microbes on rearing and cocoon parameters of silkworm, <i>Bombyx mori</i> L. H. G. Anusha, R. N. Bhaskar and K. V. Anitharani	335 - 338
12. Effect of fungicides, plant extracts and bioagents on spore germination of <i>Colletotrichum lindemuthianum</i> causing field bean anthracnose	

**The Journal is Currently Abstracted / Indexed in**

- Paryavarn Abstract, INDIA
- Indian Science Abstract, INDIA
- Cambridge Science Abstract, U.S.A.
- Zoological Record, U.K.
- Directory of Open Access Journal (DOAJ)
- Chemical Abstract, U. S. A.
- Research BIB
- Indian Science
- Journal Seek
- Scientific Indexing Service (SIS)
- Journal is currently rated by
- Index Copernicus
- Universal Impact Factor
- NAAS

**DISCLAIMER**

The Publisher and Editors cannot be held responsible for errors or any consequences arising from the use of information in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher/ Association and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher / Association and Editors of the products advertised.

- S. Narasimha Rao, S.L. Bhattiprolu, A. Vijaya Gopal and V. Sekhar— 339-344
13. Evaluation in vitro different fungicides for growth of *Rhizoctonia bataticola*  
A.M. Kadam, S.S. Chavan and A.H. Kendre — 345- 349
  14. Evaluation of gladiolus varieties for flowering and cut flower traits under indo-gangetic plains  
Girish, P. M., Anjana Sisodia and Anil K. Singh — 351- 355
  15. Effect of land configuration and different organic sources on growth, yield and quality of carrot under organic farming  
B. Solanki, A. R. Kaswala, P.K. Dubey and A.P. Italiya— 357- 362
  16. Verification and usability analysis of medium range weather forecast for the Kokrajhar district of lower Brahmaputra valley zone of Assam  
Kuldip Medhi, Kushal Sarmah, Vinod Upadhyay, Sunil Kumar Paul, Athar N. Islam and Bikash J. Gharphalia— 363 - 370
  17. Canonical root analysis and clustering for characterization and evaluation of aromatic rice germplasm based on morphological characters  
G. Parimala, Ch. Damodhar Raju, L.V. Subba Rao and K. Uma Maheswari— 371 - 374
  18. Genetic divergence analysis of sesame genotypes (*Sesamum indicum* L.)  
Dasari Rajitha, T. Srikanth, D. Padmaja and T. Kiran Babu— 375 - 379
  19. Nutrient uptake and chemical properties of soil after harvest of baby corn (*Zea mays* L.) as influenced by organic manures and fertilizers  
D.H. Roopashree, S. Kamal Bai, Nagaraju and S. Raghavendra— 381 - 384
  20. Genotypic response on growth and yield in papaya  
D. K. Varu., K. D. Patel and Sandip Makhmale— 385 - 389
  21. Studies on frequency distribution of yield and yield related traits in F<sub>2</sub>M<sub>2</sub> generation of sesame (*Sesamum indicum* L.)  
Rajesh Kumar Kar, Tapash Kumar Mishra and Banshidhar Pradhan — 391 - 395
  22. Correlation and path analysis in cowpea (*Vigna unguiculata* (L.) Walp)  
R.M. Nagalakshmi, R. Usha Kumari and R. Ananda Kumar— 397 - 401
  23. A simple and efficient method for DNA extraction from rabi sorghum [*Sorghum bicolor* (L.) Moench]"  
S. S. Gadakh, G. D. Khalekar, U. S. Dalvi, A. A. Kale and P.L. Kulwal— 403 - 406
  24. Determination of economic injury level (EIL) of sugarcane planthopper borer *Chilo tumidicostalis* Hampson (Lepidoptera: pyralidae)  
R. K. Nath and D. K. Saikia— 407- 409
  25. Performance of different summer mung (*Vigna radiata* L.) varieties sown at different dates under Manipur valley condition  
Meghna Gogoi, Jamkhogin Lhungdim, Kamal Kant, Urjashi Bhattacharya and Gauri Mohan — 411 - 414





### ZONAL CO-ORDINATORS OF THE ASSOCIATION

- Prof. N. Behera  
School of Life Science,  
Sambalpur University
- Dr. Nirmal Kumar  
ISTAR, Vallabh Vidyanagar,  
Anand, Gujarat
- Dr. P. N. Sudha  
D. K. M. College for Women,  
Vellore
- Prof. S. P. S. Dutta  
Dept. of Environmental Science,  
Jammu University, Jammu
- Dr. V. Salom Gnana Thanga  
Dept. of Env. Scs.,  
University of Kerala, Kariavattom  
Tiruvananthapuram, Kerala

### Publications of the Association

#### The Bioscan

An International Quarterly Journal of Life Sciences

#### The Ecoscan

An International Quarterly Journal of  
Environmental Sciences

**Both the Journals are  
online**

**Both the Journals are  
available on Google.com**

**Websites of the Journals are**  
[www.theecoscan.in](http://www.theecoscan.in)  
[www.thebioscan.in](http://www.thebioscan.in)

### FEATURES OF ASSOCIATION

- Association is registered under 80G of I.T.
- Prestigious fellowship of the Association (F. N. E. A.) to academicians of the nation.
- Regular annual conference of national and international levels organized by the Association.
- Young Scientist and Senior Scientist award during the conference of the Association.



\_\_\_\_\_ 062 & 082

\_\_\_\_\_ 000 & 000

<b>THE BIOSCAN : SUBSCRIPTION RATES</b>				
		India (Rs.)	SAARC Countries	Other Countries
Individuals	One Year	1,000	2,000(I:C)	US \$200
	Life Member*	10,000		
Institutions	One Year	3,000	6,000(I:C)	US \$400
	Life Member*	30,000		

\*Life Member will receive the journal for 15 years while other benefits will continue whole life

### THE BIOSCAN : MEMBERSHIP FORM

Please enter my subscription for the above journal for the year ..... / life member.

Name: .....

Address: .....

E-mail: .....

Payment Rs. : ..... by DD / MD in favour of  
National Environmentalists Association payable at Ranchi, No. .... Dated  
..... is enclosed.

**NOTE: FOR MEMBERSHIP THE ABOVE INFORMATION CAN BE SENT ON SEPARATE SHEET**