

EFFECT OF LAND CONFIGURATION AND DIFFERENT ORGANIC SOURCES ON GROWTH, YIELD AND QUALITY OF CARROT UNDER ORGANIC FARMING

B. SOLANKI¹, A. R. KASWALA^{2*}, P.K. DUBEY² AND A.P. ITALIYA²

¹Department of Soil Science and Agricultural Chemistry, N. M. Collage of Agriculture,

²Department of Natural Resource Management, ASPEE Collage of Horticulture and Forestry
Navsari Agricultural University, Navsari - 396 450, Gujarat, INDIA

e-mail: anandnau@gmail.com

KEYWORDS

Carrot
Organic Farming
Land configuration
Growth

Received on :

21.06.2020

Accepted on :

28.08.2020

*Corresponding
author

ABSTRACT

A field experiment was conducted at the certified organic farm, Navsari Agricultural University, Navsari during Rabi season of 2017-2018 to study the effect of land configuration, fertilizer level and liquid formulation on growth, yield and quality of carrot under organic farming. The treatments imposed were three levels of land configuration *i.e.* C₁: Flatbed C₂: Ridge and furrow and C₃: Broad bed, three levels of fertilizer *i.e.* F₁: 100% N through vermicompost F₂: 75% N through vermicompost, and F₃: 50% N through vermicompost and two levels of liquid formulation *i.e.* L₁: Jeevamrut and L₂: Amritpani in FRBD which replicated thrice. Overall the growth, yield attributes, yield and quality parameters were significantly highest with individual treatments C₃, F₁ and L₁ in all the cases. Significantly maximum root yield of carrot was achieved when 75% N was supplied in broad bed which was at par with 100% N application in all the treatments of land configuration. However root volume, root girth and root length was also affected significantly with the interaction effect of CxL and FxL. Further CxL interaction was also significant with shoot length while FxL interaction was found significant with quality parameters *i.e.* carotene and TSS content.

INTRODUCTION

Nowadays, the indiscriminate use of inorganic fertilizers are producing very hazardous effect on soil properties as well as crop yield. Therefore, it is essential to utilize various sources of nutrients, particularly under organic farming in order to increase the production of crop by maintaining soil fertility and quality. This can partly be accomplished through the adoption of good management techniques. Among them, land management system involving different methods of seed bed preparation plays a crucial role in enhancing crop production through improving soil-water-plant relationship. Ridge-furrow and bed-furrow land configuration systems emerge as few of the most promising sustainable management technologies which increase input use efficiency and crop production (Yadav *et al.*, 2003). Organic manures originate from both livestock waste and crop residues, with the nutrients in them being mineralized by soil microbes and slowly making them available to plants over a long period of time (Lampkin, 2000). Humus added by organic manures adsorbs large quantities of water and makes it available to plants. The organic matter activates the soil ingredients necessary for a plants healthy growth. It has a very complex effect on soil and plant growth as well as it improves the physical, chemical and biological properties of soil. Liquid formulations that are used in organic agriculture are the fermented products which are used as plant growth enhancing substances prepared from farm available material. They are rich sources of beneficial micro flora which

support, stimulate the plant growth and helping in getting better vegetative growth and also good quality yield (Devkumar *et al.*, 2014). With this view, an experiment was conducted to study the effect of land configuration, fertilizer level and liquid formulation on growth, yield and quality of carrot under organic farming.

MATERIALS AND METHODS

The experiment was laid out on carrot as a test crop in rabi season in Factorial Randomized Block Design with three replication during 2017-18 at Organic Farm, Navsari Agricultural University, Navsari, Gujarat, India. Experimental soil was clayey in texture, non-saline (EC-0.81 dS/m) and slightly alkaline (pH- 8.1) in nature, available nitrogen, phosphorus and potassium was high (284 kg/ha), medium (50 kg/ha) and high (482 kg/ha), respectively. There were total eighteen treatment combinations comprising from three land configuration (C₁- Flat Bed, C₂- Ridges and Furrow and C₃- Broad Bed), three nitrogen level (F₁- 100% N through Vermicompost, F₂- 75% N through Vermicompost and F₃- 50% N through Vermicompost) and application of two liquid formulations (L₁- Jeevamrut @ 600 l/ha and L₂- Amrutpani @ 600 l/ha). Seed was treated with each of 0.5% solution of *Trichoderma viride* and *Pseudomonas fluorescencesas* as a precautionary measure to prevent soil borne diseases. For fertilizing the crop 50% nitrogen was applied at basal and

remaining 50% nitrogen was applied at 30 days after sowing (DAS) through vermicompost whereas, liquid formulation was applied at 30, 45 and 60 DAS. The liquid formulations were prepared as per the method suggested by National Centre for Organic Farming, Ghaziabad. Jeevamrut: Mix cow dung 10 kg, cow urine 10 lit, Jaggary 2 kg, pulse grain flour 2 kg and live forest soil 1 kg in 200 lit of water. Ferment for 7 days. Stir the solution regularly three times a day. Amritpani: Mix 10 kg cow dung with 500 gm honey and mix thoroughly to form a creamy paste. Add 250 gm of cow desi ghee and mix at high speed and dilute with 200 lit. of water.

RESULTS AND DISCUSSION

Growth, yield attributes and yield

Data regarding effect of land configuration, fertilizer level and liquid formulation on plant height, root volume and root girth at harvest are presented in Table 1.

The effect of land configuration did not show any significant effect on the plant height at harvest of carrot (Table 1). Slightly higher height was recorded in the broad bed treatment, but it did not reach to the level of significance. The treatment C₃ (Broad bed) recorded significantly higher root volume and root girth of 83.29 cm³ and 3.25 cm, respectively but it remained at par with the treatment C₂ (Ridges and Furrow) while the lowest root volume and root girth was observed under treatment C₁ (Flatbed). In the broad bed and ridges and furrows system soil remains soft and moist, which ultimately resulted in increased moisture and nutrient availability to the

Table 1: Effect of land configuration, fertilizer level, liquid formulation and their interaction effect on plant height, root volume and root girth of carrot.

Treatments	Plant height (cm)	Root volume (cm ³)	Root girth (cm)
Land Configuration (C)			
C ₁ - Flat Bed	74.54	75.46	2.94
C ₂ - Ridges	76.44	81.24	3.13
C ₃ - Broad Bed	77.18	83.39	3.25
SEm ±	1.31	1.36	0.06
CD at 5%	NS	3.91	0.16
Fertilizer Level (F)			
F ₁ - 100 % N	78.6	82.56	3.22
F ₂ - 75 % N	75.72	80.58	3.15
F ₃ - 50 % N	73.85	76.94	2.95
SEm ±	1.32	1.36	0.06
CD at 5%	3.78	3.91	0.16
Liquid Formulation (L)			
L ₁ - Jeevamrut	77.83	81.83	3.18
L ₂ - Amritpani	74.28	78.23	3.04
SEm ±	1.07	1.1	0.06
CD at 5%	3.09	3.19	0.16
Interaction			
C×F SEm ±	2.28	2.35	0.1
CD at 5%	NS	6.77	0.23
C×L SEm ±	1.86	1.92	0.08
CD at 5%	NS	NS	NS
F × L SEm ±	1.86	1.92	0.08
CD at 5%	NS	5.53	0.23
C×F×L SEm ±	3.22	3.33	0.14
CD at 5%	NS	NS	NS
CV (%)	7.3	7.21	7.6

plants compared to the flat bed system (Mengel *et al.*, 2001). Lower yield attributes with flat bed may be due to less retention of water resulting in insufficient water level in the root zone during the crop growth period.

The results on growth and yield attributes revealed that among the different levels of fertilizer, F₁ (100% N through vermicompost) showed significantly higher plant height, root volume and root girth of 78.60 cm, 82.56 cm³ and 3.22 cm, respectively but remained at par with the treatment F₂ (75 % N through vermicompost) with value of 80.58 cm³ and 2.95 cm in the case of root volume and root girth at harvest. Significantly lowest value of plant height, root volume and root girth was recorded under the treatment F₃ i.e. 50 % vermicompost (Table 1). Addition of vermicompost not only enhanced the nutrients availability by producing gibberellin, cytokine and auxins resulting in uptake of nutrients that may have favoured plant growth and yield, and it also improve the soil structure and maintain moisture availability. Similarly, when chickpea crop was fertilized with 100% N through biocompost:vermicompost:castorcake and additionally foliar application cow urine (2%) received maximum grain and stover yield under south Gujarat condition (Bag *et al.*, 2015). According to Arancon *et al.* (2003) vermicompost combination with vermiwash applied at very low rates (2.5 to 5.0 t ha⁻¹) can significantly enhanced the growth parameter such as plant height, yield etc. The similar result in carrot was also reported by Kirad *et al.* (2010) and Mazed *et al.* (2015) and, Patel *et al.* (2016) in onion.

With respect to the treatment of liquid formulations more plant height (77.83 cm), root volume (81.83 cm³) and root girth (3.18 cm) was observed under L₁ (Jeevamrut) at harvest compared to L₂ (Table 1). The organic formulations activate the biological reactions in the soil and to protect the plants from diseases incidence which also become responsible to enhance the growth and crop yield. Similarly, Gore and Sreenivasa (2011) also reported highest plant growth and root length as well as yield of tomato with the application of RDF + Beejamruth + Jeevamruth + Panchgavya. These also agree with Vinaykumar and Neeraj (2015) and Sankar *et al.* (2009) who used panchgavya as liquid fertilizer in onion crop.

The results revealed that interaction of land configuration, fertilizer level and liquid formulation *viz.*, CxF, CxL, FxL and CxFxL were failed to exert any significant effect on plant height. Among the various two factor interactions, interaction between CxL was found non significant, whereas remaining interactions were significant. Interaction of CxF showed significant effect on root volume (Table 2). The combination of C₃F₁ produced significantly higher root volume (89.53 cm³) and root girth (3.47 cm) which was remained at par with treatment combination C₃F₂ (86.22 cm³ and 3.33 cm of root volume and root girth, respectively). The significantly lower root volume (74.43 cm³) was recorded under combination of C₃F₃ while significantly lower root girth was recorded with the treatment combination C₁F₃ (2.75 cm), which was at par with C₂F₁ combination.

Interaction between FxL also showed a significant effect on the root volume and root girth (Table 2 and 3) of carrot. The combination of F₂L₁ gave significantly higher root volume (85.83 cm³) but it was remained at par with the treatment

Table 2: Interaction effect of C x F and F x L on root volume (cm³) of carrot.

Land Configuration (C)	Fertilizer level (F)			Liquid formulation (L)	
	F ₁ - 100 % N	F ₂ - 75 % N	F ₃ - 50 % N	L ₁ - Jeevamrut	L ₂ - Amritpani
C ₁ - Flat Bed	75.13	75.67	75.57	F ₁ - 100 % N	84.66
C ₂ - Ridges	83.02	79.87	80.83	F ₂ - 75 % N	85.83
C ₃ - Broad Bed	89.53	86.22	74.43	F ₃ - 50 % N	74.99

Table 3 : Interaction effect of C x F and F x L on root girth (cm) of carrot

Land Configuration (C)	Fertilizer level (F)			Liquid formulation (L)	
	F ₁ - 100 % N	F ₂ - 75 % N	F ₃ - 50 % N	L ₁ - Jeevamrut	L ₂ - Amritpani
C ₁ - Flat Bed	3.17	2.9	2.75	F ₁ - 100 % N	3.41
C ₂ - Ridges	3.02	3.33	3.05	F ₂ - 75 % N	3.13
C ₃ - Broad Bed	3.47	3.23	3.05	F ₃ - 50 % N	3

Table 4: Effect of land configuration, fertilizer level, liquid and their interaction effect formulation on root length, shoot length and yield of carrot.

Treatments	Root length (cm)	Shoot length (cm)	Yield (t/ha)
Land Configuration (C)			
C ₁ - Flat Bed	16.68	76.39	9.92
C ₂ - Ridges	17.87	80.13	10.74
C ₃ - Broad Bed	19.85	79.66	11.14
S _{Em} ±	0.42	1.11	0.26
CD at 5%	1.21	3.19	0.74
Fertilizer Level (F)			
F ₁ - 100 % N	20.41	80.69	11.24
F ₂ - 75 % N	17.46	79.22	10.61
F ₃ - 50 % N	16.52	76.26	9.94
S _{Em} ±	0.42	1.11	0.26
CD at 5%	1.21	3.19	0.74
Liquid Formulation (L)			
L ₁ - Jeevamrut	18.48	78.88	11.19
L ₂ - Amritpani	17.78	78.57	10.01
S _{Em} ±	0.34	0.9	0.21
CD at 5%	NS	NS	0.6
Interaction			
C×F S _{Em} ±	0.72	1.92	0.45
CD at 5%	2.09	5.53	1.29
C×L S _{Em} ±	0.59	1.57	0.37
CD at 5%	NS	NS	NS
F × L S _{Em} ±	0.59	1.57	0.37
CD at 5%	1.71	NS	1.05
C×F×L S _{Em} ±	1.03	2.72	0.63
CD at 5%	NS	NS	NS
CV (%)	9.8	5.98	10.41

combinations of F₁L₁ (84.66 cm³) and F₁L₂ (80.47 cm³). The significantly lower root volume (74.99 cm³) was recorded under combination F₃L₁ however; it remained at par with F₃L₂ and F₁L₂ treatment combinations. The higher order interaction was found non significant. In the case with root girth F₁L₁ resulted in significantly higher value (3.41 cm) but it remained at par with F₂L₁ (3.13 cm) and F₃L₁ (3.00 cm). The significantly lower root girth (2.91 cm) was recorded with combination of F₃L₂ which was statistically similar to F₂L₁ combination.

The results were also in agreement with Shwetha and Babalad (2008) and Chandrakala (2008) when they have applied organic manures in combination with fermented organics *viz.*, Beejamrut, Jeevamrut, Panchgavya to soybean-wheat cropping system and observed significantly higher leaf area index, plant height, number of branches, dry matter accumulation, seed yield and yield parameters like number of pods per plant over

organics alone application.

Data regarding effect of land configuration, fertilizer level and liquid formulation on root length, shoot length and yield at harvest are presented in Table 4.

The mean root length was significantly affected due to land configuration. The treatment C₃ (Broad bed) recorded significantly superior root length (19.85 cm) whereas the lowest root length of 16.68 cm was recorded with the treatment C₁ (Flat bed), but the values under C₁ and C₂ were at par. Sowing carrot on ridges (C₂) enhanced the shoot length (80.13 cm) followed by C₃- broad bed (79.65 cm). Flat bed (C₁) type of configuration recorded smaller shoot length. The results of land configuration significantly affected on root weight of carrot and maximum root yield (11.14 t/ha) was observed under C₃ (Broad bed) treatment. It was significantly higher than C₁ (Flat bed) treatment (9.92 t/ha) but was similar (10.74 t/ha) to C₂ (Ridges and furrow). The increase in yield attributes with broad bed over ridges and furrows could be attributed to loose friable soil, improved physical properties such as lower bulk density, better aeration and lower penetration resistance. (Mengel *et al.*, 2001).

The different treatment of vermicompost application was found significant on the root length, shoot length and carrot yield (Table 4). The fertilizer level F₁- 100% N through vermicompost recorded significantly highest root length (20.41 cm) as compared to other levels. It was found significantly superior over other treatments. The lowest root length (16.52 cm) was obtained with F₃ (50% N through vermicompost) however, it was at par with F₂ (50% N through vermicompost). The different levels of vermicompost application also exerted significant effect on the shoot length of carrot. The significantly highest shoot length (80.69 cm) was recorded with the treatment F₁ but remain at par with the treatment F₂ and lowest shoot length (76.26 cm) was recorded with the treatment F₃. Application of N through vermicompost at different rates was affected significantly on the root yield of carrot. The values of root yield obtained at 100%, 75% and 50% N application through vermicompost were 11.24, 10.61 and 9.94 t/ha respectively. Among which values of F₁ and F₂ were at par. The ability of organic manure to improve the chemical properties of soil as well as its release its nutrient in to the soil, which make it an ideal input for good carrot crop yield. Ahmed *et al.* (2014) found the similar result in carrot, Kirad *et al.* (2010) in carrot, Mazed *et al.* (2015) in carrot. Similar results were also obtained by Gadelrab and ELAmin (2013) when they have applied

Table 5: Interaction effect of C x F and F X L on root length (cm) of carrot.

Land Configuration (C)	Fertilizer level (F)			Liquid formulation (L)	
	F ₁ - 100 % N	F ₂ - 75 % N	F ₃ - 50 % N	L ₁ -Jeevamrut	L ₂ - Amritpani
C ₁ - Flat Bed	17.11	16.28	16.63	20.94	19.87
C ₂ - Ridges	21.57	16.43	15.6	19.46	15.45
C ₃ - Broad Bed	22.54	19.67	17.34	15.04	18.01

Table 6: Interaction effect of C x F on shoot length (cm) of carrot.

Land Configuration (C)	Fertilizer Level (F)		
	F ₁ - 100 % N	F ₂ - 75 % N	F ₃ - 50 % N
C ₁ - Flat Bed	78.7	74.34	76.13
C ₂ - Ridges	81.03	79.97	79.38
C ₃ - Broad Bed	82.33	83.37	73.28

Table 7: Interaction effect of C x F and F x L on carrot yield (t/ha).

Land Configuration (C)	Fertilizer level (F)			Liquid formulation (L)	
	F ₁ - 100 % N	F ₂ - 75 % N	F ₃ - 50 % N	L ₁ -Jeevamrut	L ₂ - Amritpani
C ₁ - Flat Bed	10.9	9.47	9.4	12.54	9.95
C ₂ - Ridges	11.56	10.24	10.41	10.85	10.37
C ₃ - Broad Bed	11.28	12.12	10.01	10.16	9.71

Table 8: Effect of land configuration, fertilizer level, liquid formulation and their interaction effect on carotene, TSS and carbohydrate content of carrot.

Treatments	Carotene (mg/100 g)	TSS (°brix)	Carbohydrate (%)
Land Configuration (C)			
C ₁ - Flat Bed	5.04	14.03	8.16
C ₂ - Ridges	5.05	14.04	8.18
C ₃ - Broad Bed	5.07	14.11	8.21
SEm ±	0.05	0.05	0.03
CD at 5%	NS	NS	NS
Fertilizer Level (F)			
F ₁ - 100 % N	5.13	14.15	8.2
F ₂ - 75 % N	5.09	14.1	8.19
F ₃ - 50 % N	4.93	13.93	8.16
SEm ±	0.05	0.05	0.03
CD at 5%	0.14	0.14	NS
Liquid Formulation (L)			
L ₁ -Jeevamrut	5.11	14.12	8.19
L ₂ - Amritpani	4.99	14	8.18
SEm ±	0.03	0.04	0.03
CD at 5%	0.11	0.11	NS
Interaction			
C×F SEm ±	0.08	0.07	0.06
CD at 5%	NS	NS	NS
C×L SEm ±	0.07	0.07	0.05
CD at 5%	NS	NS	NS
F × L SEm ±	0.07	0.08	0.05
CD at 5%	0.2	0.24	NS
C×F×L SEm ±	0.12	0.12	0.08
CD at 5%	NS	NS	NS
CV (%)	4.1	1.51	1.7

compost Elshomokh and compost Alkhaseeb on onion.

The effect of liquid formulation was found non-significant and the maximum root length (18.48 cm) of carrot was observed with L₁ (Jeevamrut) and minimum (17.78 cm) with L₂ (Amritpani). However the different liquid formulation did not showed any significant response on root length. In case with root yield the significantly maximum yield (11.19 t/ha) was obtained with L₁ (Jeevamrut) treatment whereas L₂ (Amritpani) yielded 10.01 t/ha of carrot yield (Table 4). The increase in

yield might be due to application of microorganisms enriched organic sources which may create maximum nutrient availability to plant. Patil *et al.* (2012) also found highest grain and straw yield of soybean when they have applied 100 per cent RDN through vermicompost + jeevamrut which was statistically at par with the application of 100 per cent RDN through FYM + jeevamrut.

The interaction effect of land configuration and fertilizer level (CxL) as well as fertilizer level x land configuration (FxL) resulted significant effect on root length (Table 5). The treatment combination C₃F₁ registered significantly higher root length (22.54 cm) but did not differ statistically with C₂F₁. The lowest root length was recorded with C₂F₃ combination (15.60 cm). Another second order interaction of FxL was also showed significant effect on root length (Table 5). From interaction point of view the combinations F₁L₁, F₁L₂ and F₂L₁ maintain their superiority in root length while, the lowest value of 15.04 cm of root length was recorded with combination of F₃L₁. It was at par with F₂L₂ combination.

The interaction of land configuration and fertilizer level (CxL) recorded significant effect on shoot length (Table 6) and the treatment combination of C₃F₂ recorded significantly higher shoot length (83.36 cm), but remained at par with C₂F₁, C₃F₁, C₂F₂ and C₂F₃. The other interactions CxL, FxL and CxFxL on shoot length were failed to reach the level of significance.

Similar to root length of carrot, the interaction effect of CxL and FxL were also found significant with root yield of carrot. The result presented in Table 7 reflected that the C₃F₂ combination yielded significantly highest (12.12 t/ha) carrot yield which remained at par with C₂F₁ and C₃F₁. The lowest yield (9.40 t/ha) was obtained with C₁F₃ however, it was statistically at par with combination of C₁F₂, C₂F₂ and C₂F₃ with value of 9.47, 10.24 and 10.41 t/ha, respectively. Interaction of FxL significantly yielded highest (12.54 t/ha) carrot root under combination of F₁L₁ which was significantly superior over other combinations as remaining combinations yielded significantly lower root and showing no difference between them.

Table 9: Interaction effect of F x L on carotene and TSS content of carrot.

Fertilizer Level (F)	Carotene content (mg/100 g)		TSS (°brix)	
	L ₁ - Jeevamrut	L ₂ - Amritpani	L ₁ - Jeevamrut	L ₂ - Amritpani
F ₁ - 100 % N	5.16	5.1	14.19	14.1
F ₂ - 75 % N	5.08	5.1	14.08	14.11
F ₃ - 50 % N	5.09	4.77	14.09	13.77

Quality Parameters

Generally, chemical composition of plant reflects on quality of product. Inorganic elements, which are mostly supplied through fertilizer naturally, react with organic elements during metabolism process in the plant body and as a result new organic compound is formed. In the present study carotene, TSS and carbohydrate compounds present in carrot root were quantitatively determined. Its results are described in Table 8. The results indicated that treatments of different land configuration, fertilizer level, liquid formulation and their interaction produced non-significant effect on carbohydrate content in carrot. All the treatments showed almost similar carbohydrate content.

Results regarding carotene and TSS content did not influenced due to land configuration indicating land configuration has no effect on quality parameters of carrot. Numerically higher value of carotene and TSS content of 5.07 mg/100g and 14.11 °brix, respectively was noted in broad bed method (Table 8). The results are in proximity with Desai (1989) who reported non-significant effect of land configurations on quality parameters of groundnut at Rahuri. Balwinder Kumar and Gill (2009) while growing turmeric on Flat and Ridges concluded that planting method did not influence the quality characters of turmeric.

A perusal of the data presented in Table 8 indicated that the different treatment of fertilizer level significantly affected the carotene and TSS content of the carrot. Both the parameters were significantly highest (5.13 mg/100g of carotene and 14.15 °brix of TSS) in the F₁ treatment (100% N through vermicompost) but the values were at par with the treatment F₂ treatment (75% N through vermicompost) and that of lowest in treatment F₃ treatment (50% N through vermicompost). Vermicompost which is produced by earthworms is a rich source of both micro and macro nutrients, vitamins, growth hormones and enzymes increase organic carbon content of soil and improve soil physical properties. This results to better availability and uptake of nitrogen and other nutrients in combination of manures which might have lead to the balance C:N ratio and increased activity of plant metabolism which lead to increase in carotene content and TSS in carrot. Ahmed *et al.* (2014), Chatterjee *et al.* (2014) and Kirad *et al.* (2010) found similar result in carrot. Contrary with the carbohydrate content in this case it was found non significant but with the case of greater yam under organic farming having 100% N application it was found significant (Kaswala *et al.*, 2013).

The different liquid formulations induced significant effect on carotene and TSS content of carrot (Table 8). Application of Jeevamrut showed its superiority and synthesized more carotene (5.11 mg/100 g) and TSS (14.12 °brix) than the application of Amritpani which contained 4.99 mg/100g of carotene and 14.00 °brix of TSS. The organic liquid are eco-friendly organic preparations made from cow products and

contain nutrients, vitamins, amino acids, growth promoting substances like IAA, GA and beneficial microorganisms which can be responsible for quality improvement (Natrajan, 2007 and Sreenivasa *et al.*, 2009).

Among the various interactions, the interaction between fertilizer level and liquid formulation only produced significant effect on carotene and TSS content of carrot (Table 9). The perusal of data indicated that when 100% N was supplied along with Jeevamrut @600 l/ha (F₁L₁) favors to increase the carotene and TSS content. However, for carotene content it was at par with F₂L₁, F₃L₁, F₁L₂ and F₃L₂ and for TSS content F₂L₁, F₃L₁, F₁L₂ and F₂L₂. Sunanda Rani and Malareddy (2007) also reported that application of organic manure in carrot resulted in maximum carotene (4.6 mg kg⁻¹), total sugar and total soluble salts (TSS) when compared to other INM treatments.

REFERENCES

- Ahmad, T., Amjad, M., Iqbal, Q., Nawaz, A. and Iqbal, Z. 2014. Integrated nutrient management practices improve growth and yield of carrot. *Bulgarian J. Agricultural Science*. **20(6)**: 1457-1465.
- Arancon, N. Q., Edwards, C. A., Bierman, P., Metzger, J. D., Lee, S., and Welch, C. 2003. Effects of vermicomposts on growth and marketable fruits of field-grown tomatoes, peppers and strawberries: the 7th international symposium on earthworm ecology. Cardiff-Wales. 2002. *Pedobiologia*. **47(5-6)**: 731-735.
- Bag, P.A., Kaswala, A.R. and Patel, A.I. 2015. Effect of liquid organic manures on yield and soil properties in chickpea. *Trends in Biosciences*. **8(3)**:695-698.
- Balwinder Kumar and Gill, B. S. 2009. Effect of method of planting and harvesting time on growth, yield and quality of turmeric (*Curcuma longa* L.). *J. Spices and Aromatic Crops*. **18(1)**: 22 - 27.
- Chandrakala M. 2008. Effect of FYM and fermented liquid manures on yield and quality of chilli (*Capsium annum* L.). M.Sc. (Agri.) thesis submitted to the University of Agricultural Sciences, Dharwad.
- Chatterjee, Ranjit, Bandyopadhyay, S., and Jana, J. C. 2014. Evaluation of vegetable wastes recycled for vermicomposting and its response on yield and quality of carrot (*Daucus carota* L.). *Indian J. Recycle Organic Waste Agriculture*. **3(2)**: 60-65.
- Desai, M. M. 1989. Effect of planting layout and gypsum levels on the growth, yield and quality of groundnut (*Arachis hypogea* L.) var. Phule Pragati during kharif. M.Sc. (Agri.) thesis to Mahatma Phule Krishi Vishwavidyala, Rahuri (MS), India.
- Devkumar, N., Shubha, S., Gouder, S.B. and Rao, G.G. 2014. Microbial analytical studies of traditional organic preparations beejamrutha and jeevamrutha. Proceedings of the 4th ISOFAR Scientific Conference. 'Building Organic Bridges' at the Organic World Congress held at Istanbul, Turkey during 13 to 15 October, 2014.
- Gadelrab, H.M. and ELAmin, S.M. 2013. *J. Agricultural and Veterinary Science*, Vol. 14, pp. 61-67.
- Gore, N.S. and Shreenivasa, M.N. 2011. Influence of liquid organic manures on growth, nutrient content and yield of tomato (*Lycopersicon esculentum* Mill.) in the sterilized soil. *Karnataka J. Agric. Sci.* **24(2)**:

153-157.

Kaswala, A.R., Kolambe, B.N., Patel, K.G., Patel, V.S. and Patel, S.Y. 2013. Organic production of greater yam: Yield, quality, nutrient uptake and soil fertility. *J. Root Crops.* **39(1)**: 56-61.

Kirad, K. S., Swati, B. and Singh, D. B. 2010. Integrated nutrient management on growth, yield and quality of carrot. *Karnataka J. Agricultural Sciences.* **23(3)**: 542-543.

Lampkin, N. H. 2000. Organic farming. In: S. Padel (ed.). Soil sickness and soil fertility. Cab Publisher, Wallingford, USA.

Mazed, H. K., Pulok, M. A. I., Chowdhury, M. S. N. and Ferdous, J. 2015. Effect of Different Types of Organic Manure and Mulching on the Growth and Yield of Carrot (*Daucus Carota* L.). *International J. Scientific and Research Publications.* **5(2)**: 309-314.

Mengel, K., Kirkby, E. A., Kosegarten, H. and Appel, T. 2001. The soil as a plant nutrient medium. Principles of plant nutrition .pp. 15-110. Springer, Dordrecht.

Natarajan K. 2007. Panchgavya for plant. Proc. Nation. Conf. Glory Gomatha, Dec. 1-3, 2007, S.V. Veterinary Univ., Tirupati, pp. 72-75.

Patel, K.B., Kaswala, A.R., Patel, K.G. and Dubey, P.K. 2016. Influence of paddy straw and organic manures on growth, yield,

quality and economics of onion in organic farming. *Green Farming.* **7(5)**: 1192-1195.

Patil, D.U., Laharia, G.S. and Damre, P.R. 2012. Effect of different organic sources on biological properties of soil, nutrient uptake, quality and yield of soybean. *An Asian J. Soil Science.* **7(2)**: 190-193.

Sankar, V. D., Veragavathatham, M. and Kannan, M. 2009. Organic farming practices in white onion. *J. Eco-friendly Agriculture.* **4(1)**: 17-21.

Sreenivasa M.N., Naik N. and Bhat S.N 2009. Beejamrutha: A source for beneficial bacteria. *Karnataka J. Agric. Sci.* **22(5)**: 1038-1040.

Sunanda Rani, M. and Mallareddy, K. 2007. Effect of different organic manures and inorganic fertilizers on growth, yield and quality of carrot (*Daucus carota* L.). *Karnataka J. Agriculture Science.* **20**: 686-88.

Vinaykumar and Neeraj, 2015. Evaluation of the performance of onion varieties in response to organic cultivation. *International J. Advanced Research.* **3(9)**: 1558- 1562.

Yadav, A. L., Goswami, B., Aggarwal, P. and Arya, M. 2003. Few important soil management technologies for sustainable crop production- a review. *Agriculture Reviews.* **24(3)**: 211-216.



NATIONAL ENVIRONMENTALISTS ASSOCIATION

The National Environmentalists Association is chartered in Ranchi as a nonprofit scientific and educational association of like minded academician, researchers, scientists from all over the nation for the furtherance and diffusion of knowledge of Life Sciences in general and Environmental Science in particular.

The association not only honours its members but also provides FELLOWSHIP to outstanding contributors to the subject and the society.

Contact :

For Editorial Information

Prof. M. P. Sinha
Vice Chancellor
Sido Kanhu Murmu University
Dumka - 814 110
Jharkhand, INDIA

For information regarding Association :

SECRETARY,
National Environmentalists Association,
D-13, Sai Roofs, 1st Floor,
H. H. Colony,
Ranchi - 834002
Jharkhand, India

E-mails : editor.bioscan@gmail.com
dr.mp.sinha@gmail.com
nat.env.assoc@gmail.com

Cell : 94313-60645; 9572649448

Ph. : 0651-2244071

Website : www.thebioscan.com
: www.neaindia.org

NAAS Rating : 5.26

U.S.A. Office

2827 Videre Dr.,
Wilmington,
DE 19808 We, USA

Type setter Bandana Solutions Facility Management LLP
Published by Aditi Publications, Patliputra, Patna

The Bioscan

An International Quarterly Journal of Life Sciences

ISSN : 0973-7049

Volume 15(3) : 2020

Published as an official organ by

NATIONAL ENVIRONMENTALISTS ASSOCIATION

CONTENTS

Page

A. RESEARCH PAPER

1. Comparison and molecular profiling of Begomovirus infecting chilli (*Capsicum annum*) in gangetic alluvial zone of West Bengal
Uday Bikash Oraono, Lourembam Sanajaoba Singh and Jayanta Tarafdar—275 - 280
2. Assessment of probiotic characteristics of *L.plantarum*
Sravani Kandula And Rita Narayanan—281 - 286
3. Effect of rate and frequency of micronutrient on growth attributes and dry matter yield of Banana Cv. grand naine under south Gujarat condition
Narendra Singh, Sonal Tripathi, Patel V. A., Jaimin Naik and Chauhan Aditi—287 - 290
4. Character association and path analysis for seed yield and its components in Grass pea (*Lathyrus sativus* L.)
Gangishetti Ranjithkumar, Sandip Debnath and Duddukur Rajasekhar—291 - 295
5. Determination of physical and biometric properties of onion bulbs in relation to design of digger cum windrower
Shiddanagouda Yadachi and Kiran Nagajjanavar—297 - 301
6. Polygenic variation for morphological and biochemical traits of brinjal genotypes (*Solanum melongena* L.) and its wild relatives
Nisha Sharma, K. D. Bhutia, Rajesh Kumar, Sita Kumari Prasad, Ankita Debnath and Malay Marut Sharma—303 - 309
7. Effect of different sources of horizontal transmission of bacterial flacherie on Et_{50} for symptom expression and mortality of PM X CSR₂
B. L. Kavyashree., R. N. Bhaskar and C. Doreswamy—311 - 314
8. Comparative biology of *Goniozus nephantidis* (Muesbeck) on *Galleria mellonella* L. and *Corcyra cephalonica* (Stainton)
A. V. Desai, M. R. Siddhapara and N. P. Trivedi—315 - 321
9. Efficiency of ovatide on mass seed production of climbing perch (*Anabas testudineus*, Bloch, 1972) in Nalbari district, Assam
Ankur Rajbongshi, A. Ali, M. Chakravarty, M. Deka, H. Mazumdar, Pranab Kr Das and S. Baishya—323 - 326
10. Study of combining ability and gene action for yield and yield component characters in interspecific hybrids of cotton (*Gossypium hirsutum* L. X *Gossypium barbadense* L.)
S. B. Gohil., M. B. Parmar., M. P. Patel and D. A. Patel—327 - 333
11. Per oral inoculation of *Lysinibacillus sphaericus* with pathogenic microbes on rearing and cocoon parameters of silkworm, *Bombyx mori* L.
H. G. Anusha, R. N. Bhaskar and K. V. Anitharani—335 - 338
12. Effect of fungicides, plant extracts and bioagents on spore germination of *Colletotrichum lindemuthianum* causing field bean anthracnose

The Journal is Currently Abstracted / Indexed in

- Paryavarn Abstract, INDIA
- Indian Science Abstract, INDIA
- Cambridge Science Abstract, U.S.A.
- Zoological Record, U.K.
- Directory of Open Access Journal (DOAJ)
- Chemical Abstract, U. S. A.
- Research BIB
- Indian Science
- Journal Seek
- Scientific Indexing Service (SIS)
- Journal is currently rated by
- Index Copernicus
- Universal Impact Factor
- NAAS

DISCLAIMER

The Publisher and Editors cannot be held responsible for errors or any consequences arising from the use of information in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher/ Association and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher / Association and Editors of the products advertised.

- S. Narasimha Rao, S.L. Bhattiprolu, A. Vijaya Gopal and V. Sekhar— 339-344
13. Evaluation in vitro different fungicides for growth of *Rhizoctonia bataticola* A.M. Kadam, S.S. Chavan and A.H. Kendre — 345- 349
 14. Evaluation of gladiolus varieties for flowering and cut flower traits under indo-gangetic plains Girish,P. M., Anjana Sisodia and Anil K. Singh— 351- 355
 15. Effect of land configuration and different organic sources on growth, yield and quality of carrot under organic farming B. Solanki, A. R. Kaswala, P.K. Dubey and A.P. Italiya— 357- 362
 16. Verification and usability analysis of medium range weather forecast for the Kokrajhar district of lower Brahmaputra valley zone of Assam Kuldip Medhi, Kushal Sarmah, Vinod Upadhyay, Sunil Kumar Paul, Athar N. Islam and Bikash J. Gharphalia— 363 - 370
 17. Canonical root analysis and clustering for characterization and evaluation of aromatic rice germplasm based on morphological characters G. Parimala, Ch. Damodhar Raju, L.V. Subba Rao and K. Uma Maheswari— 371 - 374
 18. Genetic divergence analysis of sesame genotypes (*Sesamum indicum* L.) Dasari Rajitha, T. Srikanth, D. Padmaja and T. Kiran Babu—375 - 379
 19. Nutrient uptake and chemical properties of soil after harvest of baby corn (*Zea mays* L.) as influenced by organic manures and fertilizers D.H. Roopashree , S .Kamal Bai . Nagaraju and S.Raghavendra— 381 - 384
 20. Genotypic response on growth and yield in papaya D. K. Varu.,K. D. Patel and Sandip Makhmale— 385 - 389
 21. Studies on frequency distribution of yield and yield related traits in F₂M₂ generation of sesame (*Sesamum indicum* L.) Rajesh Kumar Kar , Tapash Kumar Mishra and Banshidhar Pradhan — 391 - 395
 22. Correlation and path analysis in cowpea (*Vigna unguiculata* (L.) Walp) R.M. Nagalakshmi, R. Usha Kumari and R. Ananda Kumar—397 - 401
 23. A simple and efficient method for DNA extraction from rabi sorghum [*Sorghum bicolor* (L.) Moench]" S. S. Gadakh, G. D. Khalekar.,U. S. Dalvi, A. A.Kale and P.L.Kulwal— 403 - 406
 24. Determination of economic injury level (EIL) of sugarcane plassey borer *Chilo tumidicostalis* hampson (Lepidoptera: pyralidae) R. K. Nath and D. K. Saikia— 407- 409
 25. Performance of different summer mung (*Vigna radiata* L.) varieties sown at different dates under Manipur valley condition Meghna Gogoi, Jamkhogin Lhungdim, Kamal Kant, Urjashi Bhattacharya and Gauri Mohan — 411 - 414



ZONAL CO-ORDINATORS OF THE ASSOCIATION

- Prof. N. Behera
School of Life Science,
Sambalpur University
- Dr. Nirmal Kumar
ISTAR, Vallabh Vidyanagar,
Anand, Gujarat
- Dr. P. N. Sudha
D. K. M. College for Women,
Vellore
- Prof. S. P. S. Dutta
Dept. of Environmental Science,
Jammu University, Jammu
- Dr. V. Salom Gnana Thanga
Dept. of Env. Scs.,
University of Kerala, Kariavattom
Tiruvananthapuram, Kerala

Publications of the Association

The Bioscan

An International Quarterly Journal of Life Sciences

The Ecoscan

An International Quarterly Journal of
Environmental Sciences

**Both the Journals are
online**

**Both the Journals are
available on Google.com**

Websites of the Journals are
www.theecoscan.in
www.thebioscan.in

FEATURES OF ASSOCIATION

- Association is registered under 80G of I.T.
- Prestigious fellowship of the Association (F. N. E. A.) to academicians of the nation.
- Regular annual conference of national and international levels organized by the Association.
- Young Scientist and Senior Scientist award during the conference of the Association.



_____ 062 & 082

_____ 000 & 000

THE BIOSCAN : SUBSCRIPTION RATES				
		India (Rs.)	SAARC Countries	Other Countries
Individuals	One Year	1,000	2,000(I:C)	US \$200
	Life Member*	10,000		
Institutions	One Year	3,000	6,000(I:C)	US \$400
	Life Member*	30,000		

*Life Member will receive the journal for 15 years while other benefits will continue whole life

THE BIOSCAN : MEMBERSHIP FORM

Please enter my subscription for the above journal for the year / life member.

Name:

Address:

E-mail:

Payment Rs. : by DD / MD in favour of
National Environmentalists Association payable at Ranchi, No. Dated
..... is enclosed.

NOTE: FOR MEMBERSHIP THE ABOVE INFORMATION CAN BE SENT ON SEPARATE SHEET