

"CANONICAL ROOT ANALYSIS AND CLUSTERING FOR CHARACTERIZATION AND EVALUATION OF AROMATIC RICE GERMPLASM BASED ON MORPHOLOGICAL CHARACTERS

G. PARIMALA^{1*}, CH. DAMODHAR RAJU², L.V. SUBBA RAO² AND K. UMA MAHESWARI³

¹Department of Genetics and Plant Breeding, College of Agriculture, PJTSAU, ²Department of Genetics and Plant Breeding, Rice Research Centre, Agricultural Research Institute, ⁴Department of Food and Nutrition, Post Graduate & Research centre, PJTSAU, Rajendranagar, Hyderabad-500030, Telangana, INDIA.
e-mail: Parimalagolla1996@gmail.com

KEYWORDS

Canonical root analysis
Genetic diversity
D²
Clusters analysis.

Received on :

20.07.2020

Accepted on :

17.08.2020

*Corresponding author

ABSTRACT

The present study was conducted to estimate the magnitude of genetic diversity among the genotypes and to identify the major morphological traits contributing for the observed variations. A total of 50 aromatic rice genotypes were evaluated in Randomized Block Design with three replications during -2018-2019 kharif season to study genetic diversity using Mahalanobis D² statistics for seed yield and its components. Cluster analysis classified the fifty genotypes into 12 groups. Maximum inter cluster distance was exhibited between clusters VIII and IX (227.03) followed by clusters I and IX (199.86), clusters VII and VIII (190.65), clusters V and IX (187.34) and cluster VII and IX (180.52). The lowest inter cluster divergence was observed between clusters IV and VII (11.01). The intra-cluster distance (D) ranged from 10.23 (cluster-I) to 25.36 (cluster-XII). The attributes, viz., 1000 grain weight (37.63%), Plant height (25.71%) and Seed yield per plant (18.61%) contributed much to the total genetic divergence. PCA revealed three most informative principal components with Eigen values of 6.222, 4.577 and 1.900, respectively, which together accounted 83.323 % of the total variance. The result ensures the existence of high genetic divergence among the studied genotypes and can be used by breeders to develop high yielding rice varieties and have immense applications in rice improvement.

INTRODUCTION

Rice is the staple food across Asia where around half of the world's poorest people live. It is most important human food crop in the world, directly feeding more people than any other crop. Although most rice is consumed in the countries where it is produced, a growing demand in some areas is feeding the international rice trade. Generally, non-aromatic rice varieties are high yielding, demonstrating good agronomic performance, and are highly adapted to the environmental condition and produced in all the rice-growing countries. Conversely, most of the aromatic rice varieties are low yielding, demonstrating inferior agronomic performance, highly prone to the environmental condition and are produced in few countries (Prodhan *et al.*, 2017). Despite inferior performance, the aromatic rice is highly regarded for their excellent aroma and superior grain quality (Wakte *et al.*, 2017). Aromatic rice fetch high price in international trade. So there is a need to develop aromatic rice varieties which are resistant to biotic and abiotic stresses and also high yielding. The development of varieties is a continuous process and the success of the plant breeding programme depends upon the selection of suitable genotypes to be utilized in breeding programme. The effectiveness of selection depends primarily upon the magnitude of genetic diversity in the breeding material at hand. Appropriate and most efficient approach should be used for germplasm evaluation and characterization.

Mahalanobis's D² statistics has been used in several crops for identifying diverse parents for hybridization programmes. It is a powerful tool used to quantify the genetic divergence between the accessions and to relate clustering pattern with the geographical origin. Principal component score strategy has been employed for the identifying the plant characters that categorize the distinctiveness among the promising genotypes. PCA helps to eradicate redundancy in data sets due to regular variation occurring regularly in the crop species (Maji and Saibu, 2012) and (Ramakrishnan *et al.*, 2016). Hence, the importance of PCA is considered and has been successfully used in the germplasm evaluation of crops for understanding the relationship and correlation among the variables studied (Zafar *et al.*, 2008). Therefore this study was organized to estimate the magnitude of genetic distance and to identify the major traits contributing for the observed variations among the studied genotypes.

MATERIALS AND METHODS

Experimental material consists of 50 selected aromatic rice genotypes on the basis of diverse grain morphological characteristics. The experiment was conducted at Rice Research Centre, Agricultural Research Institute, Professor Jayashankar Telangana State Agricultural University, Rajendranagar during Kharif 2018. Randomized Block Design

(RBD), was adopted to conduct the experiment with three replications. Healthy nursery was raised and thirty one days old seedlings were transplanted in well prepared main field. Two rows for each entry with spacing of 20 x 15 cm and each row consists of 25 plants per entry. First weeding was carried out 20 days after transplanting and second weeding 30 days after first weeding. Fertilizer application was carried at recommended dosage. Necessary crop protection measures were taken up based on need of crop. Observations were recorded for quantitative traits on randomly selected five plants for each entry in three replications. The replication wise mean values were used for statistical analysis. Differences among genotypes were tested by Wilk's criterion (Rao, 1952) for the pooled effect of all the 8 characters. The D^2 values for all the possible pairs of varieties were grouped into a number of clusters according to Tocher's method as suggested by Rao (1952). The characters were ranked on the basis of their contribution to D^2 values, in all the combinations. A measure of group distance based on multiple characters was given by Mahalanobis (1936) using D^2 statistic. With the help of this, genetic divergence between genotypes was estimated.

RESULTS AND DISCUSSION

The analysis of variance for individual characters revealed significant differences among genotypes. The value of V-statistic (1860.88) with 343 degrees of freedom showed highly significant differences among the genotypes for aggregate of 8 characters. Thus, one can proceed for further diversity analysis. Grouping of the genotypes was carried-out by following the Tocher's method (Rao, 1952) and were grouped into twelve clusters, with 50 genotypes. The distribution of genotypes into various clusters is shown in (Table 1). The composition of different clusters varied from 1 to 22 genotypes. Cluster II was largest comprising of twenty two genotypes followed by cluster III with eight genotypes, cluster I with six genotypes, cluster IX with 4 genotypes and cluster XII with 3 genotypes. The clusters IV, V, VI, VII, VIII, X and XI were represented by single genotype exhibiting high degree of heterogeneity among the genotypes. The discrimination of lines into so many discrete clusters suggested presence of high degree of genetic diversity in the

material evaluated. Similarly 75 diverse genotypes of soft rice were grouped into seven clusters by pragnya *et al.* (2018). 216 genotypes were grouped into 15 clusters by Praveen *et al.* (2019). Twenty six cultivars of rice were grouped into six clusters by Shivani *et al.* (2018). The clustering pattern of genotypes from different eco-geographical regions into these clusters was apparently random genotypes of different origins were clubbed in one cluster, whereas the genotypes belonging to same state or origin were grouped in different clusters thereby indicating non-relationship between geographical and genetic diversity. Singh *et al.* (2007) also reported that there was no parallelism between geographic distribution and genetic diversity. Pattern of distribution of genotypes among various clusters reflected the considerable genetic diversity present in the genotypes under study. The choice of suitable diverse parents based on genetic divergence analysis would be more fruitful than the choice made on the basis of geographic distances in accordance with Raut *et al.* (2009).

Number of times that each of the yield and its component characters appeared in first rank and its respective percent contribution towards genetic divergence is presented in (Table 4). The results recorded that the contribution of 1000-grain weight towards genetic divergence was highest (37.63 %) by taking 461 times ranking first similar with the results of Reddy *et al.* (2013), followed by plant height (25.71) by taking 315 times first, seed yield per plant (18.61) by taking 228 times first, number of grains per panicle (11.75) by first ranking 144 times, days to fifty percent flowering (5.87) ranking 72 times first, no. of effective tillers per plant (0.4%) by ranking 5 times first. Three characters out of the seven characters studied, namely 1000-grain weight, plant height, seed yield per plant contributed (81.95 %) to the total genetic divergence. The results obtained in the present study are in accordance to the findings of Tripathi *et al.* (2017), Ahamed *et al.* (2018).

Cluster means of yield and its component traits were presented in (Table 3). Cluster means data shows that days to 50% flowering was highest in cluster X (117.67) and lowest in cluster VI (100.33), plant height recorded highest in cluster IX (174.11) and lowest in cluster VII (101.67), no. of effective tillers per plant was observed highest in cluster VII (19.53) and lowest in

Table 1: Clustering pattern among 50 aromatic rice genotypes for yield and its component traits by Tocher method

Cluster number	No, of Genotypes	Genotypes
I	6	RNR 15462-2, RNR 15459-1, RNR 15460-1, RNR-15462-4, RNR-15453-2, Sugandh Samba
II	22	RNR-29420, RNR-29421, RNR-29422, RNR-29423, NVSR 406, RNR-28408, RNR-28405, RNR-26058, RNR-15488-2, RNR-28404, RNR-2600, RNR-28403, RNR-29301, NVSR 407, RNR-26009, RNR-26010, RNR-29296, Vasumathi, RNR-29296, RNR-26020, RNR-28402, RNR 17500.
III	8	RNR 28410, Ketekijoha, CR 3663-261-8-4, RNR17501, RNR 28410, CR 3715-119-18-9-2, CR 2981-16-2-6, CR 2982-14-6-3
IV	1	RNR-29305
V	1	PUSA-1121
VI	1	Sumathi
VII	1	RNR-29306
VIII	1	RNR-15459-6
IX	4	OR (CZ)-70, BM 4, Dubraj, JDP-K-37
X	1	CR 3715-119-18-9-2
XI	1	Chittimutyalu
XII	3	NWGR 9081, Pusa Basmathi, Shobini

Table 2: Intra (diagonal) and inter cluster distances of D² values of 50 genotypes yield and its component characters

Clusters	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5	Cluster 6	Cluster 7	Cluster 8	Cluster 9	Cluster 10	Cluster 11	Cluster 12
Cluster 1	10.23	57.54	55.61	122.42	108.95	43.38	123.45	21.28	199.86	117.28	27.21	120.72
Cluster 2		18.67	41.78	29.45	40.28	29.39	43.49	81.17	156.08	34.47	56.42	53.08
Cluster 3			22.16	66.09	68.44	42.27	75.94	73.64	115.84	43.59	68.6	54.14
Cluster 4				0	50.52	70.58	11.01	166.92	174.4	16.12	113.13	41.36
Cluster 5					0	22.52	98.39	95.84	187.34	32.62	120.33	78.42
Cluster 6						0	105.64	34.09	156.54	58.95	57.56	74.01
Cluster 7							0	190.65	180.52	36.44	105.22	58.77
Cluster 8								0	227.03	140.54	53.72	154.81
Cluster 9									22.2	152.24	138.44	161.87
Cluster 10										0	121.86	42.26
Cluster 11											0	136.38
Cluster 12												25.36

Table 3: Cluster means of 50 rice genotypes for yield and its component characters by Tocher method

Characters	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII
Days to 50% flowering	107.67	107.15	114.67	109.33	104.67	100.33	114.33	105.67	112.67	117.67	108.67	105.11
Effective tillers per plant	19.09	17.71	18.59	18.67	18.83	19.2	19.53	17.78	18.9	17.47	17.87	12.79
Plant height	102.57	105.96	112.6	102.27	106.73	111.43	101.67	103.78	174.11	103.8	125.09	103.78
Panicle length	21.91	24.82	24.23	24.33	25.83	26.63	22.53	22.31	27.8	22.43	25.25	26.3
Panicle weight	3.26	7.56	6.52	5.52	4.31	3.56	5.26	7.14	6.54	5.54	5.23	5.63
Grains per panicle	168.22	155.67	173.46	169.67	89.33	119.33	204.67	122	190.83	147.67	173.33	196.56
1000 grain weight	8.44	17.77	15.3	23.94	21.06	14.94	22.56	7.13	15.63	22.99	8.81	21.42
Seed yield per plant	23.82	27.38	16.11	29.13	23.74	22.4	31.3	20.2	18.21	21.55	32.07	17.59

Table 4: Relative contribution of yield and yield contributing characters to genetic diversity

S. No	Characters	Times ranked first	contribution (%)
1	Das to 50% flowering	72	5.87%
2	Plant height (cm)	315	25.71%
3	No. of effective tillers per plant	5	0.40%
4	Panicle length (cm)	0	0.02%
5	Panicle weight (g)	0	0.01%
6	1000 grain weight (g)	461	37.63%
7	No. of grains per panicle	144	11.75%
8	Seed yield per plant (g)	228	18.61%

Table 5: Eigen values, contribution of variability and factor loading for the principal component axis .

	PC 1	PC 2	PC 3
Eigenvalue	6.222	4.577	1.9
Variability (%)	40.823	30.032	12.467
Cumulative %	40.823	70.856	83.323
Days to 50% flowering	0.114	0.092	0.144
Productive tillers	0.008	0.113	-0.028
Plant height (cm)	0.814	0.444	-0.326
Panicle length (cm)	-0.001	-0.083	-0.065
Panicle weight (gm)	0.042	-0.023	0.091
No of filled grains per panicle	0.178	0.117	0.209
1000 Seed weight	0.494	-0.86	0.025
Yield per plant	0.219	-0.139	0.907

cluster XII (12.79), panicle length was highest in cluster IX (27.80) and lowest in cluster I (21.91), 1000 grain weight recorded highest in cluster IV (23.94) and lowest in cluster VIII (7.13), no. of grains per panicle was found to be highest in cluster VII (204.67) and lowest in cluster V (89.33), seed yield per plant was recorded was highest in cluster XI (32.07) and lowest in cluster III (16.11).

The average intra and inter cluster D² values are presented in (Table 2). Intra cluster D² values are minimum (0.00) in clusters , IV and V, VI, VII, VIII, X, XI as these were monogenotypic clusters and genotypes in these clusters were more divergent

and they could be utilized as parents for hybridization. Intra cluster distance observed in cluster I is (10.23), cluster II (18.67), cluster III (22.16), cluster IX (22.20), cluster XII (25.36) revealing that some genetic divergence still existed among the genotypes of the cluster. Similar results were seen in findings of Tripathi *et al.* (2017). Selection in these clusters might be executed based on maximum mean value for the desirable characters. The intra cluster distance was lower than inter cluster distance (Table 2), indicating the existence of genetic diversity among the genotypes under study.

Maximum inter cluster distance was exhibited between clusters VIII and IX (227.03) followed by clusters I and IX (199.86), clusters VII and VIII (190.65), clusters V and IX (187.34) and cluster VII and IX (180.52). The lowest inter cluster divergence was observed between clusters IV and VII (11.01) followed by clusters IV and X (16.12), clusters I and VIII (21.28) describing that the genotypes included in these clusters were closely related. The genotypes grouped into same cluster displayed the lowest degree of divergence as over genotypes from different clusters ,which indicates transgressive segregation is possible in the later case. Therefore, hybridization programmes should always be formulated in such a way that the parents belonging to different clusters with maximum divergence could be utilized. Ahamed *et al.* (2018) studied genetic divergence in 50 rice germplasm lines and found that the inter cluster distance in most cases was larger than intra cluster distance suggesting wider diversity among the germplasm of different groups.

The purpose of the PCA is to obtain a small number of factors which account for maximum variability out of the total variability. Eigen vector values, percentage of variance and the cumulative percentage are presented in (Table 5). In this case, PCA revealed three most informative principal components with Eigen values of 6.222, 4.577 and 1.900, respectively, which together accounted 83.323 % of the total variance for all the characters 3 components had Eigen values greater than 1.0. Percentage of variance for the three factors

was 40.823, 30.032, and 12.467 per cent. Together they accounted for 83.32 % of the variability of the genotypes used for the diversity analysis. According to principal component 1, characters such as plant height (0.814), 1000 seed weight (0.494) and seed yield per plant (0.219) had relatively higher contributions (40.823%) to the total morphological variability, while the second principal component accounted for 30.032% of the total variation with plant height (0.444) giving the highest contribution. Maji, A.T and Shaibu AA. (2012) carried out P.C.A with one hundred and twenty three rice germplasm in which first two components accounting for 78% of the total variation. The first three principal components contributed 74.20% of the total variation when 52 genotypes were analyzed with P.C.A by Enyew *et al.* (2019).

REFERENCES

- Ahamed, A., Singh, V.K., Sharma, V and Paswan, S.K. 2014. Genetic Divergence Analysis in Rice (*Oryza Sativa* L.) germplasm in sodic soil condition. *The Ecoscan*. **(6)**:51-54.
- Enyew, M., Dejene, T., Lakew, B and Worede, F. 2019. Clustering and principal component analysis of Barley (*Hordeum vulgare* L.) Landraces for major morphological traits from North Western Ethiopia. *International J. Agriculture Science and Food Technology*. **5(1)**:058-063.
- Mahalanobis, P.C. 1936. On the generalized distance in statistics. *Proceedings of National Institute of Sciences, India*. **12**: 49-55.
- Maji, A.T and Shaibu AA. 2012. Application of principal component analysis for rice germplasm characterization and evaluation. *J. Plant Breeding and Crop Science*. **4(6)**:87-93.
- Nuruzzaman, M., Rana, M.S., Ferdausi, A., Huda, M.M., Hassan, L and Begum, S.N. 2019. Clustering and Principal Component Analysis of Nerica Mutant Rice Lines Growing Under Rainfed Condition. *International J. Applied Sciences and Biotechnology*. **7(3)**: pp.327-334.
- Pachauri, A.K., Sarawgi, A.K., Bhandarkar, S. and Ojha, G.C. 2017. Agro-morphological characterization and morphological based genetic diversity analysis of Rice (*Oryza sativa* L.) germplasm. *J. Pharmacognosy and Phytochemistry*. **6(6)**: pp.75-80.
- Parhe, S.D., Harer, P.N. and Nagawade, D.R. 2014. Investigation of genetic divergence in chickpea (*Cicer arietinum* L.) genotypes. *The Bioscan*. **9(2)**: 879-882 9(2), pp.879-882.
- Parveen, S., Singh, L and Kanawjia, A. 2019. Genetic divergence in indigenous and exotic genotypes of rice (*Oryza sativa* L.). *International J. Current Microbiology and Applied Sciences*. **8**: 122-125.
- Pragnya, K., Radha Krishna, K.V., Subba Rao, L.V and Suneetha, K. 2018. Studies on genetic divergence analysis in soft rice (*Oryza sativa* L.) genotypes. *International J. Pure Applied Bioscience*. **6(2)**: 9689-75.
- Prodhan, Z.H., Faruq, G., Rashid, K.A. and Taha, R.M., 2017. Effects of temperature on volatile profile and aroma quality in rice. *Int J Agric Biol*. **19(5)**: 1065-1072.
- Ramakrishnan, P., Babu, C., Iyanar, K and N. Manivannan. 2014. Principal component analysis for evaluation of Guinea grass (*Panicum maximum* Jacq.) germplasm accessions. *The Bioscan*. **11(4)**: 2859-2862.
- Rao, C. R. 1952. *Advanced statistical method in biometri research.*, John Wiley and Sons, Inc. New York. **15(10)**: 130-134.
- Raut, K. R., Harer, P. N. and Yadav, P. S. 2009. Genetic divergence in rice (*Oryza sativa* L.). *J. Maharashtra Agril. Univ*. **34(2)**: 172-174.
- Reddy, G.E., Suresh, B.G., Sravan, T and Reddy, P.A. 2013. Interrelationship and cause-effect analysis of rice genotypes in North East plain zone. *The Bioscan*. **8(4)**: 1141-1144.
- Sheela, K.R.V., Robin, S. and Manonmani, S., 2020. Principal component analysis for grain quality characters in rice germplasm. *Electronic J. Plant Breeding*. **11(1)**:127-131.
- Shekhawat, N., Jadeja, G.C., Singh, j. and Ramesh. 2014. Genetic Diversity Analysis In Relation To Seed Yield And Its Component Traits In Indian Mustard (*Brassica Juncea* L. Czern & Coss). *The Bioscan*. **9(2)**: 713-717.
- Shivani, D.K., Husain, D.R., Gyanendra, K and Khan, N.A. 2018. Genetic Divergence for yield and other quantitative traits in rice (*Oryza Sativa* L.). *International J. Current Microbiology and Applied Sciences*. **7(1)**: 1201- 1207.
- Singh, V., Bhanjan, R. and Kumar, K. 2007. Genetic divergence in Indian mustard [*Brassica juncea*. (L.) Czern and Coss]. *Prog. Agric*. **7(1/2)**: 105-109.
- Tripathi, A., Kumar, S., Singh, M.K., Kumar, A and Karnwal, M.K. 2017. Phenotypic assessment of rice (*Oryza sativa* L.) genotypes for genetic variability and varietal diversity under direct seeded condition. *J. Applied and Natural Science*. **9(1)**: 6 - 9.
- Wakte, K., Zanan, R., Hinge, V., Khandagale, K., Nadaf, A and Henry, R. 2017. Thirty three years of 2 acetyl 1 pyrroline, a principal basmati aroma compound in scented rice (*Oryza sativa* L.): a status review. *J. the Science of Food and Agriculture*. **97(2)**: 384-395.
- Zafar, I., Arshad, M., Ashraf, M., Mahmood, T and Abdul, W. 2008. Evaluation of soybean (*Glycine max* (L.) Merrill germplasm for some important morphological traits using multivariate analysis. *Pakistan J. Botany*. **40**: 2323–2328.



**NATIONAL ENVIRONMENTALISTS
ASSOCIATION**

The National Environmentalists Association is chartered in Ranchi as a nonprofit scientific and educational association of like minded academicians, researchers, scientists from all over the nation for the furtherance and diffusion of knowledge of Life Sciences in general and Environmental Science in particular.

The association not only honours its members but also provides FELLOWSHIP to outstanding contributors to the subject and the society.

Contact :

For Editorial Information

Prof. M. P. Sinha
Vice Chancellor
Sido Kanhu Murmu University
Dumka - 814 110
Jharkhand, INDIA

For information regarding Association :

SECRETARY,
National Environmentalists Association,
D-13, Sai Roofs, 1st Floor,
H. H. Colony,
Ranchi - 834002
Jharkhand, India

E-mails : editor.bioscan@gmail.com
dr.mp.sinha@gmail.com
nat.env.assoc@gmail.com

Cell : 94313-60645; 9572649448

Ph. : 0651-2244071

Website : www.thebioscan.com
: www.neaindia.org

NAAS Rating : 5.26

U.S.A. Office

2827 Videre Dr.,
Wilmington,
DE 19808 We, USA

Type setter Bandana Solutions Facility Management LLP
Published by Aditi Publications, Patliputra, Patna

The Bioscan

An International Quarterly Journal of Life Sciences

ISSN : 0973-7049

Volume 15(3) : 2020

Published as an official organ by

NATIONAL ENVIRONMENTALISTS ASSOCIATION

CONTENTS

Page

A. RESEARCH PAPER

1. Comparison and molecular profiling of Begomovirus infecting chilli (*Capsicum annum*) in gangetic alluvial zone of West Bengal
Uday Bikash Oraono, Lourembam Sanajaoba Singh and Jayanta Tarafdar—275 - 280
2. Assessment of probiotic characteristics of *L.plantarum*
Sravani Kandula And Rita Narayanan—281 - 286
3. Effect of rate and frequency of micronutrient on growth attributes and dry matter yield of Banana Cv. grand naine under south Gujarat condition
Narendra Singh, Sonal Tripathi, Patel V. A., Jaimin Naik and Chauhan Aditi—287 - 290
4. Character association and path analysis for seed yield and its components in Grass pea (*Lathyrus sativus* L.)
Gangishetti Ranjithkumar, Sandip Debnath and Duddukur Rajasekhar—291 - 295
5. Determination of physical and biometric properties of onion bulbs in relation to design of digger cum windrower
Shiddanagouda Yadachi and Kiran Nagajjanavar—297 - 301
6. Polygenic variation for morphological and biochemical traits of brinjal genotypes (*Solanum melongena* L.) and its wild relatives
Nisha Sharma, K. D. Bhutia, Rajesh Kumar, Sita Kumari Prasad, Ankita Debnath and Malay Marut Sharma—303 - 309
7. Effect of different sources of horizontal transmission of bacterial flacherie on Et₅₀ for symptom expression and mortality of PM X CSR₂
B. L. Kavyashree, R. N. Bhaskar and C. Doreswamy—311 - 314
8. Comparative biology of *Goniozus nephantidis* (Muesbeck) on *Galleria mellonella* L. and *Corcyra cephalonica* (Stainton)
A. V. Desai, M. R. Siddhapara and N. P. Trivedi—315 - 321
9. Efficiency of ovatide on mass seed production of climbing perch (*Anabas testudineus*, Bloch, 1972) in Nalbari district, Assam
Ankur Rajbongshi, A. Ali, M. Chakravarty, M. Deka, H. Mazumdar, Pranab Kr Das and S. Baishya—323 - 326
10. Study of combining ability and gene action for yield and yield component characters in interspecific hybrids of cotton (*Gossypium hirsutum* L. X *Gossypium barbadense* L.)
S. B. Gohil, M. B. Parmar, M. P. Patel and D. A. Patel—327 - 333
11. Per oral inoculation of *Lysinibacillus sphaericus* with pathogenic microbes on rearing and cocoon parameters of silkworm, *Bombyx mori* L.
H. G. Anusha, R. N. Bhaskar and K. V. Anitharani—335 - 338
12. Effect of fungicides, plant extracts and bioagents on spore germination of *Colletotrichum lindemuthianum* causing field bean anthracnose

The Journal is Currently Abstracted / Indexed in

- Paryavarn Abstract, INDIA
- Indian Science Abstract, INDIA
- Cambridge Science Abstract, U.S.A.
- Zoological Record, U.K.
- Directory of Open Access Journal (DOAJ)
- Chemical Abstract, U. S. A.
- Research BIB
- Indian Science
- Journal Seek
- Scientific Indexing Service (SIS)

Journal is currently rated by

Index Copernicus

Universal Impact Factor

NAAS

DISCLAIMER

The Publisher and Editors cannot be held responsible for errors or any consequences arising from the use of information in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher/ Association and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher / Association and Editors of the products advertised.

- S. Narasimha Rao, S.L. Bhattiprolu, A. Vijaya Gopal and V. Sekhar— 339-344
13. Evaluation in vitro different fungicides for growth of *Rhizoctonia bataticola* A.M. Kadam, S.S. Chavan and A.H. Kendre — 345- 349
14. Evaluation of gladiolus varieties for flowering and cut flower traits under indo-gangetic plains Girish,P. M., Anjana Sisodia and Anil K. Singh— 351- 355
15. Effect of land configuration and different organic sources on growth, yield and quality of carrot under organic farming B. Solanki, A. R. Kaswala, P.K. Dubey and A.P. Italiya— 357- 362
16. Verification and usability analysis of medium range weather forecast for the Kokrajhar district of lower Brahmaputra valley zone of Assam Kuldip Medhi, Kushal Sarmah, Vinod Upadhyay, Sunil Kumar Paul, Athar N. Islam and Bikash J. Gharphalia— 363 - 370
17. Canonical root analysis and clustering for characterization and evaluation of aromatic rice germplasm based on morphological characters G. Parimala, Ch. Damodhar Raju, L.V. Subba Rao and K. Uma Maheswari— 371 - 374
18. Genetic divergence analysis of sesame genotypes (*Sesamum indicum* L.) Dasari Rajitha, T. Srikanth, D. Padmaja and T. Kiran Babu—375 - 379
19. Nutrient uptake and chemical properties of soil after harvest of baby corn (*Zea mays* L.) as influenced by organic manures and fertilizers D.H. Roopashree , S .Kamal Bai . Nagaraju and S.Raghavendra— 381 - 384
20. Genotypic response on growth and yield in papaya D. K. Varu.,K. D. Patel and Sandip Makhmale— 385 - 389
21. Studies on frequency distribution of yield and yield related traits in F₂M₂ generation of sesame (*Sesamum indicum* L.) Rajesh Kumar Kar , Tapash Kumar Mishra and Banshidhar Pradhan — 391 - 395
22. Correlation and path analysis in cowpea (*Vigna unguiculata* (L.) Walp) R.M. Nagalakshmi, R. Usha Kumari and R. Ananda Kumar—397 - 401
23. A simple and efficient method for DNA extraction from rabi sorghum [*Sorghum bicolor* (L.) Moench]" S. S. Gadakh, G. D. Khalekar.,U. S. Dalvi, A. A.Kale and P.L.Kulwal— 403 - 406
24. Determination of economic injury level (EIL) of sugarcane plassey borer *Chilo tumidicostalis* hampson (Lepidoptera: pyralidae) R. K. Nath and D. K. Saikia— 407- 409
25. Performance of different summer mung (*Vigna radiata* L.) varieties sown at different dates under Manipur valley condition Meghna Gogoi, Jamkhogin Lhungdim, Kamal Kant, Urjashi Bhattacharya and Gauri Mohan — 411 - 414



ZONAL CO-ORDINATORS OF THE ASSOCIATION

- Prof. N. Behera
School of Life Science,
Sambalpur University
- Dr. Nirmal Kumar
ISTAR, Vallabh Vidyanagar,
Anand, Gujarat
- Dr. P. N. Sudha
D. K. M. College for Women,
Vellore
- Prof. S. P. S. Dutta
Dept. of Environmental Science,
Jammu University, Jammu
- Dr. V. Salom Gnana Thanga
Dept. of Env. Scs.,
University of Kerala, Kariavattom
Tiruvananthapuram, Kerala

Publications of the Association

The Bioscan

An International Quarterly Journal of Life Sciences

The Ecoscan

An International Quarterly Journal of
Environmental Sciences

**Both the Journals are
online**

**Both the Journals are
available on Google.com**

Websites of the Journals are
www.theecoscan.in
www.thebioscan.in

FEATURES OF ASSOCIATION

- Association is registered under 80G of I.T.
- Prestigious fellowship of the Association (F. N. E. A.) to academicians of the nation.
- Regular annual conference of national and international levels organized by the Association.
- Young Scientist and Senior Scientist award during the conference of the Association.



_____ 062 & 082

_____ 000 & 000

THE BIOSCAN : SUBSCRIPTION RATES				
		India (Rs.)	SAARC Countries	Other Countries
Individuals	One Year	1,000	2,000(I:C)	US \$200
	Life Member*	10,000		
Institutions	One Year	3,000	6,000(I:C)	US \$400
	Life Member*	30,000		

*Life Member will receive the journal for 15 years while other benefits will continue whole life

THE BIOSCAN : MEMBERSHIP FORM

Please enter my subscription for the above journal for the year / life member.

Name:

Address:

E-mail:

Payment Rs. : by DD / MD in favour of
National Environmentalists Association payable at Ranchi, No. Dated
..... is enclosed.

NOTE: FOR MEMBERSHIP THE ABOVE INFORMATION CAN BE SENT ON SEPARATE SHEET