

NUTRIENT UPTAKE AND CHEMICAL PROPERTIES OF SOIL AFTER HARVEST OF BABY CORN (*Zea mays* L.) AS INFLUENCED BY ORGANIC MANURES AND FERTILIZERS

D.H. ROOPASHREE *, S.KAMAL BAI . NAGARAJU AND S. RAGHAVENDRA

Department of Agronomy, University of Agricultural Sciences, Bangalore, Karnataka, INDIA

e-mail: roopa229@rediffmail.com

KEYWORDS

Baby corn
Nutrients uptake
Organic manures
Yield

Received on :

17.05.2020

Accepted on :

11.08.2020

*Corresponding author

ABSTRACT

An experiment was conducted during Kharif season, to study the effect of combination of different source and dosage of organic manures with fertilizers on nutrients uptake, available nutrients and chemical properties of soil after harvest of baby corn and its effect on yield of baby corn. Results revealed that, application of recommended dose of fertilizer along with FYM (150:75:40 kg N: P₂O₅:K₂O ha⁻¹) + 10 t FYM recorded significantly higher uptake of nitrogen, phosphorus and potassium (204.24, 35.23 and 213.6 kg/ha, respectively) than applying only recommended dose of fertilizer. In same way, application of 100% N through different sources of organic manures, application of FYM + balance P and K through fertilizer significantly recorded next higher available nitrogen, phosphorus and potassium (233.5, 33.0 and 226.8 kg/ha, respectively) compared to other combinations of organic manure treatments. The same trend was noted with respect to yield. Highest baby corn yield (17.67q ha⁻¹) and green fodder yield (36.53 t ha⁻¹) was recorded in treatment where application of organic manure through FYM in combination of recommended dose of fertilizer (150:75:40 kg N: P₂O₅:K₂O ha⁻¹) + 10 t FYM was applied. Organic carbon content of soil was improved where organic manures along with fertilizers were applied.

INTRODUCTION

Maize, of all the cereal grains is the most highly valued for its multifarious uses, being utilized as human food, animal feed and raw materials in industry. Maize is the third most important cereal crop next to rice and wheat and has the highest production potential among the cereals. For diversification and value addition, as well as growth of food processing industries, an interesting recent development is growing maize for vegetable purpose, which is known as 'baby corn'. It is so called because young, fresh and fingerlike green ears are harvested when the silk length is of 2-3 cm but prior to fertilization (Pandey *et al.*, 2000). It is used as vegetable after dehusking and desilking. Its delicate texture, sweet flavour and crisp nature contribute to its increasing popularity making it an indispensable ingredient in many multi-cuisine dishes of present day.

Baby corn is a vegetable crop that can potentially improve the economic status of farmers (Das *et al.*, 2008). In addition to its sweet, succulent, and delicious taste, baby corn's nutritional value is comparable to other vegetables such as cauliflower, cabbage and tomato. When baby corn is harvested good quality green fodder is also obtained. The succulent green fodder of high quality adds enormously to the total returns to the farmers, resulting in higher profit per unit area per unit time compared to grain maize. Although, baby corn has been developed as an export vegetable that can generate foreign exchange, it is serving the local people with nutritious vegetable. Despite its great nutritional importance and economic security to the farmers only limited scientific research

has been reported on baby corn, resulting in insufficient knowledge and lack of standard technologies that hamper the popularization of baby corn production (Muthukumar *et al.*, 2007; Thavaprakash and Velayudham., 2008). Cultivation practices have not been standardized. With this background, to standardize the agro techniques not only for its potential yield but also for its quality babies and fodder this study was taken at farmers field.

MATERIALS AND METHODS

An experiment was conducted during Kharif 2008 and 2009, at farmer's field of Shettahalli Village, Mandya District, Karnataka, India situated in 12018' to 13004' N latitude and 76079' to 77020'E longitude with an altitude of 695 meters above the MSL located in the Agro

Climatic Zone - 6 (Southern Dry Zone) of Karnataka. The soil of experimental site was sandy clay loam in texture. The soil pH was 6.8 with an electrical conductivity of 0.14 dSm⁻¹. The soil was low in available nitrogen (210.8 kg ha⁻¹), medium in available phosphorus (24.6 kg ha⁻¹) and potassium (168.0 kg ha⁻¹). The organic carbon content was low (0.46 %). Nutrients composition of organic manures used in the experiment were N: 0.56 %, P:0.21 % and K:0.45 % (Farm yard manure), N: 1.80 %, P:1.10 % and K:1.58 % (vermicompost) and N: 3.21 %, P:2.16 % and K:1.72 % (poultry manure). The experiment was laid out in a randomized block design with eight treatments, viz. T₁: Recommended dose of fertilizer (RDF) (150:75:40 kg N:P₂O₅:K₂O ha⁻¹), T₂: 50 % N through FYM + balance N, P and K through fertilizer, T₃: 50 % N through

vermicompost + balance N, P and K through fertilizer, T₄: 50 % N through poultry manure + balance N, P and K through fertilizer, T₅: 100% N through FYM + balance P and K through fertilizer, T₆: 100% N through vermicompost + balance P and K through fertilizer, T₇: 100% N through poultry manure + balance P and K through fertilizer, T₈: RDF (150:75:40 kg N: P₂O₅: K₂O ha⁻¹) + 10 t FYM ha⁻¹ and replicated thrice. The nitrogen contents in organic manures viz., FYM, vermicompost and poultry manure were determined and the amount of these materials required for substituting recommended dose of nitrogen as per the treatment was calculated and applied 15 days prior to sowing and were thoroughly mixed in the soil. Fertilizer N was applied as per the treatments in two equal splits viz., first half at the time of sowing as basal dose and remaining half as top dressed at 20 days after emergence. Entire quantity of phosphorus and potassium were applied as basal dose. Baby corn seeds were sown with recommended spacing (45 x 20 cm) with a variety PAC 792. Five plants were randomly selected in each plot of each replication and were tagged for the purpose of recording observations. Tassels were removed as and when they emerged. This was done to avoid pollination. If the silk gets pollinated, the kernels would start developing within hours and the cob would become hard and unfit for consumption. Harvesting was done at three days after silk emergence. Similarly, baby corn and fodder yield from each net plot in each replication was harvested, weighed and recorded as baby corn and fodder yield per net plot. Further, this net plot yields were converted to yields per hectare. The analysis and interpretation of the data was done using the Fisher's method of analysis and variance techniques as given by Panse and Sukatme (1967). The level of significance used in 'F' and 't' test was P=0.05 probability level and wherever 'F' test was found significant, the 't' test was performed to estimate critical differences among various treatments.

RESULTS AND DISCUSSION

Effect of organic manures and fertilizers on nutrient uptake by baby corn

The pooled data of two years (Table 1) on uptake of nitrogen, phosphorus and revealed that, application of recommended dose of fertilizer (150:75:40 kg NPK ha⁻¹) + FYM (10 t ha⁻¹) recorded significantly maximum uptake of nitrogen, phosphorus and potassium (204.24, 35.23 and 213.6 kg/ha,

respectively) compared to other treatments. However, 50 per cent N through organic manure and balance NPK through fertilizer recorded highest nitrogen, phosphorus and potassium uptake as compared to 100 per cent N through organic manure and balance P and K through fertilizer. Total uptake of nutrients showed a pronounced positive effect with the application of combination of organic manures and fertilizers. Combined application of organic manures and fertilizers ensure the release of readily available nutrients in adequate quantity to promote early growth as compared to sole organic manure treatments. This might be due to result of steady and continuous availability of instant N in the rhizosphere. The lowest uptake of nitrogen, phosphorus and potassium (118.47, 17.09 and 140.15 kg/ha, respectively) recorded in the treatment with the application of 100 per cent N through FYM and balance P and K through fertilizer compared to other treatments. This is attributed to lesser availability of instant nutrients in rhizosphere, due to relatively slow mineralization of organic sources. This corroborates the findings of Arun Kumar *et al.* (2009).

Effect of organic manures and fertilizers on available nutrients in soil

The pooled data of two years (Table 2) revealed that, post harvest soil fertility status has been significantly improved with the application of 100 per cent N through organic manures. The highest available nitrogen was obtained with 100 per cent N through FYM and balance P and K through fertilizer (T₅) (233.55 kg ha⁻¹) and was on par with 100 per cent N through poultry manure and balance P and K through fertilizer (T₇) (229.07 kg ha⁻¹) and 100 per cent N through vermicompost and balance P and K through fertilizer (T₆) (225.70 kg ha⁻¹). The highest available P₂O₅ (38.02 kg ha⁻¹) was recorded in 100 per cent N through poultry manure and balance P and K through fertilizer (T₇) and was on par with 100 per cent N through vermicompost and balance P and K through fertilizer (T₆) (35.92 kg ha⁻¹). The highest available K₂O content (226.80 kg ha⁻¹) was noticed in 100 per cent N through FYM and balance P and K through fertilizer (T₅). This might be owed to slow mineralization of organic fraction coupled with under utilization of applied nutrients by baby corn. The lowest post harvest soil fertility status was found with supply of only recommended dose of fertilizer. These results are in conformity with the findings of Karki *et al.* (2005).

Table 1: Nutrients uptake by baby corn as influenced by organic manures and fertilizers

| Treatments | Nitrogen (kg ha ⁻¹) | | | Phosphorus (kg ha ⁻¹) | | | Potassium (kg ha ⁻¹) | | |
|----------------|---------------------------------|--------|--------|-----------------------------------|-------|--------|----------------------------------|--------|--------|
| | 2008 | 2009 | Pooled | 2008 | 2009 | Pooled | 2008 | 2009 | Pooled |
| T ₁ | 176.79 | 190.13 | 179.72 | 30 | 32.67 | 31.33 | 198.16 | 201.65 | 199.9 |
| T ₂ | 142.16 | 151.83 | 143.32 | 20.71 | 21.81 | 21.26 | 161.01 | 162.12 | 161.57 |
| T ₃ | 173.34 | 184.53 | 176.13 | 25.76 | 27.17 | 26.47 | 182.95 | 185.08 | 184.01 |
| T ₄ | 160.53 | 171.87 | 162.88 | 24.72 | 26.14 | 25.43 | 166.91 | 167.57 | 167.24 |
| T ₅ | 117.2 | 126.62 | 118.47 | 17.29 | 16.88 | 17.09 | 139.32 | 140.99 | 140.15 |
| T ₆ | 131.28 | 138.79 | 131.95 | 19.07 | 19.69 | 19.38 | 150.57 | 151.94 | 151.26 |
| T ₇ | 124.14 | 131.44 | 125.33 | 17.97 | 19.21 | 18.59 | 143.89 | 146.16 | 145.03 |
| T ₈ | 203.11 | 215.41 | 204.24 | 33.94 | 36.51 | 35.23 | 212.59 | 214.63 | 213.61 |
| S.Em + | 7.23 | 9.47 | 6.85 | 1.77 | 1.98 | 1.53 | 6.27 | 6.92 | 6.5 |
| C.D at 5 % | 21.94 | 28.73 | 20.78 | 5.36 | 6.02 | 4.65 | 19.03 | 20.98 | 19.72 |

T₁: RDF (150:75:40 kg N: P₂O₅: K₂O ha⁻¹); T₂: 50% N through FYM + balance N, P and K through fertilizer; T₃: 50% N through Vermicompost + balance N, P and K through fertilizer; T₄: 50% N through poultry manure + balance N, P and K through fertilizer; T₅: 100% N through FYM + balance P and K through fertilizer; T₆: 100% N through Vermicompost + balance P and K through fertilizer; T₇: 100% N through poultry manure + balance P and K through fertilizer; T₈: RDF (150:75:40 kg N: P₂O₅: K₂O ha⁻¹) + 10 t FYM ha⁻¹

Table 2: Available nutrients in soil after harvest of baby corn as influenced by organic manures and fertilizers

| Treatments | Nitrogen (kg ha ⁻¹) | | | Phosphorus (P ₂ O ₅ : kg ha ⁻¹) | | | Potassium (K ₂ O : kg ha ⁻¹) | | |
|----------------|---------------------------------|--------|--------|---|-------|--------|---|--------|--------|
| | 2008 | 2009 | Pooled | 2008 | 2009 | Pooled | 2008 | 2009 | Pooled |
| T ₁ | 187.81 | 172.14 | 179.97 | 16.47 | 14.62 | 15.55 | 143.43 | 141.36 | 142.4 |
| T ₂ | 217.48 | 222.78 | 220.13 | 25.56 | 27.02 | 26.29 | 184.29 | 192.73 | 188.51 |
| T ₃ | 214.55 | 218.95 | 216.75 | 28.25 | 30.87 | 29.56 | 179.85 | 188.44 | 184.15 |
| T ₄ | 215.3 | 219.71 | 217.5 | 29.48 | 31.96 | 30.72 | 171.1 | 182.01 | 176.55 |
| T ₅ | 232.4 | 234.7 | 233.55 | 32.74 | 33.33 | 33.03 | 220.91 | 232.69 | 226.8 |
| T ₆ | 224.07 | 227.33 | 225.7 | 34.22 | 37.62 | 35.92 | 206.92 | 218.96 | 212.94 |
| T ₇ | 227.84 | 230.29 | 229.07 | 36.33 | 39.7 | 38.02 | 193.88 | 205.44 | 199.66 |
| T ₈ | 213.52 | 215.85 | 214.69 | 24.44 | 25.14 | 24.79 | 175.59 | 178.47 | 177.03 |
| S.Em + | 3.97 | 4.19 | 3.69 | 0.93 | 2.08 | 1.24 | 3.58 | 5.17 | 1.68 |
| C.D at 5 % | 12.03 | 12.7 | 11.21 | 2.82 | 6.32 | 3.76 | 10.86 | 15.67 | 5.09 |

T₁: RDF (150:75:40 kg N: P₂O₅:K₂O ha⁻¹); T₂: 50% N through FYM + balance N, P and K through fertilizer; T₃: 50% N through Vermicompost + balance N, P and K through fertilizer; T₄: 50% N through poultry manure + balance N, P and K through fertilizer; T₅: 100% N through FYM + balance P and K through fertilizer; T₆: 100% N through Vermicompost + balance P and K through fertilizer; T₇: 100% N through poultry manure + balance P and K through fertilizer; T₈: RDF (150:75:40 kg N: P₂O₅:K₂O ha⁻¹) + 10 t FYM ha⁻¹

Table 3: Chemical properties of soil after harvest of baby corn as influenced by organic manures and fertilizers

| Treatments | pH | | | EC (dS m ⁻²) | | | Organic Carbon (%) | | |
|----------------|------|------|--------|--------------------------|------|--------|--------------------|------|--------|
| | 2008 | 2009 | Pooled | 2008 | 2009 | Pooled | 2008 | 2009 | Pooled |
| T ₁ | 6.73 | 6.69 | 6.71 | 0.14 | 0.15 | 0.15 | 0.39 | 0.31 | 0.35 |
| T ₂ | 6.85 | 6.87 | 6.86 | 0.19 | 0.16 | 0.18 | 0.56 | 0.58 | 0.57 |
| T ₃ | 6.83 | 6.84 | 6.84 | 0.21 | 0.17 | 0.19 | 0.52 | 0.55 | 0.54 |
| T ₄ | 6.82 | 6.85 | 6.84 | 0.24 | 0.18 | 0.21 | 0.5 | 0.53 | 0.52 |
| T ₅ | 6.94 | 6.95 | 6.95 | 0.21 | 0.2 | 0.21 | 0.74 | 0.77 | 0.76 |
| T ₆ | 6.89 | 6.91 | 6.9 | 0.23 | 0.21 | 0.22 | 0.65 | 0.68 | 0.66 |
| T ₇ | 6.88 | 6.89 | 6.89 | 0.26 | 0.24 | 0.25 | 0.61 | 0.64 | 0.63 |
| T ₈ | 6.79 | 6.81 | 6.8 | 0.16 | 0.14 | 0.15 | 0.54 | 0.57 | 0.56 |
| S.Em + | 0.03 | 0.03 | 0.03 | 0.02 | 0.01 | 0.01 | 0.02 | 0.02 | 0.01 |
| C.D at 5 % | 0.11 | 0.1 | 0.08 | 0.05 | 0.04 | 0.03 | 0.05 | 0.07 | 0.04 |

T₁: RDF (150:75:40 kg N: P₂O₅:K₂O ha⁻¹); T₂: 50% N through FYM + balance N, P and K through fertilizer; T₃: 50% N through Vermicompost + balance N, P and K through fertilizer; T₄: 50% N through poultry manure + balance N, P and K through fertilizer; T₅: 100% N through FYM + balance P and K through fertilizer; T₆: 100% N through Vermicompost + balance P and K through fertilizer; T₇: 100% N through poultry manure + balance P and K through fertilizer; T₈: RDF (150:75:40 kg N: P₂O₅:K₂O ha⁻¹) + 10 t FYM ha⁻¹

Table 4: Yield and yield parameters of baby corn as influenced by organic manures and fertilizers

| Treatments | Weight of baby corn (g) | | | Baby corn yield (q ha ⁻¹) | | | Fodder yield (t ha ⁻¹) | | |
|----------------|-------------------------|-------|--------|---------------------------------------|-------|--------|------------------------------------|-------|--------|
| | 2008 | 2009 | Pooled | 2008 | 2009 | Pooled | 2008 | 2009 | Pooled |
| T ₁ | 17.64 | 17.86 | 17.75 | 15.4 | 17.33 | 16.37 | 33.25 | 35.34 | 34.29 |
| T ₂ | 16.28 | 16.45 | 16.36 | 12.5 | 13.67 | 13.08 | 27.93 | 29.75 | 28.84 |
| T ₃ | 17.26 | 18.12 | 17.69 | 14.88 | 15.78 | 15.33 | 31.09 | 33.23 | 32.16 |
| T ₄ | 16.78 | 17.05 | 16.92 | 13.13 | 14 | 13.57 | 28.23 | 30.23 | 29.23 |
| T ₅ | 14.73 | 15.35 | 15.04 | 9.83 | 10.82 | 10.33 | 24.57 | 26.82 | 25.69 |
| T ₆ | 15.78 | 16.15 | 15.97 | 11.5 | 13.12 | 12.31 | 26.92 | 28.87 | 27.89 |
| T ₇ | 15.24 | 15.87 | 15.56 | 10.57 | 11.56 | 11.06 | 26.34 | 27.32 | 26.83 |
| T ₈ | 17.97 | 18.37 | 18.17 | 16.59 | 18.76 | 17.67 | 34.81 | 38.24 | 36.53 |
| S.Em + | 0.67 | 0.5 | 0.46 | 0.62 | 1.33 | 0.72 | 1.63 | 1.74 | 1.1 |
| C.D at 5 % | 2.04 | 1.51 | 1.39 | 1.88 | 4.02 | 2.2 | 4.93 | 5.28 | 3.33 |

T₁: RDF (150:75:40 kg N: P₂O₅:K₂O ha⁻¹); T₂: 50% N through FYM + balance N, P and K through fertilizer; T₃: 50% N through Vermicompost + balance N, P and K through fertilizer; T₄: 50% N through poultry manure + balance N, P and K through fertilizer; T₅: 100% N through FYM + balance P and K through fertilizer; T₆: 100% N through Vermicompost + balance P and K through fertilizer; T₇: 100% N through poultry manure + balance P and K through fertilizer; T₈: RDF (150:75:40 kg N: P₂O₅:K₂O ha⁻¹) + 10 t FYM ha⁻¹

DAS: Days after sowing

RDF: Recommended dose of fertilizer

Effect of organic manures and fertilizers on chemical properties of soil

pH and organic carbon content in soil after the harvest of baby corn crop (Table 3) differed significantly due to various treatments. Significantly highest pH and organic carbon content in soil were observed in the treatment with the application of 100 per cent N through FYM and balance P and K through fertilizer (6.95 and 0.76 %, respectively). The lowest pH and organic carbon content in soil were recorded in the treatment with the application of only recommended dose of fertilizer. The EC (Electrical conductivity) of the soil after the harvest of baby corn crop was significantly influenced

by different treatments. The highest EC (0.25 dSm⁻¹) was recorded by 100 per cent N through poultry manure and balance P and K through fertilizer and which was on par with 100 per cent N through vermicompost and balance P and K through fertilizer. Whereas, the lowest EC was observed in the treatment with the application of only recommended dose of fertilizer (0.15 dSm⁻¹). However, application of organic manures alone or in combination with chemical fertilizers recorded highest organic carbon content, pH and EC of the soil as compared to application of only recommended dose of fertilizer. This indicates that organic manures have predominant role in the improvement of soil fertility, physical-

chemical properties and biological activity, besides its nutrient contribution (Kulvinder *et al.*, 2005).

Effect of organic manures and fertilizers on yield parameters of baby corn

The pooled data on yield is presented in Tables 4 Results revealed that, significantly higher baby corn yield (17.69 q ha⁻¹) and green fodder yield (36.53 t ha⁻¹) were recorded in the treatment with application of recommended dose of fertilizer (150:75:40 kg NPK ha⁻¹) + FYM (10 t ha⁻¹) compared to other treatments. The extent of increase in baby corn yield was 7.94 per cent over the treatment which received only recommended dose of fertilizer (150:75:40 kg NPK ha⁻¹). The increase in yield might be due to increase in yield parameters of baby corn such as length of baby corn, girth of baby corn, weight of baby corn and number of babies per plant. The results are conformity with the findings of Natarajan (1990) and Shashidhara *et al.* (1998), where the increase in yield might be due to availability of nutrients coinciding with physiological needs of the crop, effective partitioning of the assimilates to sink that helps in higher dry matter accumulation and increase in yield (Ranjan and Preethi, 2017).

The increase in green fodder yield of baby corn recorded in this treatment was attributed to increase in plant height, number of leaves, leaf area, leaf area index and total dry matter production. The leaf area was correlated to increased fodder yield, since leaf area is an indicative of the assimilatory surface area providing the synthesis and accumulation of more photosynthates and dry matter production. These results are in conformity with the findings of Negalur (2000) and Eajas (2017)

Among the graded levels organic manures, significantly higher yield of baby corn were recorded due to application of 50 per cent N equivalent through various source of organic manure like, vermicompost, FYM and poultry manure in combination with balance NPK through inorganic fertilizers and all these treatments were on par. Application of 100 per cent N equivalent through organic manures like FYM, vermicompost and poultry manure in combination with balance P and K through commercial inorganic fertilizers recorded comparatively less yield compared to other combination of organic manures and fertilizers.

The above study indicated that baby corn responded well to combined application of organics and inorganics, especially to vermicompost which might be owing to favourable effect on soil condition and synchronized release of plant nutrients throughout the crop growth period and inorganic nutrients have positive influence on source-sink relationship as evident from remarkable improvement in plant height and dry matter accumulation and ultimately yields were increased. The results

are in conformity with the findings of Dadarwal *et al.* (2009).

REFERENCES

- Arun Kumar, K., Karuna Sagar, G., Chandrika, V. and Reddy, P. M. 2009. Influence of integrated nitrogen management on yield, nitrogen uptake, soil fertility status and economics of baby corn. *Indian J. Agric. Res.* **43(3)**: 227-229.
- Das, S., Ghosh, G., Kaleem, M.D. and Bahadur, V. 2008. Effect of different levels of nitrogen and crop geometry on the growth, yield and quality of baby corn (*Zea mays* L.) CV. 'GOLDEN BABY'. ISHS Acta Horticulturariae 809: International Symposium on the Socio-Economic Impact of Modern Vegetable Production Technology in Tropical Asia.
- Dadarwal, R.S., Jain, N.K. and Singh, D. 2009. Integrated nutrient management in baby corn. *Indian J. Agri. Sci.* **79(12)**: 1023-1025.
- Eajas Ahmed Dar, Abrar Yousuf, Mohammad Amin Bhat, Todarmal Poonia. 2017. Growth, Yield and quality of baby corn (*Zea mays* L.) and its effect on fodder as influenced by crop geometry and nitrogen application – A review, *The Bioscan.* **12(1)**:463-469
- Karki, T. B. and Ashok Kumar. 2005. Productivity potentials and economics of maize as affected by various fertility levels. *Ann. Agric. New Series.* **26(2)**: 340-341.
- Kulvinder, K., Krishan, Kapoor, K., Anand and Gupta, P. 2005. Impact of organic manures with and without mineral fertilizers on soil chemical and biological properties under tropical conditions. *J. Plant Nutr. Soil Sci.* **16(8)**: 117-122.
- Muthukumar, V.B., Velayudham, K. and Thavaprakash, N. 2007. Plant growth regulators and split application of nitrogen improves the quality parameters and green cob yield of baby corn (*Zea mays* L.). *J. Agron.* **6(1)**: 208-211.
- Natarajan, S. 1990. Standardization of nitrogen application for chilli *Capsicum annum* L.) grown under semidry conditions, South Indian Hort. **38**: 170-174.
- Negalur, R. B. 2000. Response of kharif pop sorghum (*Sorghum bicolor* L. Moench) genotypes to farm yard manure and mineral fertilizer in black soil under rainfed conditions. M.Sc. (Agri.) Thesis, University of Agricultural Sciences, Dharwad
- Pandey, A.K., Prakash, V., Mani, V.P. and Singh, R.D. 2000. Effect of rate of nitrogen and time of application on yield and economics of Baby corn. *Indian J. Agron.* **45(2)**: 338-343.
- Pansee, V.G. and Sukhatme, P.V. 1967. Statistical methods for agricultural workers. Indian Council of Agricultural Research, New Delhi, p. 347.
- Ranjana Verma and Preethi Choudhary. 2017. Nutritional Quality Evaluation of Organically vis-a-vis Conventionally grown Broccoli (*Brassica oleracea*). *The Bioscan.* **12(1)**:503-509.
- Shashidhara, G. B., Basavaraja, P.K., Basavarajappa, R., Jagadeesha, R.C and Nadagowda, V.B. 1998. Response of chilli to intercropping systems in red soils. In: *Water and Nutrient Management for sustainable production and quality of spice*, pp. 95-98.
- Thavaprakash, N., and Velayudham, K. 2008. Light interception and productivity of baby corn as influenced by crop geometry, intercropping systems and INM practices. *Asian J. Scientific Res.*, **1(1)**:72-78.



**NATIONAL ENVIRONMENTALISTS
ASSOCIATION**

The National Environmentalists Association is chartered in Ranchi as a nonprofit scientific and educational association of like minded academician, researchers, scientists from all over the nation for the furtherance and diffusion of knowledge of Life Sciences in general and Environmental Science in particular.

The association not only honours its members but also provides FELLOWSHIP to outstanding contributors to the subject and the society.

Contact :

For Editorial Information

Prof. M. P. Sinha
Vice Chancellor
Sido Kanhu Murmu University
Dumka - 814 110
Jharkhand, INDIA

For information regarding Association :

SECRETARY,
National Environmentalists Association,
D-13, Sai Roofs, 1st Floor,
H. H. Colony,
Ranchi - 834002
Jharkhand, India

E-mails : editor.bioscan@gmail.com
dr.mp.sinha@gmail.com
nat.env.assoc@gmail.com

Cell : 94313-60645; 9572649448

Ph. : 0651-2244071

Website : www.thebioscan.com
: www.neaindia.org

NAAS Rating : 5.26

U.S.A. Office

2827 Videre Dr.,
Wilmington,
DE 19808 We, USA

Type setter Bandana Solutions Facility Management LLP
Published by Aditi Publications, Patliputra, Patna

The Bioscan

An International Quarterly Journal of Life Sciences

ISSN : 0973-7049

Volume 15(3) : 2020

Published as an official organ by

NATIONAL ENVIRONMENTALISTS ASSOCIATION

CONTENTS

Page

A. RESEARCH PAPER

1. Comparison and molecular profiling of Begomovirus infecting chilli (*Capsicum annum*) in gangetic alluvial zone of West Bengal
Uday Bikash Oraono, Lourembam Sanajaoba Singh and Jayanta Tarafdar—275 - 280
2. Assessment of probiotic characteristics of *L.plantarum*
Sravani Kandula And Rita Narayanan—281 - 286
3. Effect of rate and frequency of micronutrient on growth attributes and dry matter yield of Banana Cv. grand naine under south Gujarat condition
Narendra Singh, Sonal Tripathi, Patel V. A., Jaimin Naik and Chauhan Aditi—287 - 290
4. Character association and path analysis for seed yield and its components in Grass pea (*Lathyrus sativus* L.)
Gangishetti Ranjithkumar, Sandip Debnath and Duddukur Rajasekhar—291 - 295
5. Determination of physical and biometric properties of onion bulbs in relation to design of digger cum windrower
Shiddanagouda Yadachi and Kiran Nagajjanavar—297 - 301
6. Polygenic variation for morphological and biochemical traits of brinjal genotypes (*Solanum melongena* L.) and its wild relatives
Nisha Sharma, K. D. Bhutia, Rajesh Kumar, Sita Kumari Prasad, Ankita Debnath and Malay Marut Sharma—303 - 309
7. Effect of different sources of horizontal transmission of bacterial flacherie on Et_{50} for symptom expression and mortality of PM X CSR₂
B. L. Kavyashree., R. N. Bhaskar and C. Doreswamy—311 - 314
8. Comparative biology of *Goniozus nephantidis* (Muesbeck) on *Galleria mellonella* L. and *Corcyra cephalonica* (Stainton)
A. V. Desai, M. R. Siddhapara and N. P. Trivedi—315 - 321
9. Efficiency of ovatide on mass seed production of climbing perch (*Anabas testudineus*, Bloch, 1972) in Nalbari district, Assam
Ankur Rajbongshi, A. Ali, M. Chakravarty, M. Deka, H. Mazumdar, Pranab Kr Das and S. Baishya—323 - 326
10. Study of combining ability and gene action for yield and yield component characters in interspecific hybrids of cotton (*Gossypium hirsutum* L. X *Gossypium barbadense* L.)
S. B. Gohil., M. B. Parmar., M. P. Patel and D. A. Patel—327 - 333
11. Per oral inoculation of *Lysinibacillus sphaericus* with pathogenic microbes on rearing and cocoon parameters of silkworm, *Bombyx mori* L.
H. G. Anusha, R. N. Bhaskar and K. V. Anitharani—335 - 338
12. Effect of fungicides, plant extracts and bioagents on spore germination of *Colletotrichum lindemuthianum* causing field bean anthracnose

The Journal is Currently Abstracted / Indexed in

- Paryavarn Abstract, INDIA
- Indian Science Abstract, INDIA
- Cambridge Science Abstract, U.S.A.
- Zoological Record, U.K.
- Directory of Open Access Journal (DOAJ)
- Chemical Abstract, U. S. A.
- Research BIB
- Indian Science
- Journal Seek
- Scientific Indexing Service (SIS)
- Journal is currently rated by
- Index Copernicus
- Universal Impact Factor
- NAAS

DISCLAIMER

The Publisher and Editors cannot be held responsible for errors or any consequences arising from the use of information in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher/ Association and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher / Association and Editors of the products advertised.

- S. Narasimha Rao, S.L. Bhattiprolu, A. Vijaya Gopal and V. Sekhar— 339-344
13. Evaluation in vitro different fungicides for growth of *Rhizoctonia bataticola*
A.M. Kadam, S.S. Chavan and A.H. Kendre — 345- 349
 14. Evaluation of gladiolus varieties for flowering and cut flower traits under indo-gangetic plains
Girish, P. M., Anjana Sisodia and Anil K. Singh— 351- 355
 15. Effect of land configuration and different organic sources on growth, yield and quality of carrot under organic farming
B. Solanki, A. R. Kaswala, P.K. Dubey and A.P. Italiya— 357- 362
 16. Verification and usability analysis of medium range weather forecast for the Kokrajhar district of lower Brahmaputra valley zone of Assam
Kuldip Medhi, Kushal Sarmah, Vinod Upadhyay, Sunil Kumar Paul, Athar N. Islam and Bikash J. Gharphalia— 363 - 370
 17. Canonical root analysis and clustering for characterization and evaluation of aromatic rice germplasm based on morphological characters
G. Parimala, Ch. Damodhar Raju, L.V. Subba Rao and K. Uma Maheswari— 371 - 374
 18. Genetic divergence analysis of sesame genotypes (*Sesamum indicum* L.)
Dasari Rajitha, T. Srikanth, D. Padmaja and T. Kiran Babu— 375 - 379
 19. Nutrient uptake and chemical properties of soil after harvest of baby corn (*Zea mays* L.) as influenced by organic manures and fertilizers
D.H. Roopashree, S. Kamal Bai, Nagaraju and S. Raghavendra— 381 - 384
 20. Genotypic response on growth and yield in papaya
D. K. Varu., K. D. Patel and Sandip Makhmale— 385 - 389
 21. Studies on frequency distribution of yield and yield related traits in F₂M₂ generation of sesame (*Sesamum indicum* L.)
Rajesh Kumar Kar, Tapash Kumar Mishra and Banshidhar Pradhan — 391 - 395
 22. Correlation and path analysis in cowpea (*Vigna unguiculata* (L.) Walp)
R.M. Nagalakshmi, R. Usha Kumari and R. Ananda Kumar— 397 - 401
 23. A simple and efficient method for DNA extraction from rabi sorghum [*Sorghum bicolor* (L.) Moench]"
S. S. Gadakh, G. D. Khalekar., U. S. Dalvi, A. A. Kale and P.L. Kulwal— 403 - 406
 24. Determination of economic injury level (EIL) of sugarcane planthopper borer *Chilo tumidicostalis* Hampson (Lepidoptera: pyralidae)
R. K. Nath and D. K. Saikia— 407- 409
 25. Performance of different summer mung (*Vigna radiata* L.) varieties sown at different dates under Manipur valley condition
Meghna Gogoi, Jamkhogin Lhungdim, Kamal Kant, Urjashi Bhattacharya and Gauri Mohan — 411 - 414

ZONAL CO-ORDINATORS OF THE ASSOCIATION

- Prof. N. Behera
School of Life Science,
Sambalpur University
- Dr. Nirmal Kumar
ISTAR, Vallabh Vidyanagar,
Anand, Gujarat
- Dr. P. N. Sudha
D. K. M. College for Women,
Vellore
- Prof. S. P. S. Dutta
Dept. of Environmental Science,
Jammu University, Jammu
- Dr. V. Salom Gnana Thanga
Dept. of Env. Scs.,
University of Kerala, Kariavattom
Tiruvananthapuram, Kerala

Publications of the Association

The Bioscan

An International Quarterly Journal of Life Sciences

The Ecoscan

An International Quarterly Journal of
Environmental Sciences

Both the Journals are
online

Both the Journals are
available on Google.com

Websites of the Journals are
www.theecoscan.in
www.thebioscan.in

FEATURES OF ASSOCIATION

- Association is registered under 80G of I.T.
- Prestigious fellowship of the Association (F. N. E. A.) to academicians of the nation.
- Regular annual conference of national and international levels organized by the Association.
- Young Scientist and Senior Scientist award during the conference of the Association.



_____ 062 & 082

_____ 000 & 000

| THE BIOSCAN : SUBSCRIPTION RATES | | | | |
|---|--------------|----------------|--------------------|--------------------|
| | | India (Rs.) | SAARC Countries | Other Countries |
| Individuals | One Year | 1,000 | 2,000(I:C) | US \$200 |
| | Life Member* | 10,000 | | |
| Institutions | One Year | 3,000 | 6,000(I:C) | US \$400 |
| | Life Member* | 30,000 | | |

*Life Member will receive the journal for 15 years while other benefits will continue whole life

THE BIOSCAN : MEMBERSHIP FORM

Please enter my subscription for the above journal for the year / life member.

Name:

Address:

E-mail:

Payment Rs. : by DD / MD in favour of
National Environmentalists Association payable at Ranchi, No. Dated
..... is enclosed.

NOTE: FOR MEMBERSHIP THE ABOVE INFORMATION CAN BE SENT ON SEPARATE SHEET