

SOIL AND FOLIAR APPLICATION OF BORON ON WHEAT PRODUCTIVITY IN NORTH EAST ALLUVIAL PLAIN OF EASTERN INDIA

A. K. MAURIYA, V. K. MAURYA, R. K. VERMA, SUNITA KUSHWAHA AND RADHEY SHYAM

Krishi Vigyan Kendra, Purnea, Bihar Agricultural University Sabour - 813 210 (Bihar)

e-mail: ak.mauriya12@gmail.com

KEYWORDS

Boron
Economics
Nutrient uptake
Wheat and Yield

Received on :

28.06.2016

Accepted on :

07.07.2018

*Corresponding author

ABSTRACT

An on-farm trials was conducted during *rabi* season of 2013 in a randomized block design with six treatments on boron deficits soil at farmer's field by KVK Purnea (Bihar) to validate, refine and popularize the technology developed at Bihar Agricultural University Sabour, Bhagalpur (Bihar) for enhancing the productivity of wheat [*Triticum aestivum* (L.) emend. Fiori & Paol.]. Application of 2.0 Kg/ha B as basal followed by 0.2% foliar application of B at 2-5 % spike initiation gave significant increase in yield attributes of wheat as compared to farmer's practice (*i.e.* without boron application). The highest grain yield (3.58 t/ha) was observed where boron was applied 2.0 Kg/ha as basal fb 0.2% foliar application at 2-5 % spike initiation and it was most economical to increase wheat productivity.

INTRODUCTION

Wheat is the second important food crop of India. It is grown in 29.31 m ha with the production of 85.76 million tonnes and productivity of 2.27 t/ha and contributes about 40% in the total food grain production in the country (MOF, 2011). It is also grown in 21.54 lakh ha with the production of 35.70 lakh tonnes and productivity 16.57 q/ha in the Bihar, while 39.7 thousands ha wheat area covered by Purnea district with the productivity of only 6.03 q/ha during 2014-15 (*Directorate of Statistical and Evaluation, Bihar 2014-15*). Wheat is predominately grown in rice-wheat cropping system, and this system is considered backbone of food security in India (Yadav and Kumar, 2009). This system exhausts of the soil nutrients limiting crop production (Mauriya *et al.* 2015), especially for the high yielding varieties. Intensive cropping also renders deficiency of micro-nutrients in addition to major nutrients in the soil. The farmer's perception of nutrient management until now has mostly been the use of major nutrients like N, P and K. The imbalance use of inorganic fertilizer, in the absence of organic sources of nutrients, has lead to the micronutrient deficiency especially in rice and wheat restricting upward growth of productivity as well as quality (Mishra, 2012). Boron (B) deficiency has been realized as the second most important micronutrient constraints in crop productivity after that of zinc on global scale as well as in Indian agriculture (Arora *et al.* 1985, Patil *et al.*, 1987, Sakal *et al.*, 1988 and Sinha *et al.* 1991). The deficiency symptom of B in wheat leads to thinner stems, shorter and fewer tillers, and failure to produce viable seeds (Marschner, 1995). Boron has been found to play a key

role in pollen germination and pollen tube growth, stimulate the plasma membrane, anther development, floret fertility and seed development (Wang *et al.* 2003). Boron deficient stems and leaves are brittle, whereas, B sufficient stems and leaves are flaccid. The soils of eastern states of India namely Assam, Bihar, Meghalaya, West Bengal, Jharkhand and Orissa are deficit in boron. In India, boron deficiency was initially reported 2% in the year 1980 (Katyal and Vlek 1985), which has now increased to 68% in red soils of Bihar (Sakal *et al.* 2002), with mean of 52% (Singh, 2012) for the whole country. Rice and wheat, the two most important crops in Purnea district of Bihar are sensitive to B deficiency, however, substantial yield reduction has been observed under B deficiency. Due to continuous growing of boron exhaustive crops, high pH and calcium compounds in soils, leaching and without B application to the crops and high humidity in atmosphere most of the soils are B deficient in that area. To cope with B deficiency, it can be delivered as soil and foliar application. Foliar applied B plants retain significant phloem mobility to flowering parts from senescing leaves. Therefore, foliar sprays of B not only provide a means to apply B at a particular growth stage, but also remove B deficiency.

Keeping in view the importance of boron in increasing crop yield, the present study was conducted to investigate the effects of soil and foliar application of boron on wheat yield in north eastern plain zone of India.

MATERIALS AND METHODS

Initially Participatory Rural Appraisal (PRA) was done to

identify causes of low yield of wheat and high cost of production in late sowing condition. It was found that boron deficiency in upland situation is one of the serious constraints in Koshi region of Bihar. Therefore, based on PRA two locations of five farmer's field in two locations at Purnea were selected for on-farm trials at farmer fields during the *rabi* season of 2013. Purnea district lies in the Gangaic Alluvial plain with 25°13'80" to 27°7'59" N latitude, 86°59'6" to 87°52'35" E longitude and 52.3 m above mean sea level with annual rainfall 1412 mm. The cold season commences in November when both day and night temperature drop rapidly and lasts till February with humid climate. January is the coldest month, with mean daily minimum temperature in the range of 5-10°C and mean daily maximum in the range of 20-25°C. Humidity is generally above 70% except in the first half of the summer season. Soil sample of all selected sites (at five farmer's field) were collected and analyzed. The soil of experiment sites were sandy loam to loamy having alluvial properties, with pH 6.2, organic carbon 0.39%, available N 246.7 kg/ha, P₂O₅ 42.2 kg/ha, K₂O 180.3 kg/ha and available boron 7.0 ppm. The experiment comprised six treatments viz. T₁- Farmer's practice (No application of boron), T₂- 1.0 Kg/ha B application as basal, T₃- 2.0 Kg/ha B application as basal, T₄- 0.2% foliar application of B at 2-5 % spike initiation, T₅- 1.0 Kg/ha B application as basal + 0.2% foliar application of B at 2-5 % spike initiation and T₆- 2.0 Kg/ha B application as basal + 0.2% foliar application of B at 2-5 % spike initiation. Treatments were arranged in randomized block design (RBD) with five replications in which one farmer was treated as one replication. The area of each experimental site was 0.40 ha. Wheat variety HI 1563 was sown after the harvesting of rice in the first week of December during at the spacing of 18 cm apart at the depth of 3-4 cm with the seed rate of 150 kg/ha. A uniform application of recommended dose of fertilizers 80:40:20 kg/ha N, P₂O₅ and K₂O were applied through urea, di-ammonium phosphate and muriate of potash. Half dose of nitrogen and full dose of

phosphorus and potash were drilled as basally at the time of sowing; remaining dose of nitrogen was applied in two equal splits as top dressing at 25 and 50 days after sowing. Full dose of Boron those applying in soil as per treatments (*i.e.* 1.0 and 2.0 kg/ha) was applied at the time of first ploughing as broadcasting and 0.2% foliar application at 2-5 % spike initiation stage through borax (11% B). Crop was manually harvested in the first fortnight of April. The grain yield data were recorded and adjusted to 13% of the moisture content. The harvest index (%) was calculated according to the following formula.

$$\text{Harvest index (\%)} = \frac{\text{Grain yield} \times 100}{\text{Biological yield (grain yield + straw yield)}}$$

Crop was raised using recommended cultural practices. After harvest, treatment wise grain and straw samples were collected for N, P and K nutrient analysis/nutrient uptake. Total nutrient uptake (grain + straw) in wheat was determined by multiplying the N, P and K concentration with corresponding grain and straw yields, respectively and expressed in kg/ha or g/ha. The statistical analysis of the data was carried out by using standard statistical methods of analysis of variance (Panse and Sukhatme, 1985).

RESULTS AND DISCUSSION

Yield attributes

Yield-attributing characters viz. plant height, spikes/m², spike length, grains/spike and 1,000-grain weight were positively affected by boron (B) application as soil/foliar. Crop yield mainly depends upon many yield attributing characters.

Among them number of fertile tillers (spikes/m²) are very important because higher the number of fertile tillers/m² higher will be the final yield of the crop. Application of 2.0 Kg/ha B as basal followed by 0.2% foliar application of B at 2-5 % spike initiation gave the highest values of yield attributes of wheat.

Table 1: Effect of boron application on yield attributes, yield and economics of wheat

Treatment	Plant height (cm)	Spikes/m ²	Spike length (cm)	Grains/Spike	1000 grain weight (g)	Grain Yield (t/ha)	Straw yield (t/ha)	HI (%)
Farmer's practice (No B application)	93.6	256	9.4	31.7	33.6	3.11	4.28	42.1
1.0 Kg/ha B as basal	97.1	271	9.8	38.4	38.9	3.26(4.8)	4.30	43.1
2.0 Kg/ha B as basal	99.0	292	10.4	42.0	41.5	3.55(14.1)	4.39	44.7
0.2% foliar application of B at 2-5 % spike initiation	96.8	253	10.2	38.2	38.0	3.20(2.9)	4.24	43.0
1.0 Kg/ha B as basal + 0.2% foliar application of B at 2-5 % spike initiation	97.3	276	10.4	41.1	40.7	3.47(11.6)	4.41	43.2
2.0 Kg/ha B as basal + 0.2% foliar application of B at 2-5 % spike initiation	99.8	300	10.8	43.2	42.6	3.58(15.1)	4.52	44.2
SEm±	0.66	7.1	0.22	0.80	0.74	0.093	0.119	
CD (P=0.05)	1.9	16.4	0.62	2.31	2.15	0.28	0.33	

Note: Figures in parentheses are percent increase in grain yield relative to farmer's practice (without boron application)

Table 2: Nutrients uptake by grain, straw and total as influenced by different dose and application method of boron in wheat

Treatment	Nitrogen (kg/ha)			P (kg/ha)			K (kg/ha)		
	Grain	Straw	Total	Grain	Straw	Total	Grain	Straw	Total
Farmer's practice (No B application)	53.18	20.80	73.98	10.92	3.06	13.98	10.23	53.86	64.09
1.0 Kg/ha B as basal	56.72	22.79	79.51	11.08	3.35	14.43	11.08	59.77	70.85
2.0 Kg/ha B as basal	62.13	26.34	88.47	14.56	3.93	18.49	13.14	67.54	80.68
0.2% foliar application of B at 2-5 % spike initiation	54.72	22.89	77.61	10.88	3.39	14.27	10.88	57.66	68.54
1.0 Kg/ha B as basal + 0.2% foliar application of B at 2-5 % spike initiation	60.38	24.79	85.17	13.53	3.76	17.29	12.15	64.26	76.41
2.0 Kg/ha B as basal + 0.2% foliar application of B at 2-5 % spike initiation	63.37	27.01	90.38	15.04	4.08	19.12	13.60	71.20	84.8
SEm±	2.03	1.38	3.26	0.99	0.21	1.29	0.72	3.60	4.15
CD (P=0.05)	5.88	3.94	9.36	2.87	0.61	3.74	2.11	10.47	11.9




Figure 1: Relationship between yield attributes and yield of wheat

Application of boron as soil/foliar or both have shown significant improvement in yield attributes as compared to farmer's practice (*i.e.* without boron application). Application of 1.0 Kg/ha B as basal followed by 0.2% foliar application of B at 2-5 % spike initiation was statistically non-significant with the application of 2.0 Kg/ha B as basal alone. Grain yield was positively and significantly correlated with growth and yield attributes, namely plant height ($R^2 = 0.800$), spikes/m² ($R^2 = 0.887$), spike length ($R^2 = 0.778$), grains/spike ($R^2 = 0.841$) and 1,000-grain weight ($R^2 = 0.872$). This may be due to the reason that application of B influence the vegetative growth,

flowering, longevity of spike because boron is an essential element found in the meristematic regions of plants such as root tips, emerging leaves and bud (Sharma *et al.*, 2013; Kumar *et al.*, 2015 and Singh *et al.*, 2015). Boron plays a vital role in grain setting of wheat. So, the supply of Boron at this stage helps in grain filling and ultimately sterility is reduced and number of grains per spike increased. The similar results and observations were also reported by Mitra and Jana (1991), Tahir *et al.* (2009) and Debnath and Ghosh (2011).

Grain and straw yield

Grain and straw yield of wheat was positively affected by boron application as soil or foliar or both methods. Grain yield production was significantly increased due to boron application. The maximum grain yield (3.58 t/ha) was observed in treatment where boron was applied 2.0 Kg/ha as basal followed by 0.2% foliar application at 2-5 % spike initiation which was statistically at par with the treatment where boron was applied 2.0 Kg/ha as basal and 1.0 Kg/ha as basal followed by 0.2% foliar application at 2-5 % spike initiation. However, minimum grain yield (3.11 t/ha) was observed in farmer's practice *i.e.* without boron application which was statistically at par with 1.0 kg/ha boron application as basal and 0.2% foliar application at 2-5 % spike initiation alone. Grain yield of wheat crop is the result of combined effect of various yield-attributing characters. Grain yield increased significantly due to the application of boron by which enhanced pollen tube germination and grain setting. As the boron requirement of male reproductive organ, the anthers was also greater than

carpel, since the boron requirement in the anthers for successful fertilization was met by of boron application as basal and foliar at spike initiation stage so the grain yield was higher than control. Mishra *et al.* (1989), Tahir *et al.* (2009), Ganie *et al.* (2014) and Layek (2014) also reported similar results.

The ability of wheat crop plant to convert the dry matter into grain yield of wheat is indicated harvest index. Higher the harvest index, greater will be the physiological potential for converting dry matter into grain. Harvest index (H.I.) was significantly affected by the application of Boron. The maximum H.I. (40.11%) was observed in case where Boron was applied at anthesis which is statistically at par with all other treatments where Boron was applied followed by control. These results are supported by the findings of Alam *et al.* (2000).

Nutrient uptake

Total (Grain and straw) nutrient uptake by wheat crop plant at harvest stage was positively affected by boron application as soil or foliar or both methods. Nutrient uptake was significantly increased due to boron application. Maximum total nutrients uptake (total N, P and K) was analysed with the application of 2.0 Kg/ha B as basal followed by 0.2% foliar application of B at 2-5 % spike initiation, which was statistically at par with application of 2.0 Kg/ha B as basal and 1.0 Kg/ha B as basal followed by 0.2% B foliar application at 2-5 % spike initiation. However, minimum total nutrient uptake was observed in farmer's practice *i.e.* without boron application. Nutrient uptake is a product of its content in plant and dry matter accumulation, hence, it follows the same pattern as for growth particularly grain and straw yields. It might be attributed to the vigorous growth of plants due to better absorption of nutrients from the plots supplied with balanced nutrients. The results are in conformity with those of Kapoor *et al.* (2016).

It may be concluded that application of 2.0 Kg/ha B as basal + 0.2% foliar B at 2-5 % spike initiation in wheat was more productive as compared to farmers practice (No boron).

REFERENCES

- Alam, S.M., Iqbal, Z. and Latif, A. 2000. Effect of boron application with or without zinc on the yield of wheat. *Pakistan J. Soil Science* **18**: 95-98.
- Arora, C. L., Singh, B. and Takkar, P. N. 1985. Secondary and micronutrient deficiency in crops. *Progressive Farming*. **XXI** **8**: 13.
- Debnath, P. and Ghosh, S. K. 2011. Determination of critical limit of available boron for rice in Terai zone soils of West Bengal. *J. The Indian Society of Soil Science*. **59**: 82-86.
- Directorate of Statistical and Evaluation, Bihar 2014-15. <http://www.dse.nic.in> > publication > bihar.
- Ganie, Mumtaz A., Akhter, Farida, Bhat, M. A. and Najjar, G. R. 2014. Growth, yield and quality of french bean (*Phaseolus vulgaris* L.) as influenced by sulphur and boron application on Inceptisols of Kashmir. *The Bioscan*. **9**(2): 513-518.
- Kapoor Sapna, Sharma Sanjay K, Rana, S. S. and Shankhyan Narender 2016. Effect of the Application of Nitrogen, Zinc and Boron on Micro-Nutrients Concentration and Uptake in Grain and Straw of Wheat in a Silty Clay Loam Soil of Mid Hills. *International Journal of Advances in Agricultural Science and Technology*. **3**(6): 25-39.
- Katyal, J. C. and Vlek, P. L. G. 1985. Micronutrient problem in tropical Asia. *Fertilizer Research* **7**: 69-94.
- Kumar, J., Kumar, R., Rai, Ratna and Mishra, D. S. 2015. Response of 'Pant Prabhat' guava trees to foliar sprays of zinc, boro, calcium and potassium at different plant growth stages. *The Bioscan*. **10**(2): 495-498.
- Layek, J., Shivakumar, B. G., Rana, D. S., Gangaiah, B., Lakshman, K. and Paramanik, B. 2014. Growth, nodulation, physiological indices and yield of soybean as influenced by sulphur and boron nutrition. *The Bioscan*. **9**(4): 1389-1393.
- Marschner, H. 1995. Mineral nutrition of higher plants. 2nd Edition New York: Academic Press, pp. 889-1995.
- Mauriya, A. K., Maurya, V. K., Mishra, P. K. and Verma, R. K. 2015. Site-specific primary nutrients recommendation in wheat (*Triticum aestivum* L.) for sustainable productivity and increasing profitability. *J. Soil and Soil Conservation*. **14**(3): 241-247.
- Mishra, L. K. 2012. Effect of phosphorus and zinc fertilization on biochemical composition of wheat. *The Bioscan*. **7**(3): 445-449.
- Mishra, S. S., Gulati, J. M. L., Nanda, S. S., Garnayk, L. M. and Jana, S. N. 1989. Micronutrient studies in wheat. *Orissa J. Agriculture Research*. **22**(2): 94-96.
- Mitra, A. K. and Jana, P. K. 1991. Effect of doses and method of boron application on wheat in acid trail soils of North Bengal. *Indian J. Agronomy*. **36**(1): 72-74.
- MOF. 2011. Economic Survey 2011-12 (In): Statistical Tables, Selected Indicators 1950-51 to 2011-12. *Ministry of Finance, Government of India*.
- Panse, V. G. and Sukhatme, P. V. 1967. Statistical methods for agricultural workers. *ICAR publication*. New Delhi.
- Patil, G. D., Patil, M. D., Patil, N. D. and Adsule, R. N. 1987. Effect of boronated superphosphate, single superphosphate and borax on yield and quality of groundnut. *J. Maharashtra Agriculture University*. **12**: 168-170.
- Sakal, R., Singh, A. P., Sinha, R. B. and Bhogal, N. S. 1988. Annual progress reports. ICAR All India Coordinated Scheme of Micro and Secondary Nutrients in Soils and crops of Bihar, *Research Bulletin, Department of Soil Science, RAU, Samastipur, Bihar, India*.
- Sakal, R., Singh, A. P. and Sinha, R. B. 2002. Evaluation of rate and frequency of B application in cropping systems. *Fertiliser News*. **47**(10): 37-41.
- Sharma, J., Gupta, A. K., Kumar, C. and Gautam, R. K. S. 2013. Influence of zinc, calcium and boron on vegetative and flowering parameters of gladiolus cv. Aldebran. *The Bioscan* **8**(4):1153-1158.
- Singh, M., Sirothia, P., Bhat, M.A., Tiwari, P. and Namdeo, A. (2015). Growth and yield of mungbean (*Vigna radiata* L.) in response to the application of sulphur and boron under rainfed condition. *The Bioscan*. **10**(4): 1665-1669.
- Singh, M. V. 2012. Spread of micro-nutrient deficiencies specially boron in India and response of field crops. "Brain storming workshop on soil test based nutrients including boron and other micro nutrients" Organized by ICRISAT - Agriculture Directorate, Karnataka -Rio Tinto India at Bangalore, India.
- Sinha, R. B., Sakal, R., Singh, A. P., Bhogal, N. S. 1991. Response of some field crops to boron application in calcareous soils. *J. Indian Society of Soil Science*. **39**: 342-345.
- Tahir, M., Tanveer, A., Shah, T. H., Fiaz, N. and Wasaya, A. 2009. Yield response of wheat (*Triticum aestivum* L.) to boron application at different growth stages. *Pakistan Journal of Life and Social Sciences* **7**(1): 39-42.
- Wang, Q., Longdou, L. U., Xiaoqin, W. U., Yiqin, L. I. and Jinxing, L. I. N. 2003. Boron influences pollen germination and pollen tube growth in *Picea meyeri*. *Tree Physiology*. **23**: 345-351.
- Yadav, D. S. and Kumar, A. 2009. Long-term effect of nutrient management on soil health and productivity of rice (*Oryzasativa*)-wheat (*Triticumaestivum*) System. *Indian J. Agronomy*. **54**(1): 15-23.