

**“KNOWLEDGE AND PRACTICE REGARDING SAFETY STANDARDS OF ORAL RADIOLOGY AMONG DENTAL PRACTITIONERS IN DENTAL COLLEGE, BIHAR”
CORRESPONDING AUTHOR - DR BINU SINGH**

Author :- DR. LIPI SINGH (Post Graduate, Buddha Institute of Dental Sciences and Hospital , Patna, Bihar, 800026)

Email id - lipisinghh@gmail.com

Phone no. - 8298198444

1. **Author :- DR. LIPI SINGH** (Post Graduate, Buddha Institute of Dental Sciences and Hospital , Patna, Bihar, 800026)

Email id - lipisinghh@gmail.com

Phone no. - 8298198444

2. **Co - authors :- DR. KUMAR ANAND** (Professor, Buddha Institute of Dental Sciences and Hospital , Patna, Bihar, 800026)

Email id - drkumaranand.anand@gmail.com

Phone no. - 9835013078

3. **Co - authors :- DR. ARCHANA SUDHEER** (Professor & HOD, Buddha Institute of Dental Sciences and Hospital , Patna, Bihar, 800026)

Email id - DRARCHANASUDHEER@gmail.com

phone no. - 9739577540

4. **Co - author - DR. INDRANIL MUKHOPADHYAY** (Senior Lecturer, Buddha Institute of Dental Sciences and Hospital), Patna, Bihar, 800026)

Email id - dsindra77@gmail.com

Phone no. - 7044171244

5. **Co – author - DR. BABITA KUMARI** (Postgraduate, Buddha Institute of Dental Sciences and Hospital , Patna, Bihar, 800026)

Phone no. - 828752081

Email id - Babitakumari1996@gmail.com

6. **Co – author - DR. BINU SINGH** (Postgraduate, Buddha Institute of Dental Sciences and Hospital), Patna, Bihar, 800026)

Phone no. - 8126576516

Email id - neet.binumds@gmail.com

Correspondence author - DR. LIPI SINGH (Postgraduate, Buddha Institute of Dental Sciences and Hospital), Patna, Bihar, 800026 Phone no. - 8298198444

Email id - lipisinghh@gmail.com

DOI: 10.63001/tbs.2025.v20.i04.pp1041-1061

KEYWORDS

Dental practitioners, Digital Sensor, X-Ray film

Received on:

30-09-2025

Accepted on:

05-11-2025

Published on:

09-12-2025

Abstract

In Dentistry, radiographic evaluation is required but professional negligence or ignorance can lead to radiation hazards so the knowledge regarding safety standards among Dentists is necessary. The aim of the study was to investigate the knowledge and behavior of Dental practitioner regarding safety during oral radiographic procedures. It was a survey based study enrolled 100 dentists. A 13 semi - structured, validated questionnaire was used to assess knowledge and practice related to safety standards in Oral Radiology. The study revealed that amongst 100 Practitioners; 48 had > 5 years experiences, 86 had one machine, 61 had less than 5 years old machine. 97 used RVG, 73 took less than 10 radiographs per day, 65 didn't hold the X-Ray film with their fingers, 59 didn't use lead apron and 49 stood near patient without wearing lead apron, 54 did periodic check-up only when needed, 50 had to repeat, 54 didn't know safe distance, 75 didn't know the exposure time and 52 took radiographs of pregnant women. The study conclude that the periodic training and regular monitoring should be mandatory to improve knowledge about radiation and dose reduction techniques.

INTRODUCTION

In dentistry, radiographic evaluation is required in the majority of the cases before, during, and even after treatment. Hence, both dental practitioners and patients are at an increased risk of exposure to radiation if appropriate protective measures aren't taken.¹ Knowledge of dental health professionals about radiation safety - related guidelines is crucial. The International Commission on Radiological Protection (ICRP), the National Council on Radiation Protection and

Measurements (NCRP), and the Atomic Energy Regulatory Board (AERB) provide guidelines about permissible doses, equipment, and working protocols for occupational and nonoccupational personnel. The rules and regulations about radiation safety norms are to be followed by all the dental operatories in India.² The 'As Low As Reasonably Achievable (ALARA) principle should be followed to minimize these effects.¹ Radiographer's deals with ionising radiation daily, and as a consequence,

the doses they receive are exceeded. Several harmful outcomes are much higher than the ones of the possible depending on the general public. Thus, they have a higher radiation dose received.³ Thus, besides risk in developing an ionising radiation - the extensive benefits obtained from the induced complication. General public is diagnostic application of X-rays in both not aware of the consequences of medicine and dentistry, professional ionising radiation exposure, therefore negligence or ignorance can lead to medical personnel who are directly radiation hazards which are harmful to involved need to take appropriate the living tissues in a way that is measures for radiation protection to enough to cause cancer.⁴ Unnecessary protect themselves, the general public radiation exposure to the patients and and the environment. The effects of operators can be prevented by using ionising radiation exposure fall into two good equipment, proper technique, and categories, stochastic and deterministic. optimal processing methods. The methods of radiation protection are usually taught The chance of producing a stochastic during undergraduate training, and effect raises when the radiation dose dentists are expected to follow them increases. Stochastic effects do not have in private practice.¹ a dose threshold and characterise a In physicians, partial body radiation possible outcome in any dose such as such as the hands should not exceed cancer: the adverse effects could appear 500 mSv per year. Radiation to the after 10-20 years. In deterministic effects, thyroid gland and eyes should even stay the severity of the outcome increases below 300 and 150 mSv per year. For when the radiation dose is higher. Thus, example, a surgeon is exposed to a deterministic effects have a dose radiation dose of 0.05 mSv per minute at threshold, and with the threshold

a distance of approximately 0.5 m.⁵ In some countries, dentists are required to undergo certificate courses in radiation protection apart from their undergraduate training, and a permit is required for the use of radiographic equipment in the dental office. In India, however, no separate training is required. Therefore, the aim and objective of present study was to investigate the knowledge and practice regarding safety standards of oral radiology among dental practitioners during oral radiographic procedures in dental college, Bihar.¹

MATERIAL AND METHOD

The present ethics committee - approved study (Institutional Ethics Committee, IEC-BIDSH, Ref. No :- 121/BIDSH/IEC/2024-25(2023-24) was conducted among Dental Practitioners in Buddha Institute of Dental College and Hospital, Bihar. Under inclusion criteria , Dental Practitioner getting exposed while there was a radiography procedure, working in the Buddha Institute of

Dental College and Hospital, Bihar. One hundred practitioners were selected by simple random sampling using random number tables. The study protocol was explained to the participants, and the consent was obtained. A 13 item questionnaire in English was used to assess the participant's knowledge about the need for radiation protection and safety measures and adherence to radiation safety protocol in the dental radiology. The questionnaires were distributed to these 100 dental practitioners . The participation was voluntary, and confidentiality was assured. A few questions were objective in nature with "yes" or "no" options, whereas most of the questions had multiple choices. Questions to assess knowledge and practice were related to periodic check-ups of X-ray equipment, repetition of radiographs, the method of holding the film, use of a lead apron, the position and distance rule, adjustment of the exposure time, and performing

radiographs of pregnant women.¹ The data was obtained.

responses were analyzed, and quantitative

Knowledge and Practice regarding safety standards of Oral Radiology among Dental practitioners in Dental College, Bihar

NAME OF THE PRACTITIONER-

DEPARTMENT-

1. How many years have you been involved in private practice ?
A) 1 - 5 years B) 6 - 10 years C) 11 – 12 years D) > 15 years
2. How many intraoral X-ray machines are there in your clinic ?
A) 0 B) 1 C) 2 D) > 2
3. How old is your X-ray machine ?
A) 1 - 5 years B) 6 - 10 years C) > 10 years
4. Do you have periodic check – ups performed for your X-ray equipment ?
A) Yes B) no C) only when needed
5. Which type of radiographic receptor do you use ?
A) Conventional film B) digital sensor (RVG)
6. How many IOPA radiographs do you take in your dental clinic daily ?
A) < 10 B) 10 - 20 C) 20 - 30 D) > 30
7. How many times do you need to repeat the radiograph before you obtain a good image ?
A) Do not have to repeat B) once C) twice D) more than twice
8. Do you or your assistant hold the X-ray film with your fingers while taking periapical Radiographs ?
A) yes B) no
9. Do you use a lead apron for patient protection ?
A) yes, always B) yes, occasionally C) no
10. Where do you stand during intraoral exposure ?
A) behind a protective wall C) behind a lead screen
B) near patient without lead apron D) near the patient wearing a lead apron
11. If there is no barrier between you and the patient, in which area do you stand according to the X-ray tube and what is your distance from the patient ?
A) 135° to tube and 6 feet away C) 160° to tube and 8 feet away
B) 110° to tube and 3 feet away D) Do not know
12. How do you adjust the exposure time under the following conditions?

from overweight patient to thin patient

A) increase B) decrease C) no change

from anterior teeth to posterior teeth

A) increase B) decrease C) no change

From adult patient to pediatric patient

A) increase B) decrease C) no change

13. Do you take periapical radiographs of a pregnant women

A) yes B) no

RESPONSES OF STUDY POPULATION TO QUESTIONNAIRE

QUESTIONS	RESPONSE	NUMBER	%
1. How many years have you been involved in private Practice?	1-5 years	48	48.0
	6-10 years	32	32.0
	11-12 years	7	7.0
	>15 years	13	13
2. How many intraoral x-ray machines are there in your Clinic?	0	0	0
	1	86	86
	2	14	14
	>2	0	0
3. How old is your x-ray machine?	1-5 years	61	61
	6-10 years	35	35
	>10 years	4	4
4. Do you have periodic check-ups performed for your x-ray equipment?	Yes	12	12
	No	34	34

	Only when needed	54	54
5. Which types of radiographic receptor do you use?	Conventional film	3	3
	Digital sensor (RVG)	97	97
6. How many IOPA radiographs do you take in your Dental clinic daily?	<10	73	73
	10-20	23	23
	20-30	4	4
	>30	0	0
7. How many times do you need to repeat the radiograph Before you obtain a good image?	Do not have to repeat	50	50
	Once	45	45
	Twice	4	4
	More than twice	1	1
8. Do you or your assistant hold the x-ray film with your Fingers while taking periapical radiographs?	yes	33	33
	No	65	65
9. Do you use a lead apron for patient protection?	Yes, always	11	11
	Yes, occasionally	30	30
	No	59	59
10. Where do you stand during intraoral exposure?	Behind a protective wall	17	17
	Behind a lead screen	18	18
	Near patient without wearing lead apron	49	49
	Near the patient with wearing lead apron	16	16
11. If there is no barrier between you and the patient	135° to tube and 6 feet away	33	33
	110° to tube and 3 feet away	9	9
	160° to tube and 8 feet away	4	4
	Do not know	54	54
12. (A) How do you adjust the exposure time under the Following conditions? From overweight patient To thin patient	Increase	6	6
	Decrease	19	19
	No change	75	75
(B) How do you adjust the exposure time under the Following conditions? From anterior teeth to Posterior teeth	Increase	23	23
	Decrease	11	11
	No change	66	66
(C) How do you adjust the exposure time under the Following conditions? From adult patient to Paediatric patients	Increase	4	4
	Decrease	34	34
	No change	62	62
13. Do you take periapical radiographs of a pregnant Women	Yes	52	52
	No	48	48

RESULTS

Results were analysed statistically using the t-test for binary exposure variables and one - way analysis of variance (ANOVA) for exposure variables with > 2 categories. The knowledge score was normally distributed ; Mean scores and mean differences were estimated with participant characteristics as the independent categorical variables and the knowledge scale as a numeric dependent variable, using ANOVA and the t-test, wherever applicable. For analyses involving > 2 categories based on an overall p-value for group differences were estimated by default using ANOVA, which provided p - values for each group comparison (as compared to the first / reference group). Statistical significance was assumed as p - value < 0.05 (5 % Significance Level).

Demographic profile of the study subjects

Most practitioners were early in their private practice careers, with 48 % had 1–5 years of experience and 32 % with 6 –10 years. Only 7 % had practiced for 11–12 years, while 13 % have over 15 years of experience. Overall, 80 % had 10 or fewer years in private practice, indicating a predominantly early to mid career group.

Question	Response	Number	Percentage
1. How many years have you been involved in private practice ?	A) 1 - 5 Years	48	48.0
	B) 6 - 10 Years	32	32.0
	C) 11 - 12 Years	7	7.0
	D) > 15 Years	13	13.0

Radiographic equipment

The majority of clinics (86 %) had one intraoral X-ray machine, while 14 % have two. None of the respondents reported had zero or more than two machines,

indicating that most clinics are equipped with at least one, but not more than two, intraoral X-ray units.

Question	Response	Number	Percentage
2. How many intraoral X-ray machines are there in your clinic ?	A) 0	00	00
	B) 1	86	86
	C) 2	14	14
	D) >2	00	00

While, 61 % of respondents had X-ray machines that was 1–5 years old, 35 % reported machines aged 6–10 years, and only 4% had machines older than 10 years. This suggests that most clinics were equipped with relatively new X-ray equipment.

Question	Response	Number	Percentage
3. How old is your X-ray machine ?	A) 1 – 5 Years	61	61
	B) 6 – 10 Years	35	35
	C) > 10 Years	4	4

And only 12 % of respondents conduct periodic check - ups of their X-ray equipment, while 34 % did not perform regular maintenance. Notably, the majority 54 % only carried out check - ups when needed, indicating a reactive rather than preventive approach to equipment maintenance.

Question	Response	Number	Percentage
4. Do you have periodic check - ups performed for your X-ray equipment ?	A) Yes	12	12
	B) No	34	34
	C) Only when needed	54	54

Image receptors

A vast majority of respondents (97 %) used digital sensors (RVG) as their preferred radiographic receptor, while only a small fraction (3%) used conventional film. This indicates a strong shift towards digital radiography in clinical practice.

Question	Response	Number	Percentage
5. Which type of radiographic receptor do you use ?	A) Conventional film	3	3
	B) Digital sensor (RVG)	97	97

Response of participants

most dental practitioners (73%) took fewer than 10 IOPA radiographs daily. A smaller proportion (23%) took between 10–20 radiographs, and only 4% took 20–30 radiograph daily. Notably, none of the respondents reported taking more than 30 radiographs per day. This suggests that most clinics operate at a moderate diagnostic imaging workload.

Question	Response	Number	Percentage
6. How many IOPA radiographs do you take in your dental clinic daily?	A) <10	73	73
	B) 10-20	23	23
	C) 20-30	4	4
	D) >30	00	00

50 % of respondents had no need to repeat radiographs to obtain a good image, indicating efficient technique and equipment use. Meanwhile, 45 % required a single repeat, and only a small fraction 4 % and 1 % need two or more repetitions respectively. This suggests overall satisfactory radiographic accuracy and minimal retakes in most clinical settings.

Question	Response	Number	Percentage
7. How many times do you need to repeat the radiograph before you obtain a good image ?	A) Do not have to repeat	50	50
	B) Once	45	45
	C) Twice	4	4
	D) More than twice	1	1

65 % of respondents did not hold the X-ray film with their fingers while taking periapical radiographs, adhering to radiation safety protocols. However, 33 % still report holding the film manually, indicating a need for improved awareness and training on safe radiographic practices.

Question	Response	Number	Percentage
8. Do you or your assistant hold the X-ray film with your fingers while taking periapical Radiographs ?	A) YES	33	33
	B) NO	67	67

Methods of patient protection

Practitioner showed inconsistent use of lead aprons, with 11 % always used them, 30 % occasionally, and 59 % not used them at all, highlighting gaps in radiation safety practices.

Question	Response	Number	Percentage
9. Do you use a lead apron for patient protection?	A) Yes, always	11	11
	B) Yes, occasionally	30	30
	C) No	59	59

Methods of personal protection

Most individuals (49%) stood near the patient without a lead apron, risking radiation exposure. Only 18 % positioned themselves behind a lead screen, while 17 % stood behind a protective wall for safety. A smaller 16 % wore a lead apron near the patient, providing limited protection. These findings highlight inadequate shielding measures in intraoral exposure settings.

Question	Response	Number	Percentage
10. Where do you stand during intraoral exposure ?	A) Behind a protective wall	17	17
	B) Behind a lead screen	18	18
	C) Near patient without wearing lead apron	49	49
	D) Near the patient while wearing a lead apron	16	16

Only 33 % of individuals correctly selected option of standing 135° to the tube and 6 feet away. However, the most striking observation was that 54 % of individuals indicated that they did not know the correct response. This suggests a widespread gap in knowledge regarding radiation safety measures, emphasizing the urgent need for improved training and awareness to ensure proper protective positioning during X-ray procedures.

Question	Response	Number	Percentage
11. If there is no barrier between you and the patient, in which area do you stand according to the X-ray tube and what is your distance from the patient ?	A) 135° to tube and 6 feet away	33	33
	B) 110° to tube and 3 feet away	9	9
	C) 160° to tube and 8 feet away	4	4
	D) Do not know	54	54

Participants knowledge of radiation protection

75% of respondents recommend no change in exposure time when imaging thin

versus overweight patients, while 19 % suggest decreasing it and only 6 % advocate increasing it. This indicates a strong preference for consistent exposure settings, possibly due to automated systems or standardized protocols, with minor adjustments for thinner patients.

Question	Response	Number	Percentage
12 (A). How do you adjust the exposure time under the following conditions, From overweight patient to thin patient ?	A) Increase	6	6
	B) Decrease	19	19
	C) No change	75	75

While 66% of respondents recommend no change in exposure time when transitioning from imaging anterior teeth to posterior teeth. A smaller proportion (23%) suggest increasing exposure time, while only 11 % advise decreasing it. This indicates a general preference for maintaining consistent exposure settings, likely due to standardized imaging protocols or compensating techniques. The minority advocating for increased exposure may account for the greater density of posterior teeth and surrounding structures.

Question	Response	Number	Percentage
12(B) How do you adjust the exposure time under the following conditions , from anterior teeth to posterior teeth ?	A) Increase	23	23
	B) Decrease	11	11
	C) No change	66	66

62 % of respondents keep exposure time unchanged when imaging pediatric versus adult patients, while 34 % decrease it likely due to smaller patient size and lower radiation needs. Only 4 % increase exposure, suggesting it is rarely required for pediatric cases. Most favour standardized or adjusted protocols over higher exposure.

Question	Response	Number	Percentage
12 (C) How do you adjust the exposure time under the following conditions , from adult patient to pediatric patient ?	A) Increase	4	4
	B) Decrease	34	34
	C) No change	62	62

52 % of practitioners taking periapical radiographs for pregnant patients (likely using precautions like shielding) and 48 % avoiding it, reflecting caution over fetal radiation exposure. This split underscore varying clinical judgments about necessity and safety in prenatal dental care.

Question	Response	Number	Percentage
13. Do you take periapical radiographs of a pregnant women	A) YES	52	52
	B) NO	48	48

DISCUSSION

To be certain of the radiation safety of the patient and the operator, protocols, principles and guidelines had given to achieve radiation exposure dose for the patient As Low As Reasonably Possible. Routine check up of X-Ray machine is one of the basic steps of quality assurance to ensure optimal radiation exposure without any radiation leakage. In our study 12 % of the practitioner ensured regular maintenance check-ups of their X-ray machines ; which was lower than those obtained in studies by Sheikh et al. One of the main advantages of the Digital sensor is reduce exposure of X-ray upto 80 % as compare to conventional methods. In the present study 97 % of the practitioner were using digital imaging which was more than that reported by Jacobs et al. To avoid repetition high degree of sharpness and resolution is required. This can be obtained by using Proper equipment, Proper radiographic technique, Correct

processing and Interpretation of the image. A Lead apron is a protective piece of equipment that attenuates 98 % of scattered radiation. In this study 11 % of practitioner uses lead apron for patient protection and this percentage was lower than that reported by Math et al. The preferred method for operator protection during radiographic exposure was to position the receptor with a holding device and to leave room or to use a protective barrier or to stand at least 6 feet from the patient at an angle of 90-135 degree to the central X-Ray beam. In our study only 4 % of practitioner were aware of the position and distance rule, which can be used when no protective barrier is present and this percentage was lower than Lee et al. Selecting an appropriate exposure time is very important to minimize the patient's radiation exposure and obtain an image with proper density and contrast. Pediatric patients require 25-50 % less exposure than do adults. Underweight

patients require less exposure than do overweight patients. Edentulous patient require 25 % less exposure than do dentulous patients. In this study 68 % of dentists didn't changed the exposure time according to patient's age, build and the region of the oral cavity being radiographed. Even radiographs can be taken during pregnancy if the patient wear lead apron to avoid greatest risk to the foetus for abnormality. In our study 48 % of practitioner thought that radiographs cannot be taken in pregnant women. Although the Knowledge of Dentists about Radiation Protection Techniques in our study was fair and acceptable but they should implement proper radiation protection techniques and guidelines concerning patient and personnel safety.

CONCLUSION

The implementation of recommended radiation safety

protocols and practices in the dentistry is vital for the safety of the working personnel, patients, and environment. By following the radiation safety guidelines, dental professionals can protect themselves and others from the hazards of radiation. The present study revealed the importance of awareness and adherence to radiation safety measures among dental practitioners and the need for more studies to be conducted in this domain. This study demonstrated that most of the participants revealed a reasonable knowledge of the need for personnel protection, safety measures, and adherence to radiation safety protocol. The periodic training and regular monitoring of occupationally exposed dental practitioners should be mandatory to ensure appropriate compliance with radiation safety

regulations.

ACKNOWLEDGEMENTS - Nil.

FINANCIAL SUPPORT AND

SPONSORSHIP - This manuscript did not receive any funding from public or private sectors.

CONFLICTS OF INTEREST - The authors have no conflict of interests to disclose.

REFERENCES-

1. Vikrant O. Kasat et al, 2016. Knowledge and practice regarding safety standards of oral radiology among dental practitioners in western Maharashtra, India. Oral Radiol.
2. Aarati S. Panchbhai , Prasanna R. Sonar. 2024. Assessing the Adherence to Safety Protocol Among Personnel Working in the Dental Radiology Department : A Qualitative Cross-Sectional Study. Cureus part of springer nature.
3. C. Zervides, L. Sassis, P et al. 2019. Assessing radiation protection knowledge in diagnostic radiography in the Republic of Cyprus. A questionnaire survey. Elsevier (2019) 1078 – 8174.
4. Monica Chaudhry et al. 2015. Oral Radiology Safety Standards Adopted by the General Dentists Practicing in National Capital Region (NCR) Arch Orthop Trauma Surg(2015).
5. Thorsten Jentzsc et al. 2015. The compliance with and knowledge about radiation protection in operating room personnel: a cross-sectional study with a questionnaire. Arch Orthop Trauma Surg (2015) 135:1233-1240
6. Aps J K. 2010. Flemish general dental practitioners' knowledge of dental radiology. Dentomaxillofac Radiol ; 39 : 113 – 8.
7. Binna A, Rajesh G, Denny C, Ahmed J, Nayak V. 2013. Insights into the state of radiation protection among a subpopulation of Indian dental practitioners. Imaging Sci Dent.; 43 : 253 – 9.
8. Bohay R N, Kogon S L, Stephens R G. 1994. A survey of radiographic techniques and equipment used by a sample of general dental practitioners. Oral Surg Oral Med Oral Pathol ; 78 : 806 – 10
9. Farman A G, Farman T T. 2005. A

- comparison of 18 different x-ray detectors currently used in dentistry. Oral Surg Oral Med Oral Pathol Oral Radiol Endod ; 99 (4) : 485–9.
10. Geist J R, Katz J O. 2002. Radiation dose - reduction techniques in North American dental schools. Oral Surg Oral Med Oral Pathol Oral Radiol Endod; 93(4):496–505.
 11. Haden N K, Andrieu S C, Chadwick D G, Chmar J E, Cole J R, George M C et al. 2006. The dental education environment. J Dent Educ; 70: 1265-70.
 12. Handal B, Groenlund C, Gerzina T. 2010. Dentistry students' perceptions of learning management systems. Eur J Dent Educ ; 14: 50-4.
 13. Hendricson W D, Andrieu S C, Chadwick D G, Chmar J E, Cole J R, George M C et al. 2006. Educational strategies associated with development of problem-solving, critical thinking, and self-directed learning. J Dent Educ; 70: 925-36.
 14. Ilgüy D, Ilgüy M, Dincer S, Bayırlı G. 2005. Survey of dental radiological practice in Turkey. — Dentomaxillofac Radiol ; 34 :222 – 7.
 15. Jacobs R, Vanderstappen M, Bogaerts R, Gijbels F. 2004. Attitude of the Belgian dentist population towards radiation protection. Dentomaxillofac Radiol ; 33 : 334 – 9.
 16. Kuba R, Kohli A. 2014. Continuing professional development of dentists through distant learning: An Indira Gandhi National Open University-Dental Council of India experiment a report. Indian J Dent Res ; 25 : 821 – 7.
 17. Kantor M L. 2005. Dental digital radiography: more than a fad, less than a revolution. J Am Dent Assoc; 136 (10):1358, 1360, 1362.
 18. Lee B D, Ludlow J B. 2013. Attitude of the Korean dentists towards radiation safety and selection criteria. Imaging Sci Dent ; 43: 179 – 84.
 19. Linjawi A I, Walmsley A D, Hill K B. 2012. Online discussion boards in dental education : potential and challenges. Eur J Dent Educ.
 20. Ludlow J B, Mol A. 2004. Digital imaging. In: White S C, Pharoah M J, editors. Oral radiology. Principles

- and interpretation. 5th ed. Mosby: St. Louis; p. 225–44.
21. Math S Y, Murugeshappa D G, Annigeri R, Kalra D. 2013 . Compliance of Indian dentists with oral radiology safety measures. *J Oral Maxillofac Radiol* ; 1:104 – 10.
 22. Melo M F, Melo S L. 2008. Radioprotection in dentistry offices. *Cienc Saude Colet* ; 13 (Suppl 2) : 2163 - 70.
 23. Miles D A, Langlais R P, 2004, National Council on Radiation Protection and Measurement. NCRP Report No. 145 : New dental X-ray guidelines : their potential impact on your dental practice. *Dent Today* ; 23(9):128, 130, 132, 134 passim ; quiz 134. L
 24. Neves F S, Vasconcelos T V, Bastos L C, Góes L A, Freitas D Q. 2010. Attitudes of dentists in relation to radiological protection, according to Brazilian law. *Rev Odontol Bras Central* ; 19: 301-5.
 25. Olapiriyakul K, Scher J M. 2006. A guide to establishing hybrid learning courses: employing information technology to create a new learning experience, and a — case study. *Internet Higher Educ* ; 9 : 287-301.
 26. Ruprecht A. 2009. The status of oral and maxillofacial radiology worldwide in 2007. *Dentomaxillofac Radiol* ; 38 : 98-103.
 27. Salti L, Whaites E J. 2002. Survey of dental radiographic services in private dental clinics in Damascus, Syria. *Dentomaxillofac Radiol* ; 31 : 100 – 5.
 28. Shahab S, Kavosi A, Nazarinia H, Mehralizadeh S, Mohammadpour M, Emami M. 2012. Compliance of Iranian dentists with safety standards of oral radiology. *Dentomaxillofac Radiol* ; 41 : 159 – 64.
 29. Sheikh S, Pallagatti S, Singla I, Gupta R, Aggarwal A, Singh R, et al. 2014. Survey of dental radiographical practice in States of Punjab and Haryana in India. *J Investig Clin Dent* ; 5:72–7.
 30. Svenson B, Söderfeldt B, Gröndahl H G. 1996. Attitudes of Swedish dentists to the choice of dental X-ray film and collimator for oral radiology. *Dentomaxillofac Radiol* ; 25: 157-61.
 31. Svenson B, Söderfeldt B,

- Gröndahl H G. 1996. Analysis of dentists' attitudes towards risks in oral radiology. *Dentomaxillofac Radiol* ; 25: 151-6.
32. Tosoni G M, Campos D M, Silva M R. 2003. Frequency and quality management of intraoral radiographic examinations in private dental practices. *Rev Odontol UNESP* ; 32: 25 - 9.
33. van der Stelt P F. 2005. Filmless imaging: the uses of digital radiography in dental practice. *J Am Dent Assoc* ; 136 (10) : 1379–87.
34. White S C, Pharoah M J. 2004. Health physics. Oral radiology: principles and interpretation. 5th ed. New Delhi : Elsevier; p. 47–68.