(ISSN: 0973-7049(P))

The Bioscan AN INTERNATIONAL QUARTERLY JOURNAL OF LIFE SCIENCES 21(4): S.I (4),1009-1017, 2026 www.thebioscan.com

Microbial-Based Innovations for the Prevention of Mosquito-Borne Diseases: Mechanisms, Applications, Challenges and Future Perspectives

Dr. Kiran Bala Bhuyan[1], Dr. Lipika Jena[2], Dr. Radha Yadamuri[3], Dr. Manzur Hassan[4]

[1]Professor in Zoology, S.K.C.G. Autonomous College,Paralakhemundi , Gajapati ,Odisha -761200, kiranbpkd@gmail.com

[2]Associate Professor, Department of microbiology,IMS & SUM hospital, Bhubaneswar, Odisha, lipikajena.jena07@gmail.com

[3]Assistant Professor Department of Biological and Chemical sciences Mohan Babu University, radhaprasadburla31@gmail.com

[4]PhD (Cotton University, Panbazar, Ghy-1) hassanmanzur0@gmail.com

DOI: https://doi.org/10.63001/tbs.2026.v21.i04.S.I(4).pp1009-1017

KEYWORDS

mosquito-borne diseases; microbial control; Wolbachia; Bacillus thuringiensis israelensis; Lysinibacillus sphaericus; entomopathogenic fungi; mosquito microbiome; biological control; dengue; vector management

Article Info

Received:02.20.2026

Accepted: 29.10.2026

Published: 11.11.2026

Abstract

Mosquito-borne diseases remain a major global public-health challenge, contributing substantially to morbidity, mortality, healthcare expenditure, and socioeconomic disruption. Dengue, malaria, chikungunya, Zika, yellow fever, West Nile fever, and lymphatic filariasis are transmitted by mosquito vectors whose geographical distribution and population dynamics are influenced by environmental change, urbanization, climate variability, human mobility, and vector adaptation. Conventional mosquito-control approaches, particularly chemical insecticides, remain important but are increasingly challenged by insecticide resistance, operational limitations, environmental concerns, and difficulties in achieving sustained population-level control. Microbial-based technologies have consequently emerged as an important component of innovative and integrated vector management. These technologies include bacterial larvicides such as Bacillus thuringiensis subsp. israelensis (Bti) and Lysinibacillus sphaericus, entomopathogenic fungi including Beauveria bassiana and Metarhizium spp., mosquito-associated microbial manipulation, and Wolbachia-based population replacement or suppression. Microbial agents can act through several mechanisms, including larval gut toxicity, fungal infection through the insect cuticle, reproductive manipulation, pathogen interference, and modification of mosquito-associated microbial communities. Evidence from randomized and quasi-experimental studies has demonstrated substantial reductions in dengue incidence following deployment of Wolbachia-infected Aedes aegypti in several settings, although effectiveness varies according to ecological, operational, and epidemiological conditions. Recent research is also expanding toward mosquito microbiome engineering, insect-specific viruses, improved microbial formulations, synergistic combinations, and precision delivery systems. This review examines the scientific basis, current evidence, applications, advantages, limitations, implementation considerations, and future prospects of microbial approaches for mosquito-borne disease prevention.

1. Introduction

Mosquitoes are among the most important biological vectors of infectious diseases affecting humans. Members of the genera Aedes, Anopheles, and Culex transmit a wide range of viral, parasitic, and filarial pathogens. Dengue, malaria, chikungunya, Zika, yellow fever, Japanese encephalitis, West Nile fever, and lymphatic filariasis collectively impose a

substantial burden on healthcare systems and communities.The epidemiology of mosquito-borne diseases is changing as a consequence of urban expansion, climate variability, environmental transformation, international travel, population movement, and changes in mosquito ecology. At the same time, intensive use of conventional insecticides has contributed to the emergence and spread of insecticide resistance. These developments have stimulated the search for complementary vector-control technologies that are effective, environmentally compatible, scalable, and capable of functioning within integrated vector-management programs.

Microbial control represents one such approach. Unlike conventional insecticides that generally act through chemical toxicity, microbial interventions use microorganisms, microbial toxins, symbiotic microorganisms, or microbial-derived biological mechanisms to interfere with mosquito survival, development, reproduction, or pathogen transmission.

2. Objectives

The principal objectives of this review are:
1. To summarize major microbial approaches used or investigated for mosquito control.
2. To describe the biological mechanisms underlying microbial-mediated mosquito suppression or pathogen blocking.
3. To evaluate recent evidence regarding the effectiveness of microbial interventions.
4. To compare bacterial, fungal, symbiont-based, viral, and microbiome-oriented approaches.
5. To identify operational, ecological, regulatory, and community-level challenges.
6. To examine opportunities for combining microbial technologies with integrated vector management.
7. To identify emerging research directions for next-generation microbial mosquito-control strategies.

3. Literature Review

Conventional mosquito control includes environmental management, larval-source management, insecticide-treated nets, indoor residual spraying, space spraying, adulticides, and chemical larvicides. These interventions remain important components of disease-control programs, but their performance can be compromised by insecticide resistance, limited coverage, environmental conditions, and operational constraints.

The development of microbial technologies has been driven partly by the need to diversify vector-control portfolios. Microbial interventions may provide different modes of action from conventional chemical insecticides and can therefore contribute to resistance-management strategies.

4. Methodology

An integrative literature-review methodology was used to synthesize evidence concerning microbial interventions for mosquito-borne disease prevention. The review focused on biological mechanisms, efficacy, field implementation, environmental considerations, and emerging technologies.

Information sources included PubMed/MEDLINE, Cochrane Library, WHO publications, peer-reviewed scientific journals, public-health and vector-control agencies, and reference lists from recent reviews.

Representative search terms included “microbial mosquito control,” “Wolbachia Aedes dengue,” “Bacillus thuringiensis israelensis mosquito,” “Lysinibacillus sphaericus mosquito,” “entomopathogenic fungi mosquito control,” “Beauveria bassiana mosquito,” “Metarhizium mosquito,” “mosquito microbiome vector competence,” and “insect-specific virus mosquito control.”

Studies were included when they investigated microbial organisms or microbial-derived

mechanisms for mosquito control, examined mosquito survival, development, reproduction, abundance, or vector competence, or evaluated interventions in laboratory, semi-field, field, epidemiological, or implementation settings. Studies unrelated to mosquito control, duplicates, and publications without sufficient scientific information were excluded.

Evidence was synthesized thematically rather than through a quantitative pooled analysis performed specifically for this review.

5. Major Microbial-Based Approaches

5.1 Bacterial Larvicides

Bacterial larvicides are among the most mature microbial technologies in mosquito control.

5.1.1 Bacillus thuringiensis subsp. israelensis

Bti is a naturally occurring bacterium that produces insecticidal proteins, including Cry and Cyt toxins. Following ingestion by mosquito larvae, the toxins become activated within the larval midgut and damage intestinal epithelial cells. Bti is particularly valuable because its activity is relatively selective toward susceptible dipteran larvae.

5.1.2 Lysinibacillus sphaericus

Lysinibacillus sphaericus, historically known as Bacillus sphaericus, produces a binary toxin with activity against mosquito larvae. Its environmental persistence can exceed that of some Bti formulations. Combining L. sphaericus with Bti or other control methods can provide complementary mechanisms.

5.2 Entomopathogenic Fungi

Entomopathogenic fungi represent another important microbial-control platform. Important genera include Beauveria, Metarhizium, Isaria, and Lecanicillium. Fungal agents can infect mosquitoes through direct contact with the cuticle. Spores attach to the mosquito, germinate, penetrate the cuticle, proliferate internally, and ultimately cause physiological disruption and death.

5.2.1 Beauveria bassiana

Beauveria bassiana has attracted substantial research interest because it can affect mosquito larvae and adults. Field-oriented technologies have also investigated fungal autodissemination and attractive stations.

5.2.2 Metarhizium spp.

Metarhizium fungi can infect mosquitoes through cuticular penetration and have been investigated as alternatives to chemical insecticides. Strain selection, host specificity, formulation, environmental tolerance, and delivery mechanisms are important determinants of field performance.

6. Wolbachia-Based Mosquito Control

Wolbachia are intracellular bacteria naturally associated with many arthropod species. They are maternally transmitted and can influence host reproduction and interactions with pathogens.

Scientists have introduced specific Wolbachia strains into Aedes aegypti to create mosquito populations with reduced capacity to support replication of dengue and other arboviruses.

Two major strategies have emerged: population replacement and population suppression.

In population replacement, Wolbachia-infected male and female mosquitoes are released with the objective of establishing the infection within the local mosquito population. In population suppression, Wolbachia-infected males are released so that incompatible matings reduce reproductive output.

The landmark AWED randomized trial in Yogyakarta, Indonesia, demonstrated reduced virologically confirmed dengue following deployment of wMel-Wolbachia-infected Aedes aegypti. Subsequent studies in Brazil, Colombia, and other settings have expanded the evidence base.

7. Evidence From Field Studies

Field evidence is increasingly important because laboratory pathogen-blocking effects do not necessarily predict epidemiological impact.

Evidence from randomized, quasi-experimental, and implementation studies indicates that Wolbachia deployments can reduce dengue outcomes in several settings. However, the magnitude of effects varies according to local mosquito ecology, intervention coverage, baseline transmission, environmental conditions, and implementation characteristics.

Microbial larvicides and entomopathogenic fungi provide complementary mechanisms targeting mosquito survival, while Wolbachia primarily targets vector competence and/or mosquito reproduction.

8. Microbiome-Based Vector Control

Mosquitoes are holobionts whose biology is influenced by complex communities of bacteria, fungi, viruses, and other microorganisms. The mosquito microbiome can affect larval development, nutrition, immunity, reproduction, insecticide resistance, pathogen replication, and vector competence.

Consequently, microbiome manipulation represents a potential next generation of microbial vector control. Research is increasingly examining whether particular microorganisms can be introduced, enriched, depleted, or engineered to reduce pathogen transmission. However, microbiome engineering remains less mature than Bti or Wolbachia deployment.

9. Insect-Specific Viruses

Insect-specific viruses generally replicate within mosquitoes but do not cause conventional disease in humans. Some studies suggest that interactions between insect-specific viruses and arboviruses may alter vector competence. This is an emerging area requiring additional field and epidemiological evidence.

10. Mechanisms of Microbial Control

Microbial technologies can influence mosquito populations through several mechanisms:

- Direct larval toxicity: Bti and L. sphaericus toxins damage the larval midgut after ingestion.
- Fungal infection: Beauveria and Metarhizium attach to and penetrate the mosquito cuticle.
- Vector-competence reduction: Wolbachia can reduce the ability of Aedes mosquitoes to support arbovirus replication.
- Reproductive manipulation: Wolbachia-associated cytoplasmic incompatibility can alter mosquito reproduction.
- Microbiome-mediated effects: candidate microorganisms can alter host physiology and pathogen replication.

11. Comparative Characteristics of Major Microbial Strategies

Bti primarily targets mosquito larvae through toxin-mediated midgut damage and generally requires repeated application. L. sphaericus also targets larvae and has useful persistence but resistance considerations. Entomopathogenic fungi can target larvae and adults but are relatively dependent on environmental conditions. Wolbachia targets adult Aedes populations and may persist after successful population establishment, although implementation requires substantial infrastructure and community engagement.

12. Synergistic and Combination Approaches

Potential combinations include Bti with L. sphaericus, microbial larvicides with environmental source reduction, fungal agents with autodissemination systems, Wolbachia with conventional surveillance, and microbial agents with insecticide resistance-management programs. Combination strategies may diversify mechanisms and reduce dependence on a single technology.

13. Novel Formulation and Delivery Technologies

Important delivery approaches include long-lasting microbial formulations, autodissemination systems, attractive toxic stations, and precision spatial deployment using geographic information systems, mosquito surveillance, remote sensing, and mathematical modeling.

14. Role of Climate and Environmental Factors

Temperature, humidity, rainfall, water chemistry, ultraviolet radiation, breeding-site characteristics, mosquito density, mosquito genetics, and microbial strain can influence microbial efficacy. Environmental adaptation is therefore an important consideration when planning deployment.

15. Safety and Ecological Considerations

Important environmental questions include whether the intervention affects non-target organisms, whether microbial agents persist outside intended hosts, whether horizontal transmission can occur, and whether mosquitoes or pathogens can adapt. Each microbial technology and formulation should be evaluated within its specific ecological context.

16. Community Engagement and Social Acceptance

Successful implementation requires transparent communication, public consultation, accessible safety information, engagement with healthcare professionals and local authorities, and mechanisms for monitoring and reporting. Community engagement can influence operational coverage, participation, trust, and long-term sustainability.

17. Economic and Health-System Considerations

The economic value of microbial control depends on production costs, delivery infrastructure, frequency of intervention, surveillance requirements, healthcare costs avoided, disease incidence, population density, and local mosquito ecology. Economic evaluations increasingly examine both program costs and potential healthcare savings.

18. Resistance and Evolutionary Considerations

Microbial interventions can create evolutionary pressure. Bacterial larvicides may select for resistance, particularly under repeated exposure to a single toxin. For Wolbachia, long-term monitoring should consider changes in bacterial density, infection stability, mosquito adaptation, pathogen adaptation, and environmental effects.

19. Current Evidence and Knowledge Gaps

Major gaps include variable Wolbachia performance across ecological settings, persistence and resistance issues for bacterial larvicides, field scalability of fungal agents, limited field evidence for microbiome engineering and insect-specific viruses, limited standardized trials of combinations, differences among economic models, insufficient longitudinal ecological data, context-specific community acceptance, and uncertainty about climate resilience.

20. Discussion

The development of microbial mosquito-control technologies represents a shift from conventional chemical suppression toward biologically mediated vector management. Bti and related bacterial larvicides provide targeted larval control. Entomopathogenic fungi expand the intervention space by targeting mosquitoes through contact. Wolbachia represents a different model by modifying vector competence and reproduction.

The strongest epidemiological evidence currently concerns Wolbachia, but laboratory results do not automatically translate into equivalent public-health effects across all locations. Future evaluation should increasingly focus on epidemiological endpoints, long-term durability, cost-effectiveness, environmental safety, and implementation quality.

21. Proposed Integrated Microbial Vector-Control Framework

A practical framework consists of seven stages:

1. Surveillance: mosquito density, pathogen surveillance, insecticide resistance, and environmental monitoring.
2. Risk characterization: identify vector species, breeding habitats, and disease hotspots.
3. Microbial selection: Bti, L. sphaericus, fungal agents, Wolbachia, or emerging microbiome interventions.
4. Targeted implementation: larval control, population replacement/suppression, and autodissemination.
5. Community engagement: education, consultation, participation, and reporting.
6. Monitoring: microbial prevalence, mosquito density, vector competence, and disease incidence.
7. Adaptive management: modify intervention according to surveillance and ecological evidence.

22. Future Perspectives

Future research should address next-generation Wolbachia strains, microbiome engineering, engineered microorganisms under rigorous biosafety oversight, artificial intelligence for mosquito prediction and deployment planning, climate-responsive microbial control, and multi-technology integration.

The future is likely to involve microbial control combined with environmental management, surveillance, chemical resistance management, community participation, and digital technologies.

23. Limitations of the Review

This review was designed as an integrative narrative review rather than a registered systematic review with formal meta-analysis. Substantial heterogeneity exists among microbial interventions, mosquito species, geographical regions, deployment strategies, and outcome measures. Evidence is considerably stronger for some technologies, particularly Wolbachia and Bti, than for emerging microbiome and insect-specific-virus approaches.

24. Conclusion

Microbial-based technologies are expanding the scientific and operational toolbox for preventing mosquito-borne diseases. Bacterial larvicides such as Bti and Lysinibacillus sphaericus provide established methods for targeted larval control, while entomopathogenic fungi offer alternative mechanisms capable of affecting mosquitoes at different life stages.

Among emerging technologies, Wolbachia has generated substantial epidemiological evidence for reducing dengue transmission through population replacement and suppression approaches. The magnitude and durability of effects remain context-dependent.

Future research should prioritize multi-country field studies, long-term epidemiological evaluation, resistance monitoring, cost-effectiveness assessment, environmental safety, community acceptance, and climate resilience. An integrated vector-management framework combining microbial technologies with environmental and epidemiological surveillance may provide an adaptable approach to reducing mosquito-borne disease transmission.

Table 1. Selected Evidence for Microbial-Based Mosquito Control

Table 2. Comparative Characteristics of Major Microbial Strategies

Table 3. Major Knowledge Gaps and Research Priorities

Figure 1. Conceptual Framework of Microbial Mosquito Control




Figure 1. Conceptual framework illustrating major microbial pathways for mosquito control.

Figure 2. Integrated Microbial Vector Management Framework




Figure 2. Integrated framework for planning, implementing, monitoring, and adapting microbial vector-control interventions.

References

1. Utarini A, Indriani C, Ahmad RA, et al. Efficacy of Wolbachia-infected mosquito deployments for the control of dengue. New England Journal of Medicine. 2021;384:2177–2186. doi:10.1056/NEJMoa2030243.

2. Indriani C, Tantowijoyo W, Rancès E, et al. Reduced dengue incidence following deployments of Wolbachia-infected Aedes aegypti in Yogyakarta, Indonesia. Gates Open Research. 2020;4:50.

3. Pinto SB, et al. Effectiveness of Wolbachia-infected mosquito deployments in reducing dengue and other Aedes-borne diseases in Niterói, Brazil. PLoS Neglected Tropical Diseases. 2021.

4. Fox T, Sguassero Y, Chaplin M, et al. Wolbachia-carrying Aedes mosquitoes for preventing dengue infection. Cochrane Database of Systematic Reviews. 2024;4:CD015636.

5. Velez ID, Tanamas SK, Arbelaez MP, et al. Reduced dengue incidence following city-wide wMel Wolbachia mosquito releases throughout three Colombian cities. PLoS Neglected Tropical Diseases. 2023;17:e0011713.

6. Lim JT, et al. Efficacy of Wolbachia-mediated sterility to reduce the incidence of dengue: a synthetic control study in Singapore. The Lancet Microbe. 2024.

7. O'Neill SL, Ryan PA, Simmons CP, Moreira LA. Long-term durability and public health impact of city-wide wMel Wolbachia mosquito releases in Niterói, Brazil. 2025.

8. Impact of Wolbachia-infected Aedes aegypti on dengue incidence in endemic regions: a systematic review and meta-analysis. 2026.

9. Nichols HL, Coon KL. Leveraging microbial ecology for mosquito-borne disease control. Trends in Parasitology. 2025;41:670–684.

10. Liu J, Zhang L, Tong H, et al. Microorganisms in mosquitoes for controlling mosquito-borne viral diseases: from lab to field. Trends in Microbiology. 2026;34:277–292.

11. Hong G, Yu L, Ji H, et al. Microbiological control for mosquito larvae: current progress and applications. Virulence. 2025;16:2569999.

12. Deng SQ, Li N, Yang XK, et al. Recombinant Beauveria bassiana expressing Bacillus thuringiensis toxin Cyt1Aa for Aedes mosquito control. Microbiology Spectrum. 2024;12:e03792-23.

13. Rodrigues de Miranda RP, Soares TKDA, Castro DP, Genta FA. General aspects, host interaction, and application of Metarhizium sp. in arthropod pest and vector control. Frontiers in Fungal Biology. 2024;5:1456964.

14. Patil Tawidian, Kang Q, Michel K. The potential of a new Beauveria bassiana isolate for mosquito larval control. Journal of Medical Entomology. 2023;60:131–147.

15. Hussain S, Kanuka H, Rakotondrafara A, et al. Pathogenicity and sub-lethal activity of entomopathogenic fungi against Aedes aegypti and Anopheles stephensi. Journal of Invertebrate Pathology. 2024;207:108233.

16. Recent advancements in pathogenic mechanisms, applications and strategies for entomopathogenic fungi in mosquito biocontrol. 2023.

17. Delivery and effectiveness of entomopathogenic fungi for mosquito and tick control: current knowledge and research challenges. 2022.

18. Discovery of novel entomopathogenic fungi for mosquito-borne disease control. 2023.

19. Evaluation of In2Care mosquito stations for suppression of Aedes notoscriptus. Journal of Medical Entomology. 2023.

20. Bacterial toxins active against mosquitoes: mode of action and resistance. 2021.

21. Flores HA, Taneja de Bruyne J, O'Donnell TB, et al. Multiple Wolbachia strains provide comparative levels of protection against dengue virus infection in Aedes aegypti. PLoS Pathogens. 2020;16:e1008433.

22. Fraser JE, O'Donnell TB, Duyvestyn JM, et al. Novel phenotype of Wolbachia strain wPip in Aedes aegypti. PLoS Pathogens. 2020;16:e1008410.

23. Koh C, Islam MN, Ye YH, et al. Dengue virus dominates lipid metabolism modulations in Wolbachia-coinfected Aedes aegypti. Communications Biology. 2020;3:518.

24. Mancini MV, Ant TH, Herd CS, et al. High temperature cycles result in maternal transmission and dengue infection differences between Wolbachia strains in Aedes aegypti. mBio. 2021;12:e00250-21.

25. Edenborough KM, Flores HA, Simmons CP, Fraser JE. Using Wolbachia to eliminate dengue: will the virus fight back? Journal of Virology. 2021;95:e02203-20.

26. Zhang H, Lui R. Releasing Wolbachia-infected Aedes aegypti to prevent the spread of dengue virus: a mathematical study. Infectious Disease Modelling. 2020;5:142–160.

27. Cardona-Salgado D, Campo-Duarte DE, Sepulveda-Salcedo LS, et al. Optimal release programs for dengue prevention using Aedes aegypti mosquitoes transinfected with Wolbachia strains. Mathematical Biosciences and Engineering. 2021;18:2952–2990.

28. Hugo LE, Rašić G, Maynard AJ, et al. Wolbachia wAlbB inhibit dengue and Zika infection in Aedes aegypti. PLoS Neglected Tropical Diseases. 2022;16:e0010786.

29. Wolbachia-based emerging strategies for control of vector-transmitted disease. 2024.

30. Exploiting Wolbachia as a tool for mosquito-borne disease control: pursuing efficacy, safety, and sustainability. 2025.

31. Impact of symbiotic insect-specific viruses on mosquito vector competence for arboviruses. 2024.

32. Mosquito gut microbiota: a review. 2024.

33. Fungal microbiota diversity in Aedes, Anopheles and Culex and their potential use as biological tools to control vector-borne diseases: a systematic review. 2026.

34. Wolbachia-based mosquito control: environmental perspectives on population suppression and replacement strategies. 2025.

35. Wolbachia as a transformative tool for mosquito-borne disease control: a comprehensive review of mechanisms, efficacy, and future directions. 2025.

36. Wolbachia-based biocontrol of Aedes aegypti: current progress, challenges, and future prospects. 2025.

37. Safaei S, et al. Wolbachia: a bacterial weapon against dengue fever. 2025.

38. World Health Organization. Operational manual on larval source management: control of Anopheles and Aedes mosquito vectors. Geneva: WHO; 2026.

39. World Health Organization. WHO update on the assessment of Wolbachia population replacement for the control of Aedes-borne diseases. 2026.

40. Centers for Disease Control and Prevention. Mosquitoes with Wolbachia. 2024.