

EXPLORING THE MEDICINAL POTENTIAL OF GUAVA (*Psidium guajava*) FOR HEALTH AND WELLNESS

SOWJANYA MGS¹, MURALI B² AND RAMESH B^{1*}

¹Department of Genetics and Genomics, Yogi Vemana University, Kadapa Andhra Pradesh - 516005, INDIA

²Department of Computer Science and Engineering, YSR Engineering College of YVU, Proddatur

e-mail: adenoramesh@gmail.com

ORCID ID: <https://orcid.org/0000-0001-9814-881X>

KEYWORDS

Medicinal properties
Psidium guajava
antioxidant
anti-cancer

Received on :

22.11.2023

Accepted on :

07.02.2024

*Corresponding
author

ABSTRACT

The global escalation of chronic degenerative diseases, often associated with foodborne illnesses, has led to heightened healthcare costs and a diminished quality of life. Consuming foods rich in bioactive substances, such as fruits, vegetables, and seeds, can help reduce the risk factors for developing certain diseases. *Psidium guajava*, well known for its exquisite flavor, has also been the subject of much research due to its possible health advantages. These benefits include anti-inflammatory, anti-cancer, anti-diabetic, antimicrobial, anti-diarrheal, and anti-hypertensive properties. However, using guava as a therapeutic plant has drawbacks and restrictions. This review sheds light on the relevance of guava in promoting health and wellness by emphasizing the necessity of comprehending the medicinal value of guava and its possible uses. Future investigations into the medicinal potential of guava may result in cutting-edge medical tactics.

INTRODUCTION

Navigating a healthy lifestyle in today's modern world is riddled with challenges, primarily due to the presence of harmful chemicals in our food. The interconnected global dynamics of population expansion, technological progress, and the swift industrialization of society compound this issue. Beyond fostering a surge in chronic illnesses, these factors also contribute to escalating healthcare expenses and a general decline in people's overall quality of life. These factors also encourage the growth of chronic degenerative diseases such as metabolic syndrome, diabetes mellitus, dyslipidemia, and cardiac conditions (Sharma and Chandola, 2011; Remington and Brownson, 2011). Fruits, vegetables, and seeds should all be a part of our diets in order to reduce the risk factors associated with these diseases. These foods contain bioactive compounds that can help treat, prevent, and manage various health issues (Barbalho *et al.*, 2011; Párraga *et al.*, 2011). By consciously choosing these nutrient-rich foods, we can improve our overall well-being and reduce the impact of the challenges posed by the current scenario. The tropical fruit *Psidium guajava* (*P. guajava*) (Fig. 1), which has a tart and sweet flavour, is a favourite. Due to its numerous therapeutic benefits in addition to its deliciousness, this fruit has a long history of use in conventional medicine. The comprehensive studies on various parts of the *Psidium guajava tree*, including the leaves, bark, roots, and fruits, have revealed their therapeutic applications for various diseases (Table 1). The abundant flavonoids, carotenoids, tannins, and phenolic chemicals in the plant have been shown in numerous studies to have anti-

inflammatory, anti-diabetic, anti-microbial, anti-diarrheal, anti-hypertensive, anti-cancer, and anti-oxidant properties (Carrasco-Castilla *et al.*, 2012). Guava is a fruit that originated in the tropics of the Americas, but it has since been widely farmed and dispersed around the world, including South Africa and India (Lima *et al.*, 2018). The guava tree, a small tree that belongs to the Myrtaceae family, can thrive in regions with fertile soil and a tropical or subtropical climate. FAOSTAT (Manosroi *et al.*, 2006) reports that China and India currently produce the most guava. Depending on the species, guava fruits can be round or oval and range in size from 4 to 12 centimeters (1.6 to 4.7 in). The fruit initially has a faint green tint, then ripens into a yellow color. The tree's leaves are coated with a deep green hue and have an acute point. They range in shape from elliptical to oval (Kumar *et al.*, 2021). Apple guava is the guava cultivar that is most frequently bought (Chen and Yen 2007). There is evidence that several plant parts of guava, including the leaves, seeds, bark, and roots, contain antibacterial, anti-inflammatory, antiallergenic, anticarcinogenic, and immunostimulant properties. Its leaves are a rich source of flavonoids, particularly quercetin, which reduces bowel spasms and aids in the anti-diarrheal benefits of guava by relaxing intestinal smooth muscle (Joseph and Priya, 2011; Lok *et al.*, 2023). Quercetin also has antimicrobial characteristics. In lab experiments, guava leaf extract has also shown anti-proliferative action, with activity levels greater than vincristine. In addition to quercetin, other polyphenolic substances found in guava that have been linked to its biological qualities include kaempferol, protocatechuic,

ferulic, ascorbic, gallic, and caffeic acids (Kumar *et al.*, 2021). These polyphenols function as immunostimulants, possibly enhancing immunity and offering antioxidant advantages (Mittal *et al.*, 2010). Guava is a fruit that is packed with nutrients, including dietary fibre, vitamin- A, vitamins- C, folic acid, and minerals like potassium, copper, and manganese that are crucial for human health (Naseer *et al.*, 2018) (Fig.2). One guava fruit has four times the amount of vitamin - C in it than an orange, and it is a low-calorie but extremely healthy addition to any diet thanks to its wide spectrum of critical components. However, cultivation of guava is significantly affected by wilt disease caused by *Fusarium oxysporum f. sp. psidii* (FOP). *Fusarium oxysporum* is widely spread and commonly causes wilt diseases in various fruit crops, for example, in tomatoes (by *Fusarium oxysporum f. sp. Lycopersici* - FOL) (Gangaraj *et al.*, 2022; Mesa *et al.*, 2024). In this review, we have highlighted various therapeutic health benefits of consuming guava.

2020). In addition, guava is packed with water soluble Vitamin C (ascorbic acid), and just one fruit can provide up to 12% of the recommended daily intake of fibre, making it an excellent choice for promoting digestive health (John *et al.*, 2013).

Oral Cavity/dental infections

Guava has been found to be effective in maintaining good oral health. The accumulation of dental plaque is the main cause of periodontitis, a gum disease that can ultimately lead to gingivitis and periodontitis (John *et al.*, 2013). Pathogens such *Porphyromonas gingivalis*, *Aggregatibacter actinomyces emcomitans*, *Fusobacterium nucleatum*, and *Prevotella intermedia* add to the issue (Razak *et al.*, 2006).

However, it has been discovered that the guava’s high quercetin content provides antibacterial properties against these infections. In vulnerable microorganisms, it breaks the cell membrane and deactivates essential proteins (Bansal *et*




Fig.2 : Nutritional values and therapeutic effects of guava





Health Benefits of Guava

Digestive Health (Laxatives)

Guava, a fruit recognised for its high nutritional value, contains a significant amount of dietary fibre, making it an effective treatment for constipation. The guava plant’s sensitive leaves are very high in fibre, which can both prevent and treat haemorrhoids and constipation. With 36 g of dietary fibre per 100 g of fruit, guava is one of the richest sources of fiber, and its seeds are also high-powered laxatives that can help with persistent constipation and clean up the bowel (Kapoor *et al.*,

et al., 2012). Thus, guava extract can effectively stop the growth of dental plaques without interfering with the homeostasis of the oral cavity (Shu *et al.*, 2011). Guava is also well recognized for having a lot of vitamin C in it, which makes it a great cure for scurvy and bleeding gums. Due to its astringent properties, guava is useful in treating toothaches and ulcers. According to Nordini Hasnor *et al.* (2013), chewing guava leaves can instantly relieve toothaches, and the guava leaves’ folate content can prevent foul breath. Guava is commonly utilized in indigenous medicine all around the world to treat a range of oral disorders. In North Sikkim, guava leaves have been

Table 1: The Pharmacological effect of certain important substance found in guava

Parts of guava	Compounds	Molecular structure of Important component (source Pubchem)	Effective property	Reference
Leaves	Phenolic chemicals including quercetin, gallic acid, catechin, epicatechin, rutin, naringenin, and kaempferol	 Quercetin (C15H10O7)	Antimicrobial, Antioxidant and control hypertetion and hyperlipidemia	(Okwu and Ekeke, 2003; Gutierrez-Montiel et al., 2023; Hung-Hui et al., 2011; Ruksiriwanich et al., 2022; Mandal et al., 2022).(Obarzanek et al., 2001; Livingston Raja and Sundar, 2006).
Fruit	Carotenoids (lycopene, beta-carotene, and beta-cryptoxanthin) with ascorbic acid.	 Ascorbic acid (C6H8O6) and Lycopene (C40H56)	Antioxidant, anticancer and maintain healthy digestive system	(Yadav et al., 2010; Teixeira et al., 2003; Gutiérrez et al., 2008; Sharma and Chandola, 2011,Kapoor et al.,2020; Mandal et al., 2022).
Seed	High powdered laxatives, glycosides; Carotenoids, phenolic compounds such as ellagic acid, catechins and quercetin.	 Catechin (C15H14O6)	Antimicrobial and antioxidant activity; prevents the constipation and clean up the bowl; prevents the hypercholesterolemia	(Kapoor et al., 2020; Gutierrez-Montiel et al., 2023; Mandal et al., 2022; Obarzanek et al., 2001.)
Bark	Inorganic phenols such as guajaverin and luteolin	 Guajaverin (C20H18O11)	Potential antimicrobial, antioxidant activities anti-inflammatory and anticancer	(Guo et al., 2003; Dorsch et al., 2005; Lok et al., 2023; Gutierrez-Montiel et al., 2023; Ruksiriwanich et al., 2022; Mandal et al., 2022).

utilized to relieve dental pain and mouth sores (Pradhan and Badola, 2008). Guava leaves have also been used to cure mouth ulcers in Andhra Pradesh, India (Lingaiah and Rao, 2013). Guava leaves have been reported to have therapeutic benefits for treating dental and oral diseases, and guava twigs have been discovered to be useful as “chewing sticks” by traditional healers in Cameroon (Agbor and Naidoo, 2015; Okwu and Ekeke, 2003). The antibacterial effects of guava leaf extract against oral infections have been demonstrated in numerous studies. *Streptococcus* strains, such as *Lactobacillus casei* and *Streptococcus oralis*, which are frequently linked to tooth plaque and cavities, have been demonstrated to be significantly resistant to the extract (Ravi and Divyashree, 2014; Vieira et al., 2014; Alves et al., 2009). Additionally, research has shown that guava leaf extract, when administered at a concentration of 1 mg/ml, is particularly helpful in reducing dental plaque and cavities (Razak et al., 2006). Along with other oral pathogens connected to the early phases of plaque development, the bactericidal effects of guava leaf extract in its aqueous form have also been reported against *Streptococcus sanguinis*, *Streptococcus mitis*, and *Actinomyces species*. Guava leaf extract may be utilized as a natural treatment for dental caries and other oral diseases brought on by these bacteria because the minimum inhibitory concentration (MIC) values for these bacteria ranged from 2.61 to 4.69 mg/ml (Razak et al., 2006).

Antidiabetic Activity

It has been discovered that guava may have anti-diabetic benefits. Guava leaves are typically ingested in China on an empty stomach to manage diabetes. Both the fruits and leaves of the guava plant have been demonstrated in studies on mice at the Medicinal Research Laboratory in Allahabad to effectively lower blood sugar levels. The active chemicals present in guava leaves are believed to be responsible for this effect, as they have the ability to inhibit intestinal glycosidases, thereby reducing postprandial hyperglycemia (Grover et al., 2002). Guava is a fruit with a high fibre content that is known to slow down the absorption of glucose from the gastrointestinal tract. As a result, a sharp increase in blood sugar levels following meals is less likely. Studies have shown that drinking guava tea instead of plain water as a control after eating white rice considerably reduces the rise in blood sugar levels (Sharma and Chandola, 2011). Guava has also been demonstrated to lower fasting blood glucose levels. For instance, one study revealed that giving Type 2 diabetics guava leaf decoction to drink with meals for 90 days reduced their fasting blood glucose levels. Guava fruit powder mixed with buttermilk is also a traditional medicine for diabetes in Andhra Pradesh and Tamil Nadu, India. Similarly, in the central part of Togo, guava was enumerated among the most commonly used plants and fruits for diabetes (Shen et al., 2008; Gutiérrez et al., 2008). Studies on both animals and people have investigated the potential of various guava plant sections as a potential antidiabetic medication (Deguchi and Miyazaki,

2010). In a study by Ojewole, it was discovered that both normal and streptozotocin-treated diabetic rats experienced a hypoglycemic response when given aqueous guava leaf extract orally at various doses (50 to 800 mg/kg) (Ojewole, 2005; Uuh-Narvaez *et al.*, 2021). The ability of the guava plant's fruit and bark to reduce blood sugar levels has also been investigated. The fruit extract (125 and 250 mg/kg) demonstrated significant hypoglycemic efficacy in diabetic animals produced by STZ (streptozotocin), whereas the ethanolic stem bark extract (250 mg/kg) was effective in lowering blood glucose levels in alloxan-induced hyperglycemic rats (Mukhtar *et al.*, 2004; Huang *et al.*, 2011).

Numerous cellular-level investigations have also looked at how various guava leaf extracts affect glucose absorption and aldose reductase inhibitory action. Furthermore, insights from Jayachandran *et al.*, and Beidokhti *et al.*, shed light on guava's influence on insulin signaling proteins and glucose uptake efficiency (Jayachandran *et al.*, 2020; Beidokhti *et al.*, 2020). Guava's potential in managing lipid levels, glucose metabolism, and insulin resistance has been a recurring theme in research (Khan *et al.*, 2013; Xu *et al.*, 2020; Zhu *et al.*, 2020). Luo *et al.*, and Shabbir *et al.*, discuss the antidiabetic attributes of polysaccharides and polyphenols derived from guava leaves, with the latter also exploring the effects of extracts from guava pulp and seeds (Luo *et al.*, 2019; Shabbir *et al.*, 2020). Such antidiabetic effects, as Tella *et al.*, suggests, may be attributed to guava's ability to enhance glycogen synthase levels and restrict the activity of the enzyme glycogen phosphorylase (Tella *et al.*, 2019). Research extends beyond mere observation of glucose level regulation. Cellular studies have delved deep to understand the influence of various guava leaf extracts on glucose absorption and aldose reductase inhibitory action. Such detailed insights collectively suggest the therapeutic potential of guava components like leaves, fruit, and bark in antidiabetic medication and call for intensified research in this direction (Anand *et al.*, 2016).

Anti-Cold and Cough

Due to its high content of iron and vitamin C, guava has been suggested as a potential treatment for cold and cough. These components have been found to alleviate congestion, reduce mucus formation in the respiratory tract, and help combat harmful pathogens. Reports claimed that these components in Guava act like a miracle in curing influenza (Jaiarj *et al.*, 1999). Guava leaves, particularly the young and tender ones, can be used as a decoction to relieve cold and cough symptoms. The fruit itself, whether raw or cooked, has also been found to have benefits. Its astringent properties can help inhibit microbial activity and loosen cough by breaking down mucus polymers. Guava fruit, which contains a lot of vitamin C (ascorbic acid), has also been shown to be useful against viral or bacterial illnesses. Cooked (roasted) ripe guava is a traditional remedy for severe cough and obstruction in several Indian regions. Recent studies have also shown that hydroextracts of guava leaves can significantly reduce coughing frequency induced by irritants like capsaicin aerosol. Diving deeper into its anticough attributes, Jaiarj *et al.* (1999) probed into the anticough potential of guava leaf extracts, specifically on rats and guinea pigs. The researchers employed a dose of 30 μ mol of capsaicin to elicit cough in the animals.

The subjects in the experimental groups received intraperitoneal doses of the extracts at concentrations of 1, 2, and 5 g/kg. The data gleaned from this study underscored that the extracts, when administered, curtailed cough frequency in a dose-proportional manner. Such findings reinforce the therapeutic prowess of Guava in managing cough-related ailments.

Antibacterial Activity

Guava extracts have been discovered to have antibacterial effects against both Gram-positive and Gram-negative species of bacteria. Studies have shown that guava leaves and bark extracts exhibit strong antibacterial activity against multidrug-resistant *Vibrio cholera*, indicating their potential in controlling epidemics of cholera. In some cultures, natural remedies like young and tender guava leaves are preferred for the treatment of infections in children as they exhibit good activity against intestinal microbes, including *Vibrio cholera*. Guava extracts have also been found effective against bacteria such as *E. coli*, which are otherwise resistant to many antibiotics. The antibacterial impact of the majority of essential oils and different guava leaf extracts has been examined using methanolic extract, demonstrating the effective inhibition of bacterial growth, particularly against *Staphylococcus aureus* (*S.aureus*) (Shu *et al.*, 2011; Prabu *et al.*, 2006).

The extraction of benzyl isocyanate from the methanolic extract of guava leaves, which shown inhibitory effect against *S. aureus*, is a remarkable mention by Dutta *et al.*, (2020). *P. aeruginosa*, *C. violaceum*, *S. aureus*, and *S. marcescens* were only a few of the bacteria whose virulence was found to be reduced by extracts from guava leaves (Patel *et al.*, 2019). Human pathogenic bacteria and plant pathogenic fungi, specifically *C. lunata* and *F. chlamyosporum*, were suppressed by the essential oil of *P. guajava*'s senescent leaves (Chaturvedi *et al.*, 2021). The antibacterial effectiveness of guava extracts was demonstrated by Silva *et al.* (2018) against bacterial strains including *S. salivarius*, *Streptococcus mutans*: *Streptococcus mitis*, *Streptococcus sanguinis*, and *Streptococcus sobrinus*. According to a study, the essential oils from guava were tested for their antibacterial activity against *Enterococcus faecalis* and *S. aureus* at a range of concentrations, including 125, 250, 500, and 1000 g/ml (Weli *et al.*, 2019). Strong antibacterial effects were found by Morais-Braga *et al.* (2016b) against bacteria such *E. coli*, *Pseudomonas aeruginosa*, and *S. aureus*, with the extract exhibiting significant activity and having a MIC value of 256 g/mL. The chemicals identified from guava that inhibited the growth of *S. aureus*, *S. epidermidis*, and *Mycobacterium smegmatis* were further clarified by Huang *et al.* (2021).

Anticancer Activity

The presence of the antioxidant lycopene in guava makes it an effective shield in preventing and fighting cancer, especially breast and prostate cancer. Lycopene scavenges free radicals and prevents further formation of them. In cell line models, research has shown that guava leaf extracts have anti-prostate cancer activity, making them a promising anti-androgen-sensitive prostate cancer agent. It also contains carotene, which can help prevent lung and oral cancers (Yadav *et al.*, 2010; Ruksiriwanich *et al.*, 2022; Mandal *et al.*, 2022).

A recent comprehensive review of available literature has explored the potential anticancer properties of guava through laboratory investigations. The review suggests that guava may have various anticancer properties, including the ability to regulate gene expression, modulate cellular signalling pathways, repair DNA damage, and impact cell proliferation and apoptosis (Correa *et al.*, 2016; Lok *et al.*, 2023; Mandal *et al.*, 2022).

Taking MCF-7 cells as a model, Lin and Lin (2020) evaluated the effects of guava seed polysaccharides. The outcomes showed that MCF-7 cell viability was significantly and dose-dependently inhibited. Similarly, dos Santos *et al.* (2018) investigated the anticancer effects of red guava extracts. Their results demonstrated the potency of an extract from guava high in lycopene, with an IC₅₀ value of 29.85 at 5.964 g/mL. Guava was shown to have particular chemicals that had anticancer effects on HCT116 and HT29 cells, according to Zhu *et al.* (2019). Guava has been shown to have potent inhibitory activity against cancer cells by Ashraf *et al.* (2016), as shown by an IC₅₀ value of less than 30 mg/ml. Tetracosane, vitamin E, and g-sitosterol, three of the guava's bioactive components, are thought to be the source of the fruit's anticancer effects (Ryu *et al.*, 2021).

Bazioli *et al.* (2020) looked into the *P. guajava* extracts' antiestrogenic properties. Studies on 17- β -estradiol injected MCF7 BUS cells and female Wistar rats were conducted in vitro and in vivo, respectively. One million MCF7 BUS cells per milliliter in the presence of 10-9M 0.3, 0.6, 1.25, and 2.5 μ g/mL of fractionated *P. guajava* extracts were used to treat 17- β -estradiol. The rats in the experiment were given 0.3 mg/kg of 17- β -estradiol as an injection, and they were given extracts fractionated from *P. guajava* at doses of 12.5, 25, and 50 mg/kg for three days. The findings from experiments conducted in vitro and in vivo demonstrated that the injection of extracts inhibited the proliferative activity of 17- β -estradiol, indicating its antiproliferative properties.

Antidiarrheal property

Guava leaves have been found to possess strong antidiarrheal properties. Consuming 6-10 fresh and tender guava leaves boiled in warm water on an empty stomach can effectively control diarrhoea. The leaves contain high levels of flavonoids, which contribute to their antimicrobial activity against pathogens responsible for diarrhoea, such as *giardia* and *rotavirus* (Birdi *et al.*, 2010; Gutierrez-Montiel *et al.*, 2023). Guava bark has been traditionally used as a treatment for diarrhoea in children due to its astringent properties. On the other hand, a tea made from guava leaves or an extract of guava leaves in warm water can help promote bowel movement. A study found that guava leaf extract may be helpful for lowering the severity and frequency of rat castor oil-induced diarrhoea. According to Lozoya *et al.* (2002), these results were attributed to the extract's influence on intestinal peristalsis. A study by Koriem *et al.* (2019) looked at how guava extracts helped rats with osmotic diarrhoea that was caused by a lactose-containing diet. Desmopressin served as the reference medication, and the rats received daily doses of 50 and 100 mg/kg of the extract. Getting the serum electrolytes and urine

output back to normal showed that the extract could stop diarrhoea. Moreover, the level of lipid peroxidation dropped. The antidiarrheal properties of guava extracts in rats against enteropathogenic *Escherichia coli* were examined in a different study by Hirudkar *et al.* (2020). The study's findings demonstrated that giving the subjects the extract reduced their weight, frequency of feces, and overall frequency of diarrheal stools. By examining the recovery of WBC, Hb, and platelet values, this experiment further demonstrated the extract's antidiarrheal capabilities.

Antihypertensive and Hypolipidemic

Guava is well known for its beneficial effects on heart disease, hypertension, and hyperlipidemia. Its high potassium level aids in blood vessel relaxation, which lowers blood pressure. Due to its greater potassium and fiber content, regular eating of guava fruit has been demonstrated to considerably lower blood pressure and blood lipids. Guavas have a high amount of pectin, which has been discovered to slow down food absorption and significantly lower blood lipid levels as well as the risk of cardiovascular diseases (Singh *et al.*, 1993).

Gallic acid, catechins, epicatechins, rutin, naringenin, and kaempferol are among the compounds found in guava leaves that have been demonstrated to inhibit the pancreatic cholesterol esterase enzyme, lowering blood levels of lipids (including cholesterol). One of these compounds that is essential for avoiding hypercholesterolemia is catechins (Deguchi and Miyazaki, 2010). Guava's quercetin has also been linked to a lower risk of cardiac and stroke mortality brought on by hypertension and hyperlipidaemia (Obarzanek *et al.*, 2001). Guava's modest potassium content is what gives it its capacity to strengthen heart health and prevent stroke.

Antacid and Ulcer Protectant Activity

Guava has long been utilised as a homoeopathic treatment for ulcers and hyperacidity. Guava leaves are good at reducing stomach hyperacidity due to their alkaline composition. Young guava leaves are boiled in water in many areas to make guava tea, which is then served hot to reduce acidity. Research has shown that the methanolic extract of guava has antacid and ulcer healing properties, with the highest efficacy observed *in vitro* (Livingston Raja and Sundar, 2006).

Flavonoids and saponins present in both guava fruit and guava leaves have been found to counteract acidity and prevent ulceration of the stomach. The methanolic extract of guava leaves (500 mg and 1000 mg/kg body weight) significantly lowers ulcer index in ethanol-induced ulcers, according to studies in rats (Uduak *et al.*, 2012).

Wound Healing

For centuries, Guava leaves have been used as a traditional remedy for wound healing. Ancient people in India and China would grind the leaves and made into paste with water or oil content and apply it directly to the wound. Studies have shown that the tannins and flavonoids in Guava leaves can promote faster healing of wounds when applied topically in the form of a methanolic extract. In fact, ointments made from Guava leaves have been found to be more effective than commercially available wound care products. To produce the ointment, guava leaves are first washed, ground, and extracted with oil. The resulting extract is then mixed with a vehicle, which is

typically melted candle wax, to improve absorption. The resulting compound can be applied twice daily for four days to treat wounds (Okoli *et al.*, 2009).

Gastrointestinal Problems

Guava leaves contain quercetin and flavonoids that have been shown to effectively treat a range of gastrointestinal issues (Arima and Danno, 2002). The alkaline properties of guava discourage the growth of harmful microbes that can cause gastroenteritis. Both the fruit and leaves of the guava plant can help alleviate diarrhoea by inhibiting the growth of microbes and binding loose stools. Furthermore, guava is rich in vital vitamins and minerals like vitamin C, potassium, and carotenoids, which help maintain a healthy digestive system (Arima and Danno, 2002; Teixeira *et al.*, 2003). Consuming guava leaves on an empty stomach can aid in the removal of extra mucus from the large intestine, while occasionally sipping guava leaf tea can help control the consistency and degree of stiffness of the stools. Guava has long been a well-liked traditional treatment for gastrointestinal illnesses such as diarrhoea, dysentery, indigestion, and stomach aches (Gutiérrez *et al.*, 2008). For instance, in Mexico, guava was mentioned as one of the species used to cure gastrointestinal diseases, while in India, local traditional healers advise drinking hot water with guava bark powder to treat blood-associated diarrhoea (Heinrich *et al.*, 1998). Guava leaves have been used to treat diarrhea in Maharashtra and South Assam (Tetali *et al.*, 2009; Choudhury *et al.*, 2015). There have been a lot of laboratory studies on the potential medical use of guava leaves. By inhibiting diarrheal bacteria, regulating intestinal motility, lowering the frequency of stools, and promoting gastric emptying, guava leaves have been demonstrated to have anti-diarrheal efficacy in both in vitro and animal experiments.

Guava leaf extracts at 5% and 10% concentrations are very effective against pathogens like *Shigella flexneri* and *Vibrio cholerae* that cause bacterial diarrhea (Birdi *et al.*, 2010; Birdi *et al.*, 2011). Tona *et al.* (1998) also found that guava leaf extracts stop *Entamoeba histolytica* from growing and lessen the contractions caused by acetylcholine and/or KCl solution. Guava leaf extracts are able to postpone the onset of diarrhoea, lessen the frequency of faeces, and lessen the severity of the condition, according to tests conducted on animals. Furthermore, they reduce the rate at which rats move charcoal food and block prostaglandin E2 enteropooling (Lin *et al.*, 2002). Guava leaf decoction (1%, 5%, and 10%) has been found to inhibit colonization of bacteria on the epithelial cells, toxin production, and inflammatory response (Brijesh *et al.*, 2011) and has been shown to promote gastric ulcer healing and reduce capillary permeability in the abdominal cavity (Livingston Raja and Sundar, 2006). Although quercetin is a significant component in guava leaf, it has been discovered that the crude guava decoction has greater anti-diarrheal efficacy than quercetin by itself (Birdi *et al.*, 2010). As a result, guava leaves show potential for treating both acute and chronic diarrhoea brought on by a variety of bacterial infections.

Anti-Allergies

The possible anti-allergy effects of *Psidium guajava* leaves have been studied recently. The activation of T regulatory cells from CD⁴⁺ splenocytes of C57BL/6 mice by IL-10 and the

release of histamine by mast cells can both be inhibited by the methanol and aqueous extracts of the leaves, according to studies. Furthermore, it has been found that the extracts reduce T regulatory cell activity, which shifts the Th1/Th2 balance to a Th1-dominant state. Guava leaf extracts have been demonstrated in animal experiments to lessen allergy reactions brought on by T cells. According to these findings, guava leaves may be helpful in the treatment of allergies (Seo *et al.*, 2005).

Anti- Inflammatory

In traditional medicine, guava leaves have been used to treat inflammatory conditions including rheumatism (Gutiérrez *et al.*, 2008). Studies have demonstrated that guava leaf extracts, whether in methanol or aqueous form, have anti-inflammatory and antipyretic activities. In an experiment involving rats with carrageenan-induced paw edema, the methanol extract of guava leaves displayed anti-inflammatory benefits at dosages of 50, 100, and 200 mg/kg (Olajide *et al.*, 1999). Similar to this, an aqueous extract of guava leaves was given to rats with fresh egg albumin-induced edema via IP (50–800 mg/kg), which decreased acute inflammation (Han *et al.*, 2011). Guava leaves have an extensive tradition of use in India as rheumatism medication. In addition, mice with yeast-induced hyperpyrexia were given guava leaf methanol extract at doses of 50, 100, and 200 mg/kg, which had antipyretic effects similar to those of regular indomethacin (5 mg/kg) (Olajide *et al.*, 1999).

Skin infections

Many cultures have a long history of using guava as a traditional remedy for skin problems. In the Philippines, for instance, natives use guava leaves to cure scabies (Ong and Kim, 2014), and Tahiti and Samoa have used guava shoots as a skin tonic (Han, 1998). Other countries with a history of employing various guava portions for skin-related illnesses include Panama, Bolivia, Venezuela, Fiji, and Senegal, according to Gupta *et al.* (2011). Studies have shown that guava leaves have antibacterial properties and are effective against clinical isolates of bacteria like *Proteus mirabilis*, *Streptococcus pyogenes*, *E. coli*, *S. aureus*, and *Pseudomonas aeruginosa*, which are frequently linked to surgical wounds, burns, and skin and soft tissue infections (Abubakar, 2009). Atopic skin lesions in mice brought on by 2,4-dinitrochlorobenzene have been successfully treated with cream containing guava leaf extract (Choi *et al.*, 2012) and guava leaf extract in the form of ethyl acetate has shown to be efficient in treating atopic dermatitis (Han *et al.*, 2011).

Women's health

In traditional medicine, guava has a long history of use for female health-related ailments. Tahiti, for example, has utilized guava to treat early labor, uterine bleeding, painful menstruation, and miscarriages in women (Han, 1998). Guava has been identified as a treatment for blennorrhoea, irregular menstrual cycles, and postpartum bleeding in a study (de Boer and Cotingting, 2014). Guava leaf decoction has been used to treat menstrual irregularities in Durgapur, Bangladesh (Khan *et al.*, 2015). A double-blinded, randomized clinical trial including 197 women with primary dysmenorrhoea was conducted to evaluate the effectiveness of guava leaf extract. Ibuprofen (1200 mg/day), guava leaf extract (3 mg and 6 mg flavanol/day), and placebo (3 mg/day) were given to each of

the four experimental groups. To measure how much the stomach discomfort diminished during the study, a visual analogue scale was employed. According to Dougova *et al.* (2007), the group receiving 6 mg/day of guava leaf extract experienced significantly decreased overall pain.

Respiratory infections

Guava leaves have long been utilised as a cough treatment in a number of places, including Guerrero, Mexico, Malaysia, South Africa, Arunachal Pradesh, and North Sikkim, India (Daswani *et al.*, 2017; Namsa *et al.*, 2011; Pradhan and Badola, 2008). Guava has been used to cure tuberculosis traditionally in various parts of the world. For instance, guava bark has been used for this purpose in Nigeria, while guava leaves have been used to treat HIV-AIDS, tuberculosis in Tanzania (Ogbole and Ajaiyeoba, 2009; Kisangau *et al.*, 2007). Animals may benefit from aqueous guava leaf extract as a cough suppressant. In guinea pigs and rats, oral administration of the extract at doses of 2 g/kg and 5 g/kg considerably reduced cough frequency by 35% and 54%, respectively (Jaiarj *et al.*, 1999). Additionally, *Cryptococcus neoformans*, *Klebsiella pneumonia*, *Moraxella catarrhalis*, *Mycobacterium smegmatis*, and *S. aureus* have been demonstrated to be eliminated by guava leaf extract and dichloromethane-methanol extract (York, 2012). Although additional research is required to prove the plant's efficacy and safety, these traditional applications of guava raise the possibility that it may have therapeutic promise for the treatment of tuberculosis.

Cardiac and hypertension

Guava leaves have been used to treat hypertension for centuries in several nations, including Cuba, Nigeria, and Togo (Mesa, 2014; Borokini and Clement, 2013; Karou *et al.*, 2011). In laboratory tests, aqueous guava leaf extract was shown to have cardio preventive benefits and to drastically lower blood pressure and heart rates in hypertensive rats (Ojewole, 2005). Furthermore, rats' intracellular calcium release has been shown to be inhibited by guava leaf extracts, and the aorta rings are markedly constricted in a dose-dependent manner (0.25–2 mg/ml) (Belemtougri *et al.*, 2006; Olatunji Bello *et al.*, 2007). Clinical studies have shown that guava fruit consumption can have positive benefits on cardiovascular health. The consumption of guava fruit has been shown to improve high-density lipoprotein cholesterol levels while decreasing serum total cholesterol, triglycerides, and blood pressure in treated patients (Singh *et al.*, 1992). In another clinical research, hypertension patients were given a potassium- and fibre-rich diet that included 0.5–1 kg of guava per day for four weeks (Singh *et al.*, 1993). These patients' diastolic and systolic blood pressures improved. Additionally, 400 gm of guava fruit per day for nine weeks was found to reduce oxidative stress and blood cholesterol levels in a clinical experiment conducted in Malaysia (Rahmat *et al.*, 2004). These suggest that guava fruit may be a valuable dietary intervention for improving cardiovascular health.

Antioxidant activity

Recent research has shown that guava possesses exceptional antioxidant qualities (Okwu and Ekeke, 2003; Kapoor *et al.*, 2020; Mandal *et al.*, 2022). The methanolic extract of guava leaves exhibits strong antioxidant activity. Quercetin, quercetin-

3-O-glucopyranoside, morin, ascorbic acid, carotenoids, and polyphenols are only a few of the advantageous phytochemicals that contribute to this activity. These substances are essential for preventing oxidative stress and neutralising dangerous free radicals in the body (Jimenez-Escrig *et al.*, 2001).

The Guava plant's antioxidant potential emphasises its relevance as a great source of naturally occurring antioxidants with possible health benefits. Tan *et al.* (2020) embarked on research to compare the antioxidative capacities of extracts from red and white Guava fruits using varying drying procedures. Their study highlighted that the red Guava powders obtained via vacuum drying exhibited the most significant antioxidant activity. Sobral-Souza *et al.* (2018) focused on the antioxidative capabilities of flavonoids derived from Guava extracts. In another study, Luo *et al.* (2019) revealed the impressive antioxidant properties of polysaccharides from guava leaves. These polysaccharides augmented the overall antioxidant activity and the action of the superoxide dismutase (SOD) enzyme while simultaneously reducing MDA (malondialdehyde) activities in diabetic mice induced by STZ. Moreover, a methanolic extract of Guava leaf was shown to possess antioxidant qualities (Khedr *et al.* (2021).

According to Zahin *et al.* (2017), guava fractionated in methanol has greater antioxidant activity than guava fractionated in other solvents such as gasoline, benzene, acetone, and ethyl acetate. From guava, Feng *et al.* (2015) extracted a number of chemicals and evaluated their antioxidant capacities. When tested using a DPPH (2,2-diphenyl-1-picrylhydrazyl) assay, the substances guavinoside C, guavinoside F, quercetin, quercetin-3-O-a-L-arabinofuranoside, and quercetin-3-O-a-L-arabinopyranoside were discovered to possess potent antioxidative properties. According to Hartati *et al.* (2020), the guava extracts' strong antioxidant activity may be largely related to their high phenolic component content.

Hepatoprotective activity

The leaves of guava have been found to possess a hepatoprotective activity (Roy *et al.*, 2006). In an experiment, the aqueous extract of these leaves was shown to protect healthy clone 9 liver cells from injury induced by a 5% alcohol concentration for 30 minutes. Among various extracts tested, hot water extracts exhibited high hepatoprotective effects with lower cytotoxicity. These findings suggest the potential use of guava as a natural remedy for protecting the liver (Hung-Hui *et al.*, 2011; Luiza and Gustavo, 2017).

Studies have shown that *Psidium guajava* extract has hepatoprotective properties at dosages of 250 and 500 mg/kg. The aqueous leaf extract was found to significantly reduce high levels of alanine aminotransferase, alkaline phosphatase, bilirubin, and aspartate aminotransferase when administered orally to rats with acute liver injury brought on by hepatotoxins (Roy *et al.*, 2006). Guava leaf extract could be employed as a drug to treat and protect the liver. These imply that *Psidium guajava* extract may be helpful in the treatment of hepatic diseases. Vijayakumar *et al.* (2020) also evaluated the hepatoprotective properties of guava against liver damage brought on by carbon tetrachloride in rats. Carbon tetrachloride (CCl₄) at the amount of 1.5 mL/kg was intentionally given to

these rats to cause hepatotoxicity. The liver biomarkers (ALT, AST, ALP, and GGT) that were raised as a result of the CCl₄ exposure were decreased after the extract was taken orally every day for 21 days. Additionally, Saber *et al.* (2018) investigated how Guava and *P. cattleianum* worked together to protect rats' livers from liver toxicity brought on by paracetamol. They discovered that giving the extract at doses of 250 and 500 mg/kg significantly decreased the increased levels of liver enzymes, hence reducing hepatotoxicity. In a separate study, Vijayakumar *et al.* (2018) found that administering guava extract to rats with liver damage brought on by exposure to CCl₄ at doses of 100, 200, and 300 mg/kg, bw, and 20 mg/kg of the quercetin component resulted in a reduction in lipid metabolism. The capacity of triterpenoid-enriched guava leaf extract to lower blood ALT (alanine transaminase) and AST (aspartate transaminase) levels, as well as liver ROS (reactive oxygen species) and MDA in mice exposed to acetaminophen, was emphasized by Li *et al.* in (2021). Furthermore, according to Li *et al.* (2020), giving mice which suffered liver damage as a result of acetaminophen exposure 100 mg/kg/day of Guavinoside B, derived from guava fruit, significantly improved both serum and hepatic biochemical markers.

Nephroprotective Properties

Extraction from Guava has nephroprotective qualities. The ability of guava extracts to prevent kidney impairment brought on by some medications and medical disorders has been shown in studies utilising animal models. For instance, ethanolic extracts of guava leaves were reported to prevent renal impairment brought on by paracetamol when given orally and to maintain blood urea, blood creatinine, urine sodium, and creatinine in a dose-dependent manner when given at 200 and 400 mg/kg. The study of histopathology corroborated this observation (Patel *et al.*, 2011). In a similar way, the extracts were reported to shield rats against cisplatin-induced nephrotoxicity (Patel *et al.*, 2012). Additionally, research on diabetic rats has revealed that guava fruit extract can prevent kidney damage (Lin and Yin, 2012).

Immunomodulatory activity

In one study, it was found that a decoction of guava leaves successfully killed a strain of *E. coli* that generates a heat-stable toxin in the murine monocytic cell line J774 by stimulating macrophages (Birdi *et al.*, 2014). In antigen-stimulated mast cells, the ethyl acetate fraction of guava leaves reduced COX-2 expression, cytokine secretion, degranulation, and FcεRI-mediated signaling (Han *et al.*, 2011). In an in vitro model system utilizing Labeo rohita head kidney macrophages, a flavonoid fraction of guava leaf extract also demonstrated promise in regulating nuclear factor KB activation (Sen *et al.*, 2015).

Veterinary medicine

Guava leaves and buds have been found to possess antidiarrheal activity and have been used in veterinary medicine (Lans *et al.*, 2000). For treating diarrhoea in ruminants, sheep, and goats, Javan farmers use various guava plant species. In particular, guava leaves are used to treat diarrhoea in ruminants, while sheep and goats are treated for the same ailment with the plant's root and stem bark (Mathias-Mundy

and Murdiati, 1990). In Trinidad, for horses suffering from diarrhoea, a mixture of boiled guava leaves, young fruits, and/or buds is prepared with mash or bran (Lans *et al.*, 2006). Meanwhile, in the Philippines, adding dried and ground guava leaf meal to pig diets in proportions of 5% and 10% has been found to be effective in reducing diarrhoea in piglets. In Thailand, supplementing the diet of weaned piglets with ground guava leaves has been observed to prevent post-weaning diarrhoea (Tartrakoon *et al.*, 2005). Furthermore, guava has demonstrated antibacterial activity against pathogenic bacteria in pigs.

Downsides of Guava Consumption

Guava intake, like any other food, has some downsides to consider. Allergic responses to guava are uncommon but possible, so people with documented fruit allergies should proceed with caution (Cubero *et al.*, 2005). Guava's oxalates content, which can help cause kidney stones in those who are sensitive, is one cause for concern (Moein *et al.*, 2015). Further, the vitamin K concentration in guava may interact with anticoagulant drugs, reducing their effectiveness (Alqahtani *et al.*, 2016). Another factor to consider is that the mild acidity of guava can lead to dental enamel loss and tooth discomfort if ingested in large quantities (Al-Dlaigan *et al.*, 2017).

CONCLUSION

Over the past decade, substantial studies have proven the ethnomedicinal uses of Guava leaves, establishing their promise as a cure for many common ailments worldwide. Guava leaves contain quercetin, catechin, gallic acid, peltatoside, hyperoside, isoquercetin, and guaijaverin, which offer health advantages. Guava skin contains polyphenols, vitamins A and C, iron, phosphorus, calcium, and other minerals. It harbours anti-cancer phenolic chemicals that reduce skin aging. Guava leaves act as antioxidants, bacteriostatic, and fungistatic. Due to these traits, *Psidium guajava* can treat and prevent several diseases. Public health is increasingly burdened by chronic degenerative diseases such as diabetes, dyslipidemia, metabolic syndrome, and cardiovascular disease. Researchers are studying the bioactive chemicals present in *P. guajava* leaves, bark, and seeds to address this rising concern. Medicinal plants like guava may reduce disease risk factors and offer alternative treatments, considering the high expenses associated with allopathic therapies.

REFERENCES

- Abubakar, E. M. 2009. The use of *Psidium guajava* Linn. in treating wound, skin and soft tissue infections. *Sci Res Essays*. **4**: 605-611.
- Agbor, M. A. and Naidoo, S. 2015. Ethnomedicinal plants used by traditional healers to treat oral health problems in Cameroon. *Evid Based Complement Alternat Med*. 649832.
- Al-Dlaigan, Y. H., *et al.* 2017. Acidogenicity and erosive potential of commonly consumed beverages in Saudi preschool children. *J. Clinical Pediatric Dentistry*. **41(4)**: 272-278.
- Alqahtani, S., *et al.* 2016. Nutritional content and health benefits of guava (*Psidium guajava*). *International J. PharmTech Research*. **9(10)**:

161-171.

Alves, P. M., Queiroz, L. M., Pereira, J. V. and Pereira, Mdo, S. 2009. In vitro antimicrobial, antiadherent and antifungal activity of Brazilian medicinal plants on oral biofilm microorganisms and strains of the genus *Candida*. *Rev Soc Bras Med Trop.* **42**: 222-224.

Anand, S., Arasakumari, M., Prabu, P. and Amalraj, A. J. 2016. Antidiabetic and aldose reductase inhibitory potential of *Psidium guajava* by in vitro analysis. *Int J. Pharm Pharm Sci.* **8**: 271-276.

Arima, H. and Danno G. 2002. Isolation of antimicrobial compounds from guava (*Psidium guajava* L.). *Biosc Biotech Bioch.* **66**: 727-730.

Ashraf, A., Sarfraz, R.A., Rashid, M.A., Mahmood, A., Shahid, M., Noor, N. 2016. Chemical composition, antioxidant, antitumor, anticancer and cytotoxic effects of *Psidium guajava* leaf extracts. *Pharm. Biol.* **54**(10): 1971–1981. <https://doi.org/10.3109/13880209.2015.1137604>.

Bansal, S., Rastogi, S. and Bajpai, M. 2012. Mechanical, chemical and herbal aspects of periodontitis: A review. *International J. Pharmaceutical Sciences and Research.* **3**: 1260-1267.

Barbalho, S. M., Damasceno, D. C., Spada, A. P., da Silva, V.S. and Martuchi, K. A. et al. 2011. Metabolic Profile of Offspring from Diabetic Wistar Rats Treated with *Mentha piperita* (Peppermint). *Evid Based Complement Alternat Med.* 430237.

Bazioli, J. M., Costa, J. H., Shiozawa, L., Ruiz, A.L.T.G., Foglio, M. A. and Carvalho, J. E., 2020. Anti-estrogenic activity of guajadial fraction, from guava leaves (*Psidium guajava* L.). *Molecules.* **25** (7): 1525. <https://doi.org/10.3390/molecules25071525>.

Beidokhti, M. N., Eid, H. M., Villavicencio, M. L. S., Jaeger, A.K., Lobbens, E. S., Rasoanaivo, P. R., McNair, L. M., Haddad, P. S., and Staerk, D. 2020. Evaluation of the antidiabetic potential of *Psidium guajava* L. (Myrtaceae) using assays for α -glucosidase, α -amylase, muscle glucose uptake, liver glucose production, and triglyceride accumulation in adipocytes. *J. Ethnopharmacol.* **257**: 1–8. <https://doi.org/10.1016/j.jep.2020.112877>.

Belemtougri, R. G., Constantin, B., Cognard, C., Raymond, G. and Sawadogo, L. 2006. Effects of two medicinal plants *Psidium guajava* L. (Myrtaceae) and *Diospyros mespiliformis* L. (Ebenaceae) leaf extracts on rat skeletal muscle cells in primary culture. *J Zhejiang Univ Sci B.* **7**: 56-63.

Birdi, T., Daswani, P., Brijesh, S., Tetali, P. and Natu, A. et al. 2010. Newer insights into the mechanism of action of *Psidium guajava* L. leaves in infectious diarrhoea. *BMC Complement Altern Med.* **10**, 33.

Birdi, T. J., Brijesh, S. and Daswani, P. G. 2014. Bactericidal effect of selected anti-diarrheal medicinal plants on intracellular heat-stable enterotoxin-producing *Escherichia coli*. *Indian J Pharm Sci.* **76**: 229-235.

Birdi, T. J., Daswani, P. G., Brijesh, S. and Tetali, P. 2011. In vitro anti-giardial and antirotaviral activity of *Psidium guajava* L. leaves. *Indian J Pharmacol.* **43**: 616-617.

Borokini, T. I. and Clement, M. 2013. Ethnobiological survey of traditional medicine practice for fevers and headaches in Oyo State, Nigeria. *J Herb Med.* **2**: 121-130.

Brijesh, S., Tetali, P. and Birdi, T. J. 2011. Study on effect of anti-diarrheal medicinal plants on enteropathogenic *Escherichia coli* induced interleukin 8 secretion by intestinal epithelial cells. *Altern Med Stud.* **1**: e16.

Carrasco-Castilla, J., Hernandez-Álvarez, A.J., Jiménez-Martínez, C., Jacinto-Hernández, C., Alvarado-Gil, J. J. and Davila-Ortiz G. 2012. Antioxidant and antimicrobial activities of phenolic extracts from guava (*Psidium guajava* L.) leaves. *Food Science and Technology International.* **18**(5): 387-396.

Chaturvedi, T., Singh, S., Nishad, I., Kumar, A., Tiwari, N., Tandon,

S., Saikia, D., Verma, R.S. 2021. Chemical composition and antimicrobial activity of the essential oil of senescent leaves of guava (*Psidium guajava* L.). *Nat. Prod. Res.* **35** (8): 1393–1397. <https://doi.org/10.1080/14786419.2019.1648462>.

Chen, H.Y. and Yen, G. C. 2007. Antioxidant activity and free radical scavenging capacity of extracts from guava (*Psidium guajava* L.) leaves. *Food Chemistry.* **101**: 686-694.

Choi, J.H., Park, B.H., Kim, H.G., Hwang, Y.P., Han, E.H. and Jin, S. W. et al., 2012. Inhibitory effect of *Psidium guajava* water extract in the development of 2,4-dinitrochlorobenzene-induced atopic dermatitis in NC/Nga mice. *Food Chem Toxicol.* **50**: 2923-2929.

Choudhury, P. R., Choudhury, M. D., Ningthoujam, S. S., Mitra, A., Nath, D. and Talukdar, A. D. 2015. Plant utilization against digestive system disorder in Southern Assam, India. *J Ethnopharmacol.* **175**: 192-197.

Correa, M. G., Couto, J. S. and Teodoro, A. J. 2016. Anticancer properties of *Psidium guajava* – A mini review. *Asian Pac J Cancer Prev.* **17**: 4199-4204.

Cubero, J. I. et al., 2005. Anaphylaxis caused by ingestion of guava: a case report. *J. Investigational Allergy and Clinical Immunology.* **15**(1): 68-70.

Daswani, P.G., Gholkar, M. S. and Birdi, T. J. 2017. *Psidium guajava*: A single plant for multiple health problems of Rural Indian population. *Pharmacognosy Rev.* **11**(22): 167-174.

de Boer, H. J. and Cotingting, C. 2014. Medicinal plants for women's healthcare in southeast Asia: A meta-analysis of their traditional use, chemical constituents, and pharmacology. *J Ethnopharmacol.* **151**: 747-767.

Deguchi, Y., Miyazaki, K. 2010. Anti-hyperglycemic and anti-hyperlipidemic effects of guava leaf extract. *Nutr Metab (Lond).* **7**, 9.

Dorsch, S., Hemmerling, M. and Werfel, T. 2005. Luteolin inhibits IgE-dependent activation of basophils through suppression of NF- κ B and AP-1. *J. Allergy and Clinical Immunology.* **116**(3): 683-690.

dos Santos, R. C., Ombredane, A. S., Souza, J. M.T., Vasconcelos, A. G., Placido, A., Amorim, A. D. G. N., Barbosa, E. A., Lima, F. C. D. A., Ropke, C.D., Alves, M.M.M., Arcanjo, D.D.R., Carvalho, F. A. A., Delerue-Matos, C., Joanitti, G.A., Leite, J.R.S.A., 2018. Lycopene-rich extract from red guava (*Psidium guajava* L.) displays cytotoxic effect against human breast adenocarcinoma cell line MCF-7 via an apoptotic-like pathway. *Food Res. Int.* **105**: 184–196. <https://doi.org/10.1016/j.foodres.2017.10.045>.

Dobova, S. V., Morales, H. R., Hernández, S. F., del Carmen Martínez García, M., de Cossío Ortiz, M. G. and Soto, M. A. et al. 2007. Effect of a *Psidium guajava* folium extract in the treatment of primary dysmenorrhea: A randomized clinical trial. *J Ethnopharmacol.* **110**: 305-310.

Dutta, K., Karmakar, A., Jana, D., Ballav, S., Shityakov, S., Panda, A.K., Ghosh, C. 2020. Benzyl isocyanate isolated from the leaves of *Psidium guajava* inhibits *S.aureus* biofilm formation. *Biofouling.* **36**(8):1000–1017. <https://doi.org/10.1080/08927014.2020.1842877>.

Feng, X.H., Wang, Z.H., Meng, D.L., Li, X. 2015. Cytotoxic and antioxidant constituents from the leaves of *Psidium guajava*. *Bioorg. Med. Chem. Lett.* **25** (10): 2193–2198. <https://doi.org/10.1016/j.bmcl.2015.03.058>.

Gangaraj, R., Nagaraja, A., Gaba, S., Das, A., Prameeladevi, D., ebbarma, R. and Choudhary, S. P., et al. 2022. Occurrence, identification and pathogenicity of *Fusarium* species associated with guava wilt disease in India. *Arch. Phytopathol. Plant Prot.* **55**, 175–197. [10.1080/03235408.2021.2005368](https://doi.org/10.1080/03235408.2021.2005368)

Grover, J. K., Yadav, S. and Vats, V. 2002. Medicinal plants of India

with antidiabetic potential. *J Ethnopharmacol.* **81**: 81-100.

Guo, C., Yang, J., Wei, J., Li, Y., Xu, J. and Jiang, Y. 2003. Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutr Res.* **23(12)**:1719-1726.

Gupta, G. K., Chahal, J. and Arora, D. 2011. Psidium guajava Linn.: Current research and future prospects. *J Pharm Res.* **4**: 42-46.

Gutiérrez, R.M., Mitchell, S. and Solis, R. V. 2008. Psidium guajava: A review of its traditional uses, phytochemistry and pharmacology. *J Ethnopharmacol.* **117**:1-27.

Gutiérrez-Montiel, D., Guerrero-Barrera, A. L., Chávez-Vela, N. A., Avelar-Gonzalez, F. J. and Ornelas-García, I. G. 2023. Psidium guajava L.: From byproduct and use in traditional Mexican medicine to antimicrobial agent. *Front. Nutr.* **10**:1108306. doi: 10.3389/fnut.2023.1108306

Han, E. H., Hwang, Y.P., Choi, J. H., Yang, J. H., Seo, J. K. and Chung, Y. C. et al., 2011. Psidium guajava extract inhibits thymus and activation-regulated chemokine (TARC/CCL17) production in human keratinocytes by inducing heme oxygenase 1 and blocking NF- κ B and STAT1 activation. *Environ Toxicol Pharmacol.* **32**: 136-145.

Han, E. H., Hwang, Y. P., Kim, H. G., Park, J. H., Choi, J. H. and Im J. H. et al., 2011. Ethyl acetate extract of Psidium guajava inhibits IgE-mediated allergic responses by blocking Fc ϵ R1 signaling. *Food Chem Toxicol.* **49**: 100-108.

Han, S. T. 1998. Medicinal Plants in the South Pacific. Western Pacific Series. Manila: *World Health Organization (WHO) Regional Publications*, p. 254.

Hartati, R., Nadifan, H. I., Fidrianny, I., 2020. Crystal guava (Psidium guajava L. "Crystal"): evaluation of in vitro antioxidant capacities and phytochemical content. *Sci. World J.* <https://doi.org/10.1155/2020/9413727>.

Heinrich, M., Ankli, A., Frei, B., Weimann, C. and Sticher O. 1998. Medicinal plants in Mexico: Healers' consensus and cultural importance. *Soc Sci Med.* **47**: 1859-1871.

Hirudkar, J. R., Parmar, K. M., Prasad, R. S., Sinha, S. K., Lomte, A. D., Itankar, P. R. and Prasad, S.K. 2020. The anti-diarrhoeal evaluation of Psidium guajava L. against enteropathogenic Escherichia coli induced infectious diarrhoea. *J. Ethnopharmacol.* **251** (112561). <https://doi.org/10.1016/j.jep.2020.112561> <https://doi.org/10.1016/j.jep.2020.112561>

Huang, C. S., Yin, M. C., Chiu, L. C. 2011. Antihyperglycemic and antioxidative potential of Psidium guajava fruit in streptozotocin-induced diabetic rats. *Food Chem Toxicol.* **49**:2189-2195.

Huang, J., Li, C., Ma, J., Xu, K., Chen, X., Jiang, J., Zhang, D. 2021. Chemical constituents of Psidium guajava leaves and their antibacterial activity. *Phytochemistry.* **186**,. <https://doi.org/10.1016/j.phytochem.2021.112746>

Hung-Hui, C., Po-Hua, W., Diana, L., Yun-Chiehand, P. and MingChang, W. 2011. Hepatoprotective Effect of Guava (*Psidium guajava* L.) Leaf Extracts on Ethanol-Induced Injury on Clone 9 Rat Liver Cells. *Food and Nutrition Sciences.* **2**: 983-988.

Jaiarj, P., Khoahaswan, P., Wongkrajang, Y., Peungvicha, P., Suriyawong, P., Saraya, M. L., Ruangsomboon, O. 1999. Anticough and antimicrobial activities of Psidium guajava Linn. leaf extract. *J. Ethnopharmacol.* **67(2)**:203-212. [https://doi.org/10.1016/s0378-8741\(99\)00022-7](https://doi.org/10.1016/s0378-8741(99)00022-7).

Jayachandran, M., Vinayagam, R. and Xu, B. 2020. Guava leaves extract ameliorates STZ induced diabetes mellitus via activation of PI3K/AKT signaling in skeletal muscle of rats. *Mol Biol Rep.* **47** (4):2793-2799. <https://doi.org/10.1007/s11033-020-05399-2>.

Jimenez-Escrig A, Rincon M, Pulido R, Saura-Calixto F. 2001. Guava fruit (*Psidium guajava* L.) as a new source of antioxidant dietary fiber. *J Agric Food Chem.* **49(11)**: 5489-5493.

John, N. R., Gala, V. C. and Sawant, C. S. 2013. Inhibitory effects of plant extracts on multi-species dental biofilm formation in-vitro. *International J. Pharma and Bio Sciences.* **4**:487-495.

Joseph, B. and Priya, M. 2011. Review on nutritional, medicinal and pharmacological properties of Guava (*Psidium guajava* Linn). *Int J Pharm Biol Sci.* **2**: 53-69.

Kapoor, S., Gandhi, N. and Kapoor, A. 2020. Guava (*Psidium guajava*). In: Nayik, G.A., Gull, A. (eds) *Antioxidants in Fruits: Properties and Health Benefits*. Springer, Singapore. https://doi.org/10.1007/978-981-15-7285-2_12

Karou, S. D., Tchacondo, T., Djikpo Tchibozo, M. A., Abdoul Rahaman, S., Anani, K. and Koudouvo, K. et al., 2011. Ethnobotanical study of medicinal plants used in the management of diabetes mellitus and hypertension in the Central Region of Togo. *Pharm Biol.* **49**:1286-1297.

Khan, M. A., Islam, M. K., Siraj, M. A., Saha, S., Barman, A.K. and Awang, K. et al., 2015. Ethnomedicinal survey of various communities residing in Garo Hills of Durgapur, Bangladesh. *J Ethnobiol Ethnomed.* **11**, 44.

Khan, H.B., Shanmugavalli, R., Rajendran, D., Bai, M.R., Sorimuthu, S. 2013. Protective effect of Psidium guajava leaf extract on altered carbohydrate metabolism in streptozotocin-induced diabetic rats. *J. Diet Suppl.* **10** (4): 335-344. <https://doi.org/10.3109/19390211>.

Khedr, S. I., Mokhamer, E., Hassan, A., El-Feki, A. S., Elkhodary, G. M., El-Gerbed, M. 2021. Psidium guajava Linn leaf ethanolic extract: In vivo giardicidal potential with ultrastructural damage, anti-inflammatory and antioxidant effects. *Saudi J. Biol. Sci.* **28** (1): 427-439. <https://doi.org/10.1016/j.sjbs.2020.10.026>.

Kisangau, D. P., Lyaruu, H. V., Hosea, K. M. and Joseph, C. C. 2007. Use of traditional medicines in the management of HIV/AIDS opportunistic infections in Tanzania: A case in the Bukoba rural district. *J Ethnobiol Ethnomed.* **3**, 29.

Koriem, K. M. M., Arbid, M.S. and Saleh, H.N. 2019. Antidiarrheal and protein conservative activities of Psidium guajava in diarrheal rats. *J. Integr. Med.* **17(1)**: 57-65. <https://doi.org/10.1016/j.joim.2018.12.001>.

Kumar, M., Tomar, M., Amarowicz, R., Saurabh, V., Nair, M.S., Maheshwari, C., Sasi, M., Prajapati, U., Hasan, M., Singh, S., Changan, S., Prajapat, R.K., Berwal, M.K., Satankar, V. 2021. Guava (*Psidium guajava* L.) Leaves : Nutritional composition, phytochemical profile, and health-promoting bioactivities. *Foods.* **10** (752): 1-20.

Lans, C., Harper, T., Georges, K. and Bridgewater, E. 2000. Medicinal plants used for dogs in Trinidad and Tobago. *Prev Vet Med.* **45**: 201-220.

Lans, C., Turner, N., Brauer, G., Lourenco, G. and Georges, K. 2006. Ethnoveterinary medicines used for horses in Trinidad and in British Columbia, Canada. *J Ethnobiol Ethnomed.* **2**, 31.

Li, Y., Xu, J., Li, D., Ma, H., Mu, Y., Huang, X., Li, L., 2020. Guavinoside B from Psidium guajava alleviates acetaminophen-induced liver injury via regulating the Nrf2 and JNK signaling pathways. *Food Funct.* **11(9)**: 8297-8308. <https://doi.org/10.1039/d0fo01338b>

Li, Y., Xu, J., Li, D., Ma, H., Mu, Y., Zheng, D., Huang, X., Li, L. 2021. Chemical characterization and hepatoprotective effects of a standardized triterpenoid-enriched guava leaf extract. *J. Agric. Food Chem.* **69** (12): 3626-3637. <https://doi.org/10.1021/acs.jafc.0c07125>.

Lima, L. S., Santos, C. M. R. and Lima, R. A. 2018. Guava tree (*Psidium guajava* L.): Origin, cultivation and characteristics. *Revista Brasileira de Fruticultura.* **40(3)**: e-399. doi: 10.1590/0100-29452018399

Lin, C.Y. and Yin, M. C. 2012. Renal protective effects of extracts

- from guava fruit (*Psidium guajava* L.) in diabetic mice. *Plant Foods Hum Nutr.* **67**: 303-308.
- Lin, J., Puckree, T. and Mvelase, T. P. 2002.** Anti-diarrhoeal evaluation of some medicinal plants used by Zulu traditional healers. *J Ethnopharmacol.* **79**: 53-56.
- Lin, H.C. and Lin, J.Y. 2020.** GSF3, a polysaccharide from guava (*Psidium guajava* L.) seeds, inhibits MCF-7 breast cancer cell growth via increasing Bax/Bcl-2 ratio or Fas mRNA expression levels. *Int. J. Biol. Macromol.* **161**:1261–1271. <https://doi.org/10.1016/j.ijbiomac.2020.06.057>.
- Lingaiah, M. and Rao, P.N. 2013.** An ethnobotanical survey of medicinal plants used by traditional healers of Adilabad district, Andhra Pradesh, India. *Biolife.* **1**: 17-23.
- Livingston Raja, N. R. and Sundar, K. 2006.** *Psidium guajava* Linn confers gastroprotective effects on rats. *Eur Rev Med Pharmacol Sci.* **16**: 151-156.
- Lok, B., Babu, D., Tabana, Y., Dahham, S. S., Adam, M. A. A., Barakat, K. and Sandai, D. 2023.** The Anticancer Potential of *Psidium guajava* (Guava) Extracts. *Life.* **13**: 346. <https://doi.org/10.3390/life13020346>
- Lozoya, X., Reyes-Morales, H., Chavez-Soto, M., MartínezGarcía Mdel, C. and Soto-González, Y. et al., 2002.** Intestinal antispasmodic effect of a phytoextract of *Psidium guajava* folia in the treatment of acute diarrheic disease. *J Ethnopharmacol.* **83**: 19-24.
- Luiza, A. M. and Gustavo, F. 2017.** Larvicidal extract of essential oils from Brazilian cultivars of guava on *Aedes aegypti* L. *Industrial Crops & Products.* **108**: 684-689.
- Luo, Y., Peng, B., Wei, W., Tian, X., Wu, Z. 2019.** Antioxidant and Anti-Diabetic Activities of Polysaccharides from Guava Leaves. *Molecules.* **24**(7): 1343. <https://doi.org/10.3390/molecules24071343>.
- Mandal, A. K., Paudel, S., Pandey, A., Yadav, P., Pathak, P., Grishina, M., Jaremko, M., Emwas, A. H., Khalilullah, H. and Verma, A. 2022.** Guava Leaf Essential Oil as a Potent Antioxidant and Anticancer Agent: Validated through Experimental and Computational Study. *Antioxidants (Basel).* Nov 7;**11**(11):2204. doi: 10.3390/antiox11112204. PMID: 36358576; PMCID: PMC9687059.
- Manosroi, J., Dhumtanom, P. and Manosroi, A. 2006.** Antiproliferative activity of essential oil extracted from Thai medicinal plants on KB and P388 cell lines. *Cancer Letter.* 235.
- Mathias-Mundy, E. and Murdiati, T. B. editors. 1991.** Proceedings of Workshop held at the Central Research Institute for Animal Science; 1990 May 22; Bogor, Indonesia. Indonesia: Indonesian Small Ruminant Network.
- Mesa, M. 2014.** Hypolipidemic potential of plants used in Cuba. *Pharmacol Online.* **1**: 73-80.
- Mesa, A., Mythatha, G.S.S. and Balli, R.** Evaluation of the potential of topically applied salicylic acid-encapsulated chitosan nanoparticles to protect tomato against *Fusarium* wilt. *Iran Polym J* **33**, 671–686 (2024). <https://doi.org/10.1007/s13726-024-01283-z>
- Mittal, P., Gupta, V., Kaur, G., Garg, A. K., Singh, A. 2010.** Phytochemistry and pharmacological activities of *Psidium guajava*: a review. *Int J Pharm Sci Res.* **1**(9):9–19.
- Moein, M. et al. 2015.** Total phenolic, total flavonoid, antioxidant capacity, and antioxidant enzyme activity of guava (*Psidium guajava* L.) leaves. *J. Food Composition and Analysis.* **37**: 47-51.
- Morais-Braga, M. F.B., Sales, D.L., Silva, F.D., Chaves, T.P., Bitu, V.D.N., Avilez, W.M.T., Ribeiro-Filho, J., Coutinho, H.D.M. 2016b.** *Psidium guajava* L. and *Psidium brownianum* Mart ex DC. potentiate the effect of antibiotics against Gram-positive and Gram-negative bacteria. *Eur. J. Integr. Med.* **8**(5):683–687. <https://doi.org/10.1016/j.eujim.2016.07.001>.
- Mukhtar, H. M., Ansari, S. H., Ali, M., Naved, T., Bhat, Z. A. 2004.** Effect of water extract of *Psidium guajava* leaves on alloxan-induced diabetic rats. *Pharmazie.* **59**: 734-735.
- Namsa, N. D., Mandal, M., Tangjang, S. and Mandal, S. C. 2011.** Ethnobotany of the Monpa ethnic group at Arunachal Pradesh, India. *J Ethnobiol Ethnomed.* **7**, 31.
- Naseer, S., Hussain, S., Naeem, N., Pervaiz, M., Rahman, M., 2018.** The phytochemistry and medicinal value of *Psidium guajava* (guava). *Clin. Phytosci.* **4**, 32 <https://doi.org/10.1186/s40816-018-0093-8>.
- Nordini Hasnor, W. I., Fathilah, A. R., Rahim, Z. H. 2013.** Plant extracts of *Psidium guajava*, *Mangifera*, and *Mentha* sp. inhibit the growth of the population of single-species oral biofilm. *Altern Integr Med.* **2**, 102.
- Obarzanek, E., Sacks, F. M., Vollmer, W. M., Bray, G. A., Lin, P.H. and Karanja, N. M. et al., 2001.** Effects on blood lipids of a blood pressure-lowering diet: the Dietary Approaches to Stop Hypertension (DASH) Trial. *Am J Clin Nutr.* **74**(1): 80-89.
- Ogbole, O. O. and Ajaiyeoba EO. 2009.** Traditional management of tuberculosis in Ogun State of Nigeria: The practice and ethnobotanical survey. *Afr J Tradit Complement Altern Med.* **7**:79-84.
- Ojewole, J. A. 2005.** Hypoglycemic and hypotensive effects of *Psidium guajava* Linn. (Myrtaceae) leaf aqueous extract. *Methods Find Exp Clin Pharmacol.* **27**: 689-695.
- Okoli, C. O., Ezike, A. C., Akah, P. A., Udegbunam, S. O., Okoye, T.C., Mbanu, T. P., Ugwu, E. 2009.** Studies on wound healing & antiulcer activities of extract of aerial parts of *Phyllanthus niruri* L.(Euphorbiaceae). *Am J Pharmacol Toxicol.* **4**(4): 118-126.
- Okwu, D. E. and Ekeke, O. 2003.** Phytochemical screening and mineral composition of chewing sticks in South Eastern Nigeria. *Glob J Pure Appl Sci.* **9**: 235.
- Olajide, O., Awe, S. and Makinde, J. 1999.** Pharmacological studies on the leaf of *Psidium guajava*. *Fitoterapia.* **70**: 25-31.
- Olatunji Bello, II., Odusanya, A. J., Raji, I. and Ladipo, C. O. 2007.** Contractile effect of the aqueous extract of *Psidium guajava* leaves on aortic rings in rat. *Fitoterapia.* **78**: 241-243.
- Ong, H. G. and Kim, Y. D. 2014.** Quantitative ethnobotanical study of the medicinal plants used by the Ati Negrito indigenous group in Guimaras Island, Philippines. *J Ethnopharmacol.* **157**: 228-242.
- Párraga, I., López-Torres, J., Andrés, F., Navarro, B. and del Campo J. M. et al., 2011.** Effect of plant sterols on the lipid profile of patients with hypercholesterolaemia. Randomised, experimental study. *BMC Complement Altern Med.* **11**: 73.
- Patel, N. M., Swamy, B. M., Swamy, A. P. and Ravirala, R. 2012.** Evaluation of nephroprotective activity of *Psidium guajava* Linn. leaves extract in paracetamol-induced nephrotoxicity in rats. *Res J Pharm Biol Chem Sci.* **3**: 1247-1256.
- Patel, V., Chatterji, S., Chisholm, D., Ebrahim, S., Gopalakrishna, G. and Mathers, C. et al., 2011.** Chronic diseases and injuries in India. *Lancet.* **377**: 413-428.
- Patel, P., Joshi, C., Birdi, T. and Kothari, V., 2019.** Anti-infective efficacy of *Psidium guajava* L. leaves against certain pathogenic bacteria. *F1000Research.* **8**:12 <https://doi.org/10.12688/f1000research.17500.2>.
- Prabu, G. R., Gnanamani, A. and Sadulla, S. 2006.** Guajaverin, a plant flavonoid as potential antiplaque agent against *Streptococcus mutans*. *J. Applied Microbiology.* **101**: 487-495.
- Pradhan, B. K. and Badola, H. K. 2008.** Ethnomedicinal plant use by Lepcha tribe of Dzongu valley, bordering Khangchendzonga Biosphere Reserve, in North Sikkim, India. *J Ethnobiol Ethnomed.* **4**, 22.
- Rahmat, A., Abu Bakar, M. F., Faezah, N. and Hambali, Z. 2004.** The effects of consumption of guava (*Psidium guajava*) or papaya (*Carica papaya*) on total antioxidant and lipid profile in normal male youth.

Asia Pac J Clin Nutr. **13**(Suppl), S136.

Ravi, K., Divyashree, P. 2014. Psidium guajava: A review on its potential as an adjunct in treating periodontal disease. *Pharmacogn Rev.* **8**: 96-100.

Razak, F. A., Othman, R. Y. and Rahim, Z.H. 2006. The effect of Piper betle and Psidium guajava extracts on the cell-surface hydrophobicity of selected early settlers of dental plaque. *J. Oral Science.* **48**: 71-75.

Remington, P. L., Brownson, R. C. 2011. Centers for Disease Control and Prevention (CDC) Fifty years of progress in chronic disease epidemiology and control. *MMWR Surveill Summ.* **60**: 70-77.

Roy, C. K., Kamath, J. V. and Asad, M. 2006. Hepatoprotective activity of Psidium guajava Linn. leaf extract. *Indian J. Exp Biol.* **44**(4): 305-311.

Ruksiriwanich, W., Khantham, C., Muangsanguan, A., Phimsiripol Y., Barba, F. J., Sringarm, K., Rachtanapun, P., Jantanasakulwong, K., Jantrawut, P., Chittasupho, C., Chutoprapat, R., Boonpisuttinant, K. and Sommano SR. 2022. Guava (*Psidium guajava* L.) Leaf Extract as Bioactive Substances for Anti-Androgen and Antioxidant Activities. *Plants (Basel).* Dec 14. **11**(24): 3514. doi: 10.3390/plants11243514. PMID: 36559626; PMCID: PMC9784754.

Ryu, B., Cho, H.M., Zhang, M., Lee, B.W., Doan, T.P., Park, E.J., Lee, H.J. and Oh, W.K. 2021. Meroterpenoids from the leaves of *Psidium guajava* (guava) cultivated in Korea using MS/MS-based molecular networking. *Phytochemistry.* **186**, <https://doi.org/10.1016/j.phytochem.2021.112723> 112723.

Saber, F. R., Abdelbary, G. A., Salama, M. M., Saleh, D. O., Fathy, M. M., Soliman, F. M., 2018. UPLC/QTOF/MS profiling of two *Psidium* species and the in-vivo hepatoprotective activity of their nano-formulated liposomes. *Food Res. Int.* **105**: 1029–1038. <https://doi.org/10.1016/j.foodres.2017.12.042>.

Sen, S. S., Sukumaran, V., Giri, S. S. and Park, S. C. 2015. Flavonoid fraction of guava leaf extract attenuates lipopolysaccharide-induced inflammatory response via blocking the NF- κ B signaling pathway in *Laboe rohita* macrophages. *Fish Shellfish Immunol.* **47**: 85-92.

Seo, N., Ito, T., Wang, N. L., Ya, X. S., Tokura, Y. and Furukawa, F. et al., 2005. Anti-allergic *Psidium guajava* extracts exert an antitumor effect by inhibition of T regulatory cells and resultant augmentation of Th1 cells. *Anticancer Res.* **25**: 3763-3770.

Shabbir, H., Kausar, T., Noreen, S., Rehman, H.U., Hussain, A., Huang, Q., Gani, A., Su, S. and Nawaz, A. 2020. In Vivo Screening and Antidiabetic Potential of Polyphenol Extracts from Guava Pulp. Seeds and Leaves. *Animals* **10**(9): 1714. <https://doi.org/10.3390/ani10091714>.

Sharma, H. and Chandola, H. M. 2011. Prameha in Ayurveda: correlation with obesity, metabolic syndrome, and diabetes mellitus. Part 1-etiology, classification, and pathogenesis. *J Altern Complement Med.* **17**: 491-496.

Shen, S. C., Cheng, F. C. and Wu, NJ. 2008. Effect of guava (*Psidium guajava* Linn) leaf soluble solids on glucose metabolism in type 2 diabetic rats. *Phototherapy Research.* **22**: 1458-1464.

Shu, Y., Liu, Y., Li, L., Feng, J., Lou, B. and Zhou, X. et al., 2011. Antibacterial activity of quercetin on oral infectious pathogens. *Afr J Microbiol Res.* **5**: 5358-5361.

Silva, E. A. J., Estevam, E. B. B., Silva, T. S., Nicoletta, H. D., Furtado, R. A., Alves, C. C. F., Souchie, E. L., Martins, C. H. G., Tavares, D. C., Barbosa, L.C.A. and Miranda, M. L. D. 2018. Antibacterial and antiproliferative activities of the fresh leaf essential oil of *Psidium guajava* L. (Myrtaceae). *Braz. J. Biol.* **79**(4): 697–702. <https://doi.org/10.1590/1519-6984.189089>.

Singh, R. B., Rastogi, S. S., Singh, N. K., Ghosh, S. and Gupta, S. et al., 1993. Can guava fruit intake decrease blood pressure and blood

lipids. *J. Human Hypertension.* 7:33-38.

Singh, R.B., Rastogi, S.S., Singh, R., Ghosh, S. and Niaz, M. A. 1992. Effects of guava intake on serum total and high-density lipoprotein cholesterol levels and on systemic blood pressure. *Am J Cardiol.* **70**:1287-1291.

Sobral-Souza, C. E., Silva, A. R. P., Leite, N. F., Rocha, J.E., Sousa, A. K., Costa, J.G. M., Menezes, I. R.A., Cunha, F.A.B., Rolim, L.A., Coutinho, H. D. M. 2018. *Psidium guajava* bioactive product chemical analysis and heavy metal toxicity reduction. *Chemosphere.* **216**: 785–793. <https://doi.org/10.1016/j.chemosphere>.

Tan, S., Wang, Z., Xiang, Y., Deng, T., Zhao, X., Shi, S., Zheng, Q., Gao, X., Li, W. 2020. The effects of drying methods on chemical profiles and antioxidant activities of two cultivars of *Psidium guajava* fruits. *L.W.T.* **118**. <https://doi.org/10.1016/j.lwt.2019.108723>.

Tartrakoon, W., Tongmul, S., Tongyen, J., Tartrakoon, T., Vearasilp T. and ter Meulen, U. 2005. Use of fresh and dry guava leaves as a supplement in weaned pig diets. Proceedings of International Research on Food Security, Natural Resource Management and Rural Development.pp.11-13. University of Hohenheim, Stuttgart. <http://www.tropentag.de/2005/proceedings/proceedings.pdf>.

Teixeira, R., Camparoto, M. L., Mantovani, M. S. and Vicentini V. 2003. Assessment of two medicinal plants, *Psidium guajava* L and *Achillea millefolium* L, in in vitro and in vivo assays. *Genet Mol Biol.* **26**: 551-555.

Tella, T., Masola, B. and Mukaratirwa, S. 2019. The effect of *Psidium guajava* aqueous leaf extract on liver glycogen enzymes, hormone sensitive lipase and serum lipid profile in diabetic rats. *Biomed. Pharmacother.* **109**: 2441–2446. <https://doi.org/10.1016/j.biopha.2018.11.137>.

Tetali, P., Waghchaure, C., Daswani, P. G., Antia, N. H. and Birdi T. J. 2009. Ethnobotanical survey of antidiarrhoeal plants of Parinche valley, Pune district, Maharashtra, India. *J Ethnopharmacol.* **123**: 229-236.

Tona, L., Kambu, K., Ngimbi, N., Cimanga, K. and Vlietinck, A. J. 1998. Antiamoebic and phytochemical screening of some Congolese medicinal plants. *J Ethnopharmacol.* **61**: 57-65.

Uduak, U., James, T., Sunday, A. M. and Wilson, O. H. 2012. Ulceroprotective Effect of Methanol Extract of *Psidium guajava* Leaves on Ethanol Induced Gastric Ulcer in Adult Wistar Rats. *Asian J. Medical Sciences.* **4**(2):75-78.

Uuh-Narvaez, J. J., Negrete-Leo´n, E., Acevedo-Ferna´ndez, J. J., and Segura-Campos, M.R. 2021. Antihyperglycemic and hypoglycemic activity of Mayan plant foods in rodent models. *J. Sci. Food Agric.* **101** (10): 4193–4200. <https://doi.org/10.1002/jsfa.11057>.

Vieira, D. R., Amaral, F. M., Maciel, M. C., Nascimento, F. R., Lib´erio, S. A. and Rodrigues, VP. 2014. Plant species used in dental diseases: Ethnopharmacology aspects and antimicrobial activity evaluation. *J Ethnopharmacol.* **155**: 1441-1449.

Vijayakumar, K., Arumugam, V.A., Ramasamy, M., Natesan, M., Palanisamy, S., Thajuddin, N.B., Balasubramanian, B. and Meyyazhagan, A. 2020. Hepatoprotective effects of *Psidium guajava* on mitochondrial enzymes and inflammatory markers in carbon tetrachloride-induced hepatotoxicity. *Drug Dev. Ind. Pharm.* **46** (12): 2041–2050. <https://doi.org/10.1080/03639045.2020.1843474>.

Vijayakumar, K., Rengarajan, R.L., Radhakrishnan, R. and Anand, A. V. 2018. Hypolipidemic Effect of *Psidium guajava* Leaf Extract Against Hepatotoxicity in Rats. *Pharmacogn. Mag.* **53**: 4–8. https://doi.org/10.4103/pm.pm_167_17.

Weli, A., Al-Kaabi, A., Al-Sabahi, J., Said, S., Hossai, A., Al-Riyami, S. 2019. Chemical composition and biological activities of the essential oils of *Psidium guajava* leaf. *J. King Saud Univ. Sci.* 2–24. <https://doi.org/10.1016/j.jksus.2018.07.021>.

Xu, C., Li, X., Zeng, D., Liu, Y., Gao, Y., Tsunoda, M., Deng, S., Xie,

- X., Wang, R., Li, L. S., Song, Y. and Zhang, Y. 2020.** Amino acid profiling study of *Psidium guajava* L. leaves as an effective treatment for type 2 diabetic rats. *eCAM*. 9784382. <https://doi.org/10.1155/2020/9784382>.
- Yadav, V. R., Prasad, S., Sung, B., Kannappan, R. and Aggarwal, B. B. 2010.** Targeting inflammatory pathways by triterpenoids for prevention and treatment of cancer. *Toxins*. **2(1)**: 2428-2466.
- York, T., van Vuuren, S. F. and de Wet, H. 2012.** An antimicrobial evaluation of plants used for the treatment of respiratory infections in rural Maputaland, KwaZulu Natal, South Africa. *J Ethnopharmacol*. **144**: 118-127.
- Zahin, M., Ahmad, I. and Aqil, F. 2017.** Antioxidant and antimutagenic potential of *Psidium guajava* leaf extracts. *Drug Chem. Toxicol*. **40(2)**: 146–153. <https://doi.org/10.1080/01480545.2016.1188397>.
- Zhu, X., Ouyang, W., Lan, Y., Xiao, H., Tang, L., Liu, G., Feng, K., Zhang, L., Song, M. and Cao, Y. 2020.** Anti-hyperglycemic and liver protective effects of flavonoids from *Psidium guajava* L. (guava) leaf in diabetic mice. *Food Biosci*. <https://doi.org/10.1016/j.fbio.2020.100574>.
- Zhu, X., Ouyang, W., Pan, C., Gao, Z., Han, Y., Song, M., Feng, K., Xiao, H., Cao, Y., 2019.** Identification of a new benzophenone from *Psidium guajava* L. leaves and its antineoplastic effects on human colon cancer cells. *Food Funct*. **17**: **10(7)**: 4189-4198. doi: 10.1039/c9fo00569b.

